

9-1-2019

Outcomes of Bilateral Shoulder Arthroplasties: A Comparison of Bilateral Total Shoulder Arthroplasties and Bilateral Reverse Shoulder Arthroplasties.

Ryan Matthew Cox
Thomas Jefferson University

Tyler James Brolin
Thomas Jefferson University

Eric Michael Padegimas
Thomas Jefferson University

Mark D. Lazarus
Additional works at: https://jdc.jefferson.edu/rothman_institute
Thomas Jefferson University
Part of the [Orthopedics Commons](#), and the [Surgery Commons](#)

Charles Lonnie Getz
[Let us know how access to this document benefits you](https://jdc.jefferson.edu/rothman_institute/116)
Thomas Jefferson University

Recommended Citation

See next page for additional authors.
Cox, Ryan Matthew; Brolin, Tyler James; Padegimas, Eric Michael; Lazarus, Mark D.; Getz, Charles Lonnie; Ramsey, Matthew Lee; Williams, Gerald Ross; and Abboud, Joseph Albert, "Outcomes of Bilateral Shoulder Arthroplasties: A Comparison of Bilateral Total Shoulder Arthroplasties and Bilateral Reverse Shoulder Arthroplasties." (2019). *Rothman Institute Faculty Papers*. Paper 116.

https://jdc.jefferson.edu/rothman_institute/116

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Rothman Institute Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Ryan Matthew Cox, Tyler James Brolin, Eric Michael Padegimas, Mark D. Lazarus, Charles Lonnie Getz, Matthew Lee Ramsey, Gerald Ross Williams, and Joseph Albert Abboud



Outcomes of Bilateral Shoulder Arthroplasties: A Comparison of Bilateral Total Shoulder Arthroplasties and Bilateral Reverse Shoulder Arthroplasties

Ryan Matthew Cox, BS, Tyler James Brodin, MD, Eric Michael Padegimas, MD, Mark David Lazarus, MD, Charles Lonnie Getz, MD, Matthew Lee Ramsey, MD, Gerald Ross Williams, Jr., MD, Joseph Albert Abboud, MD

Division of Shoulder and Elbow Surgery, Rothman Institute of Orthopaedics, Thomas Jefferson University Hospitals, Philadelphia, PA, USA

Background: The purpose of this study was to compare outcomes of patients who underwent bilateral total shoulder arthroplasties (TSAs) for osteoarthritis (OA) versus bilateral reverse shoulder arthroplasties (RSAs) for cuff tear arthropathy (CTA).

Methods: Inclusion criteria were patients who underwent bilateral TSAs for OA or bilateral RSAs for CTA with at least 2 years of follow-up. Twenty-six TSA patients (52 shoulders) were matched 2 to 1 with 13 RSA patients (26 shoulders) by sex, age at first surgery, and time between surgeries. Outcomes measured were shoulder range of motion (ROM), complications, and patient-reported scores.

Results: Preoperatively, TSA patients had significantly better forward elevation (FE) of both shoulders than RSA patients (dominant side [Dom]: $103^\circ \pm 32^\circ$ vs. $81^\circ \pm 31^\circ$, $p = 0.047$; nondominant side [non-Dom]: $111^\circ \pm 28^\circ$ vs. $70^\circ \pm 42^\circ$, $p = 0.005$) without significant differences in external (ER) or internal rotation (IR). Postoperatively, TSA patients had significantly better FE (Dom and non-Dom: $156^\circ \pm 12^\circ$, $156^\circ \pm 14^\circ$ vs. $134^\circ \pm 24^\circ$, $137^\circ \pm 23^\circ$; $p = 0.006$, $p = 0.019$) and ER ($42^\circ \pm 11^\circ$, $43^\circ \pm 10^\circ$ vs. $24^\circ \pm 12^\circ$, $25^\circ \pm 10^\circ$; $p < 0.001$, $p < 0.001$) bilaterally and IR of their dominant arm (L1 vs. L4, $p = 0.045$). TSA patients had significantly better activities of daily living external and internal rotations (ADLEIR) scores (Dom and non-Dom: 35.3 ± 1.0 , 35.5 ± 0.9 vs. 32.1 ± 2.4 , 32.5 ± 2.2 ; $p = 0.001$, $p = 0.001$), American Shoulder and Elbow Surgeons scores (94.2 ± 8.4 , 94.2 ± 8.2 vs. 84.7 ± 10.0 , 84.5 ± 8.0 ; $p = 0.015$, $p = 0.004$), and Single Assessment Numerical Evaluation (SANE) scores (93.5 ± 7.6 , 93.8 ± 11.8 vs. 80.5 ± 14.2 , 82.3 ± 13.1 ; $p = 0.014$, $p = 0.025$), with no significant difference in visual analog scale pain scores (0.4 ± 1.0 , 0.3 ± 1.0 vs. 0.7 ± 1.3 , 0.8 ± 1.2) bilaterally.

Conclusions: Overall, patients with bilateral TSAs and RSAs exhibited improved ROM and patient-reported outcomes. Those with bilateral TSAs had better functional outcomes than those with bilateral RSAs.

Keywords: Total shoulder arthroplasty, Total shoulder replacements, Osteoarthritis, Cuff tear arthropathy, Activities of daily living

Anatomic total shoulder arthroplasty (TSA) has proven to be a reliable procedure for a variety of shoulder patholo-

gies with good long-term results.¹⁻⁶⁾ TSA is most commonly performed for degenerative shoulder conditions such as glenohumeral osteoarthritis (OA). Reverse shoulder arthroplasty (RSA) has been used in Europe since the 1980s and was approved by the Food and Drug Administration in 2004.⁷⁾ RSA is a treatment option for patients with cuff tear arthropathy (CTA), which has yielded consistent results. The number of TSA and especially RSA procedures performed each year continues to increase.⁸⁾ The cause of the increase in RSAs performed is multifactorial, but

Received December 23, 2018; Accepted February 25, 2019

Correspondence to: Joseph Albert Abboud, MD

Division of Shoulder and Elbow Surgery, Rothman Institute of Orthopaedics, Thomas Jefferson University Hospitals, 925 Chestnut St, 5th floor, Philadelphia, PA 19107, USA

Tel: +1-267-339-3738, Fax: +1-267-479-1379

E-mail: abboudj@gmail.com

Table 1. Surgical Procedure Details

Patient	Side	Surgical approach	Subscapularis management	Other procedure	Glenoid component
Bilateral TSA					
1	Dom	Deltopectoral	Peel		Standard
	Non-Dom	Deltopectoral	Peel		Standard
2	Dom	Deltopectoral	Peel		Standard
	Non-Dom	Deltopectoral	Peel		Standard
3	Dom	Deltopectoral	Peel		Standard
	Non-Dom	Deltopectoral	Peel		Standard
4	Dom	Deltopectoral	Peel		Standard
	Non-Dom	Deltopectoral	LTO		Standard
5	Dom	Deltopectoral	LTO		Standard
	Non-Dom	Deltopectoral	LTO		Standard
6	Dom	Deltopectoral	LTO		Standard
	Non-Dom	Deltopectoral	LTO		Standard
7	Dom	Deltopectoral	Peel		Standard
	Non-Dom	Deltopectoral	LTO		Standard
8	Dom	Deltopectoral	Peel		Standard
	Non-Dom	Deltopectoral	Peel		Standard
9	Dom	Deltopectoral	LTO		StepTech ^{®*}
	Non-Dom	Deltopectoral	LTO		StepTech [®]
10	Dom	Deltopectoral	LTO		Standard
	Non-Dom	Deltopectoral	LTO		Standard
11	Dom	Deltopectoral	LTO		Standard
	Non-Dom	Deltopectoral	LTO		Standard
12	Dom	Deltopectoral	LTO		Standard
	Non-Dom	Deltopectoral	LTO		Standard
13	Dom	Deltopectoral	Peel		Standard
	Non-Dom	Deltopectoral	Peel		Standard
14	Dom	Deltopectoral	LTO		Standard
	Non-Dom	Deltopectoral	LTO		Mini [†]
15	Dom	Deltopectoral	LTO	Glenoid bone graft	Standard
	Non-Dom	Deltopectoral	Peel		Standard
16	Dom	Deltopectoral	LTO		Standard
	Non-Dom	Deltopectoral	LTO		Standard

Table 1. Continued 1

Patient	Side	Surgical approach	Subscapularis management	Other procedure	Glenoid component
17	Dom	Deltopectoral	Peel		Standard
	Non-Dom	Deltopectoral	Peel		Standard
18	Dom	Deltopectoral	Peel		Standard
	Non-Dom	Deltopectoral	Peel		Standard
19	Dom	Deltopectoral	Peel		Standard
	Non-Dom	Deltopectoral	Peel		Standard
20	Dom	Deltopectoral	LTO		Standard
	Non-Dom	Deltopectoral	LTO		Standard
21	Dom	Deltopectoral	Peel	Rotator cuff repair	Standard
	Non-Dom	Deltopectoral	Peel		Standard
22	Dom	Deltopectoral	LTO		StepTech®
	Non-Dom	Deltopectoral	LTO		StepTech®
23	Dom	Deltopectoral	LTO		Standard
	Non-Dom	Deltopectoral	Peel		Standard
24	Dom	Deltopectoral	LTO		Standard
	Non-Dom	Deltopectoral	LTO		Standard
25	Dom	Deltopectoral	Peel		Standard
	Non-Dom	Deltopectoral	Peel		Standard
26	Dom	Deltopectoral	LTO		Standard
	Non-Dom	Deltopectoral	LTO		Standard
Bilateral RSA					Lateralized vs. Grammont
1	Dom	Deltopectoral	Torn		Lateralized
	Non-Dom	Deltopectoral	Torn		Lateralized
2	Dom	Deltopectoral	LTO	Bone graft; latissimus transfer	Lateralized
	Non-Dom	Deltopectoral	LTO	Bone graft; latissimus transfer	Lateralized
3	Dom	Deltopectoral	Peel		Lateralized
	Non-Dom	Deltopectoral	Peel		Lateralized
4	Dom	Deltoid split		Distal clavicle excision	Grammont
	Non-Dom	Deltopectoral	Peel	Bone graft; distal clavicle excision	Grammont
5	Dom	Deltopectoral	Peel		Lateralized
	Non-Dom	Deltopectoral	LTO	Rotational deltoid plasty	Lateralized
6	Dom	Deltopectoral	Torn		Lateralized
	Non-Dom	Deltopectoral	Peel		Lateralized

Table 1. Continued 2

Patient	Side	Surgical approach	Subscapularis management	Other procedure	Glenoid component
7	Dom	Deltoid split			Grammont
	Non-Dom	Deltoid split			Grammont
8	Dom	Deltoid split			Grammont
	Non-Dom	Deltoid split			Grammont
9	Dom	Deltoid split			Grammont
	Non-Dom	Deltoid split			Grammont
10	Dom	Deltoid split			Grammont
	Non-Dom	Deltopectoral	LTO		Grammont
11	Dom	Deltopectoral	Peel	Bone graft	Grammont
	Non-Dom	Deltopectoral	Torn		Grammont
12	Dom	Deltopectoral	Peel		Lateralized
	Non-Dom	Deltopectoral	Peel		Lateralized
13	Dom	Deltoid split			Grammont
	Non-Dom	Deltoid split			Lateralized

TSA: total shoulder arthroplasty, Dom: dominant side, Non-Dom: nondominant side, RSA: reverse shoulder arthroplasty, LTO: lesser tuberosity osteotomy. *StepTech®: Global® StepTech® Anchor Peg Glenoid (DePuy Synthes, Warsaw, IN, USA). †Mini: ArthroSurface Inlay Glenoid Replacement (ArthroSurface, Franklin, MA, USA).

includes increased surgeon's confidence in the procedure, as well as expanding indications such as use in revision shoulder arthroplasty, posterior glenoid bone loss, and fracture sequelae.^{9,10)}

Given the increase in utilization, prevalence of shoulder OA and rotator cuff disease, increasing life span of patients, and expanding indications, it is common for patients to require staged bilateral shoulder arthroplasties. Historically, one concern of bilateral RSAs has been the impact on the performance of activities of daily living (ADLs), especially maintenance of perineal hygiene. However, recent evidence suggests that patients with bilateral RSAs are able to adequately perform these activities.¹¹⁻¹⁴⁾ Several other studies have reported the results of bilateral TSAs and RSAs independently.¹⁵⁻¹⁹⁾ However, no studies have directly compared the outcomes of patients with bilateral TSAs to bilateral RSAs.

The purpose of this study was to compare the clinical and functional outcomes of patients who underwent bilateral TSAs for OA to patients who underwent bilateral RSAs for CTA. We hypothesized that patients who underwent bilateral TSAs would have better clinical and functional outcomes than patients who underwent bilateral RSAs.

METHODS

Institutional Review Board approval was obtained from Thomas Jefferson University Hospital prior to conducting this retrospective analysis (IRB No. 45 CFR 46.110, Control #17D.020). All data collection was retrospective and all patients provided informed consent preoperatively for retrospective data review. Our institutional billing database was queried using the CPT code 23472 (arthroplasty, glenohumeral joint; total shoulder) to identify all patients who underwent a TSA or RSA from 2004 to 2015. A retrospective chart review was performed to identify all patients who underwent staged bilateral TSAs for OA or staged bilateral RSAs for CTA. Inclusion criteria were patients who underwent bilateral TSAs for OA or bilateral RSAs for CTA with at least 2 years of clinical follow-up from the latest arthroplasty surgery. Exclusion criteria were patients (1) undergoing bilateral TSAs for any diagnosis other than OA, (2) undergoing RSAs for any diagnosis other than CTA, (3) undergoing revision arthroplasty surgery, and (4) having chronic neurologic conditions (cerebrovascular accident, multiple sclerosis, and Parkinson disease). Rotator cuff integrity was assessed intraoperatively for all

patients undergoing a TSA procedure. If their rotator cuff was determined to be insufficient, they received an RSA and were thus excluded from the study. For primary OA, all patients had Smilson-Prieto grade 2 or greater changes; for primary CTA, all patients had Hamada grade 2 or greater changes.^{20,21)} From 2004 to 2015, 5,261 primary shoulder arthroplasties were performed at our institution. After retrospective chart review, 26 TSA patients (52 shoulders) were matched 2 to 1 with 13 RSA patients (26 shoulders) according to sex, age at first surgery, and time from first arthroplasty surgery.

Preoperative and postoperative active ranges of motion (ROMs) in forward elevation (FE), external rotation (ER), and internal rotation (IR) were assessed by chart reviews of clinician-documented ROM. Postoperative complications, readmissions, and reoperations were recorded. Identified patients were then assessed at follow-up visits or contacted via phone for activities of daily living external and internal rotations (ADLEIR; 0–36 point scale of ADL performance) scores, American Shoulder and Elbow Surgeons (ASES) shoulder scores, Single Assessment Numerical Evaluation (SANE), and visual analog scale (VAS) pain and VAS patient satisfaction scores of both shoulders.^{22–25)} All procedures were performed by one of five fellowship-trained shoulder and elbow surgeons (CLG, MDL, MLR, GRW, JAA). All surgeries were performed either through a standard deltopectoral approach or through a superolateral deltoid-splitting approach if prior surgical incisions required this approach.²⁶⁾ Full details of the surgical procedures can be found in Table 1. All patients were immobilized in a sling for 2–4 weeks postoperatively and then were enrolled in a standardized postoperative physical therapy protocol. The main difference in rehabilitation

protocols was that patients undergoing TSA begin passive FE and passive ER starting the day of surgery while RSA patients typically do not. All patients begin active assisted ROM at 6 weeks and then begin strengthening at 12 weeks.

Study population characteristics are reported by using measures of central tendency (mean) and variability (standard deviation). Patients' shoulders were compared by using independent samples *t*-tests for continuous variables and chi-square analyses for categorical variables to detect differences between groups. For all statistical tests, a *p*-value < 0.05 was used to determine significance. All statistics were calculated with Microsoft Excel (2013; Microsoft, Redmond, WA, USA).

RESULTS

Patient Demographics and Clinical Characteristics

A summary of patient demographics and clinical characteristics is shown in Table 2. After application of the matching criteria, there were no significant differences in age at first arthroplasty surgery (71.2 ± 5.5 vs. 73.4 ± 5.8 years, $p = 0.507$), percentage of male or female patients ($p = 1.00$), or body mass index (28.9 ± 7.7 vs. 28.3 ± 6.0 kg/m², $p = 0.821$) between bilateral TSA and bilateral RSA cohorts. There were also no significant differences in time between arthroplasties (20.4 ± 18.0 vs. 18.6 ± 18.2 months, $p = 0.772$), follow-up from first arthroplasty (86.8 ± 19.4 vs. 84.1 ± 22.3 months, $p = 0.743$), or follow-up from second arthroplasty (63.4 ± 15.8 vs. 62.7 ± 28.1 months, $p = 0.947$) between these two cohorts. There were no intraoperative complications.

Table 2. Patient Demographics and Clinical Characteristics

Variable	Bilateral TSA	Bilateral RSA	<i>p</i> -value
Age at first arthroplasty (yr)	71.2 ± 5.5	73.4 ± 5.8	0.507
Sex			1.00
Male	6 (23.1)	3 (23.1)	
Female	20 (76.9)	10 (76.9)	
BMI (kg/m ²)	28.9 ± 7.7	28.3 ± 6.0	0.821
Time between arthroplasties (mo)	20.4 ± 18.0	18.6 ± 18.2	0.772
Follow-up of first arthroplasty (mo)	86.8 ± 19.4	84.1 ± 22.3	0.743
Follow-up of second arthroplasty (mo)	63.4 ± 15.8	62.7 ± 28.1	0.947

Values are presented as mean ± standard deviation or number (%).
TSA: total shoulder arthroplasty, RSA: reverse shoulder arthroplasty, BMI: body mass index.

Shoulder ROM

Shoulder ROM was compared between dominant (Dom) and nondominant (non-Dom) sides of the shoulder instead of right versus left to avoid the confounding effect of hand dominance (Table 3). Patients who underwent bilateral TSA had significantly better preoperative FE of their Dom ($103^\circ \pm 32^\circ$ vs. $81^\circ \pm 31^\circ$, $p = 0.047$) and non-Dom ($111^\circ \pm 28^\circ$ vs. $70^\circ \pm 42^\circ$, $p = 0.005$) arms compared to patients who underwent bilateral RSAs. There were no significant differences in preoperative ER between Dom ($24^\circ \pm 15^\circ$ vs. $24^\circ \pm 29^\circ$, $p = 0.983$) and non-Dom ($28^\circ \pm 15^\circ$ vs. $27^\circ \pm 18^\circ$, $p = 0.870$) arms or preoperative IR between Dom (L5 vs. sacrum, $p = 0.684$) and non-Dom (L5 vs. L5, $p = 0.600$) arms in patients who underwent bilateral TSAs or RSAs.

In regards to postoperative ROM, patients who underwent bilateral TSAs had significantly better FE of their Dom ($156^\circ \pm 12^\circ$ vs. $134^\circ \pm 24^\circ$, $p = 0.006$) and non-Dom ($156^\circ \pm 14^\circ$ vs. $137^\circ \pm 23^\circ$, $p = 0.019$) arms, ER of their Dom ($42^\circ \pm 11^\circ$ vs. $24^\circ \pm 12^\circ$, $p < 0.001$) and non-Dom ($43^\circ \pm 10^\circ$ vs. $25^\circ \pm 10^\circ$, $p < 0.001$) arms, and IR of their Dom arm (L1 vs. L4, $p = 0.045$) compared to patients who underwent bilateral RSAs. There was no significant difference in IR of their non-Dom arms (L1 vs. L2, $p = 0.177$).

Patients who underwent bilateral TSAs had significantly better overall improvement in ER of their Dom ($17^\circ \pm 13^\circ$ vs. $-5^\circ \pm 29^\circ$, $p = 0.032$) and non-Dom ($19^\circ \pm$

13° vs. $-6^\circ \pm 18^\circ$, $p = 0.002$) arms compared to patients who underwent bilateral RSAs. There were no significant differences in overall ROM improvement between bilateral TSAs and bilateral RSAs in FE of their Dom ($50^\circ \pm 36^\circ$ vs. $53^\circ \pm 43^\circ$, $p = 1.000$) or non-Dom ($51^\circ \pm 30^\circ$ vs. $72^\circ \pm 39^\circ$, $p = 0.131$) arms or IR of their Dom (3 ± 3 vertebral levels vs. 1 ± 4 vertebral levels, $p = 0.206$) or non-Dom (4 ± 3 vertebral levels vs. 2 ± 5 vertebral levels, $p = 0.22$) arms.

Functional Outcomes Scores

Functional outcome scores were also compared according to hand dominance to avoid any confounding effect. Patients who underwent bilateral TSAs had significantly better ADLEIR scores of their Dom (35.3 ± 1.0 vs. 32.1 ± 2.4 , $p = 0.001$) and non-Dom (35.5 ± 0.9 vs. 32.5 ± 2.2 , $p = 0.001$) arms, ASES scores of their Dom (94.2 ± 8.4 vs. 84.7 ± 10.0 , $p = 0.015$) and non-Dom (94.2 ± 8.2 vs. 84.5 ± 8.0 , $p = 0.004$) arms, and SANE scores of their Dom (93.5 ± 7.6 vs. 80.5 ± 14.2 , $p = 0.014$) and non-Dom (93.8 ± 11.8 vs. 82.3 ± 13.1 , $p = 0.025$) arms, and were more satisfied with their non-Dom arms (9.7 ± 0.8 vs. 8.6 ± 1.1 , $p = 0.012$) compared to bilateral RSA patients (Table 4). There were no significant differences in satisfaction of their Dom arms (9.7 ± 0.9 vs. 8.5 ± 1.8 , $p = 0.057$) or VAS pain scores of their Dom (0.4 ± 1.0 vs. 0.7 ± 1.3 , $p = 0.461$) and non-Dom (0.3 ± 1.0 vs. 0.8 ± 1.2 , $p = 0.253$) arms.

Table 3. Shoulder Range of Motion

Variable	Dom TSA	Dom RSA	<i>p</i> -value	Non-Dom TSA	Non-Dom RSA	<i>p</i> -value
Preoperative						
FE (°)	103 ± 32	81 ± 31	0.047	111 ± 28	70 ± 42	0.005
ER (°)	24 ± 15	24 ± 29	0.983	28 ± 15	27 ± 18	0.870
IR (vertebral level)	L5	Sacrum	0.684	L5	L5	0.600
Postoperative						
FE (°)	156 ± 12	134 ± 24	0.006	156 ± 14	137 ± 23	0.019
ER (°)	42 ± 11	24 ± 12	< 0.001	43 ± 10	25 ± 10	< 0.001
IR (vertebral level)	L1	L4	0.045	L1	L2	0.177
Change in ROM						
FE (°)	50 ± 36	53 ± 43	1.000	51 ± 30	72 ± 39	0.131
ER (°)	17 ± 13	-5 ± 29	0.032	19 ± 13	-6 ± 18	0.002
IR (vertebral level)	3 ± 3	1 ± 4	0.206	4 ± 3	2 ± 5	0.220

Values are presented as mean ± standard deviation.

Dom: dominant side, Non-Dom: nondominant side, TSA: total shoulder arthroplasty, RSA: reverse shoulder arthroplasty, FE: forward elevation, ER: external rotation, IR: internal rotation.

Table 4. Functional Outcomes Scores

Outcome score	Dom TSA	Dom RSA	<i>p</i> -value	Non-Dom TSA	Non-Dom RSA	<i>p</i> -value
ADLEIR	35.3 ± 1.0	32.1 ± 2.4	0.001	35.5 ± 0.9	32.5 ± 2.2	0.001
ASES shoulder score	94.2 ± 8.4	84.7 ± 10.0	0.015	94.2 ± 8.2	84.5 ± 8.0	0.004
SANE	93.5 ± 7.6	80.5 ± 14.2	0.014	93.8 ± 11.8	82.3 ± 13.1	0.025
Satisfaction	9.7 ± 0.9	8.5 ± 1.8	0.057	9.7 ± 0.8	8.6 ± 1.1	0.012
Pain	0.4 ± 1.0	0.7 ± 1.3	0.461	0.3 ± 1.0	0.8 ± 1.2	0.253

Values are presented as mean ± standard deviation.

Dom: dominant side, Non-Dom: nondominant side, TSA: total shoulder arthroplasty, RSA: reverse shoulder arthroplasty, ADLEIR: activities of daily living external and internal rotation, ASES: American Shoulder and Elbow Surgeons, SANE: Single Assessment Numerical Evaluation.

Complications and Reoperations

There were no intraoperative complications for patients in either cohort. In the TSA cohort, 1 patient had a postoperative hematoma that required an irrigation and debridement 6 weeks postoperatively, 1 patient underwent an arthroscopic distal clavicle excision 2 years postoperatively for acromioclavicular joint arthritis, and 1 patient underwent an open reduction and internal fixation (ORIF) for a clavicle fracture sustained over 3 years postoperatively. In the RSA cohort, 1 patient sustained an acromioclavicular joint separation 4 years postoperatively that was treated nonoperatively and 1 patient sustained a periprosthetic fracture 2 years postoperatively that was treated with ORIF. There were no dislocations, infections, or nerve injuries in either cohort.

DISCUSSION

This retrospective cohort comparison shows that in general patients who received bilateral TSAs had better postoperative ROM and patient-reported functional outcomes than patients who received bilateral RSAs. Patients with bilateral TSAs had better postoperative ROM in every measured plane except IR of their non-Dom arms. However, the only plane in which bilateral TSA patients had significantly better improvement in ROM was ER. Bilateral TSA patients also had significantly better functional outcome scores (ASES, SANE, and ADLEIR) than did their bilateral RSA counterparts. Despite the superior functional outcome scores in the bilateral TSA cohort, these two groups had equal final satisfaction for their Dom arms and equal pain relief (VAS pain score) in both arms.

Surgeons remain concerned with the ability of patients to perform all of their ADLs after RSA given the inherent limitations in IR with this arthroplasty design. However, recent evidence suggests that these deficits may

not be as profound as previously thought. Stevens et al retrospectively reviewed the outcomes of 15 consecutive patients who underwent staged bilateral RSA for CTA and found that all patients were able to adequately perform ADLs that require IR including maintaining perineal hygiene.¹³ Wirth et al.¹⁴ reported the results of 57 staged bilateral RSAs at least 1 year from their second arthroplasty surgery and found that at 1 year postoperatively 21% of patients had insufficient IR after their first surgery and 33% had insufficient IR after their second surgery. However, at 2 years postoperatively, only 5% of patients lacked sufficient IR bilaterally. At a mean follow-up of 48.4 months, Levy et al.¹¹ reported the results of 19 patients who underwent staged bilateral RSAs and all patients had no limitations in their leisure activities or ADLs including maintaining perineal hygiene. In the largest cohort study published to date, Mellano et al.¹² retrospectively reviewed 50 patients who underwent staged bilateral RSA for CTA, OA, massive rotator cuff tear, or revision shoulder arthroplasty and found that all patients remained independent with ADLs and personal hygiene 2 years postoperatively. All of these studies clearly demonstrate that patients can remain functional with bilateral RSAs. Similarly, all of the patients in our bilateral RSA cohort reported that they were able to adequately maintain personal hygiene. While the bilateral TSA cohort had superior functional outcomes and superior ADL scores (over 35/36 in ADLEIR testing for both arms), the bilateral RSA cohort still achieved reasonable ADL scores (over 32/36 in both arms).

Certainly, this study has several weaknesses. First, this is a purely retrospective study and is therefore subject to the limitations of retrospective data specifically on the inability to elucidate causal relationships rather than simply correlative ones. The ROM measurements were obtained by retrospectively reviewing the physical examinations dictated by physicians. These measurements

would be more accurate if they were assessed using a goniometer. We did not have preoperative patient-reported functional outcome scores or ADLEIR scores to compare with the postoperative scores and therefore could not assess net improvement in functional outcomes between cohorts. There were also some variations in surgical details and specific TSA and RSA systems patients received may have influenced the results. However, all procedures were performed by high-volume fellowship-trained shoulder and elbow surgeons, so we do not believe these variations would significantly affect the results. Additionally, this sample size may be underpowered to show a significant difference between cohorts. This is an inherent limitation of a retrospective analysis and cannot be modified. Despite these weaknesses, we were able to achieve a direct matched comparison of bilateral TSA to RSA. We found superior functional outcomes in the bilateral TSA cohort,

equal pain relief in both cohorts, and a satisfactory ability to perform ADLs in both cohorts.

Overall, patients who underwent bilateral TSAs for OA and bilateral RSAs for CTA exhibited improved ROM and patient-reported functional outcomes. Patients who underwent bilateral TSAs had superior functional results to patients who underwent bilateral RSAs. Both groups exhibited equal pain relief. Despite the difference in functional results, patients with bilateral RSAs were satisfied with their results and able to perform their ADLs with minimal limitations.

CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

REFERENCES

- Cil A, Veillette CJ, Sanchez-Sotelo J, Sperling JW, Schleck CD, Cofield RH. Survivorship of the humeral component in shoulder arthroplasty. *J Shoulder Elbow Surg.* 2010;19(1):143-50.
- Deshmukh AV, Koris M, Zurakowski D, Thornhill TS. Total shoulder arthroplasty: long-term survivorship, functional outcome, and quality of life. *J Shoulder Elbow Surg.* 2005;14(5):471-9.
- Khan A, Bunker TD, Kitson JB. Clinical and radiological follow-up of the Aequalis third-generation cemented total shoulder replacement: a minimum ten-year study. *J Bone Joint Surg Br.* 2009;91(12):1594-600.
- Raiss P, Bruckner T, Rickert M, Walch G. Longitudinal observational study of total shoulder replacements with cement: fifteen to twenty-year follow-up. *J Bone Joint Surg Am.* 2014;96(3):198-205.
- Raiss P, Schmitt M, Bruckner T, et al. Results of cemented total shoulder replacement with a minimum follow-up of ten years. *J Bone Joint Surg Am.* 2012;94(23):e1711-10.
- Schoch B, Schleck C, Cofield RH, Sperling JW. Shoulder arthroplasty in patients younger than 50 years: minimum 20-year follow-up. *J Shoulder Elbow Surg.* 2015;24(5):705-10.
- Boileau P, Watkinson DJ, Hatzidakis AM, Balg F. Grammont reverse prosthesis: design, rationale, and biomechanics. *J Shoulder Elbow Surg.* 2005;14(1 Suppl S):147S-161S.
- Padegimas EM, Maltenfort M, Lazarus MD, Ramsey ML, Williams GR, Namdari S. Future patient demand for shoulder arthroplasty by younger patients: national projections. *Clin Orthop Relat Res.* 2015;473(6):1860-7.
- Boileau P, Watkinson D, Hatzidakis AM, Hovorka I. Neer Award 2005: The Grammont reverse shoulder prosthesis: results in cuff tear arthritis, fracture sequelae, and revision arthroplasty. *J Shoulder Elbow Surg.* 2006;15(5):527-40.
- Smith CD, Guyver P, Bunker TD. Indications for reverse shoulder replacement: a systematic review. *J Bone Joint Surg Br.* 2012;94(5):577-83.
- Levy O, Walecka J, Arealis G, et al. Bilateral reverse total shoulder arthroplasty-functional outcome and activities of daily living. *J Shoulder Elbow Surg.* 2017;26(4):e85-96.
- Mellano CR, Kupfer N, Thorsness R, et al. Functional results of bilateral reverse total shoulder arthroplasty. *J Shoulder Elbow Surg.* 2017;26(6):990-6.
- Stevens CG, Struk AM, Wright TW. The functional impact of bilateral reverse total shoulder arthroplasty. *J Shoulder Elbow Surg.* 2014;23(9):1341-8.
- Wirth B, Kolling C, Schwyzer HK, Flury M, Audige L. Risk of insufficient internal rotation after bilateral reverse shoulder arthroplasty: clinical and patient-reported outcome in 57 patients. *J Shoulder Elbow Surg.* 2016;25(7):1146-54.
- Fabricant PD, Chin CS, Grawe BM, Dines JS, Craig EV, Dines DM. Staged bilateral total shoulder arthroplasty: improved outcomes with less than 6 months between surgeries. *J Shoulder Elbow Surg.* 2016;25(11):1774-9.
- Gerber C, Lingenfelter EJ, Reischl N, Sukthankar A. Single-stage bilateral total shoulder arthroplasty: a preliminary study. *J Bone Joint Surg Br.* 2006;88(6):751-5.

17. Gruson KI, Pillai G, Vanadurongwan B, Parsons BO, Flattow EL. Early clinical results following staged bilateral primary total shoulder arthroplasty. *J Shoulder Elbow Surg.* 2010;19(1):137-42.
18. Morris BJ, Haigler RE, O'Connor DP, Elkousy HA, Gartsman GM, Edwards TB. Outcomes of staged bilateral reverse shoulder arthroplasties for rotator cuff tear arthropathy. *J Shoulder Elbow Surg.* 2015;24(3):474-81.
19. Wiater BP, Boone CR, Koueiter DM, Wiater JM. Early outcomes of staged bilateral reverse total shoulder arthroplasty: a case-control study. *Bone Joint J.* 2013;95(9):1232-8.
20. Samilson RL, Prieto V. Dislocation arthropathy of the shoulder. *J Bone Joint Surg Am.* 1983;65(4):456-60.
21. Hamada K, Yamanaka K, Uchiyama Y, Mikasa T, Mikasa M. A radiographic classification of massive rotator cuff tear arthritis. *Clin Orthop Relat Res.* 2011;469(9):2452-60.
22. Bodian CA, Freedman G, Hossain S, Eisenkraft JB, Beilin Y. The visual analog scale for pain: clinical significance in postoperative patients. *Anesthesiology.* 2001;95(6):1356-61.
23. Boileau P, Chuinard C, Roussanne Y, Bicknell RT, Rochet N, Trojani C. Reverse shoulder arthroplasty combined with a modified latissimus dorsi and teres major tendon transfer for shoulder pseudoparalysis associated with dropping arm. *Clin Orthop Relat Res.* 2008;466(3):584-93.
24. Michener LA, McClure PW, Sennett BJ. American Shoulder and Elbow Surgeons Standardized Shoulder Assessment Form, patient self-report section: reliability, validity, and responsiveness. *J Shoulder Elbow Surg.* 2002;11(6):587-94.
25. Williams GN, Gangel TJ, Arciero RA, Uhorchak JM, Taylor DC. Comparison of the Single Assessment Numeric Evaluation method and two shoulder rating scales: outcomes measures after shoulder surgery. *Am J Sports Med.* 1999;27(2):214-21.
26. Zlotolow DA, Catalano LW 3rd, Barron OA, Glickel SZ. Surgical exposures of the humerus. *J Am Acad Orthop Surg.* 2006;14(13):754-65.