

7-1-2020

Improving nutrition and immunity with dry chain and integrated pest management food technologies in LMICs.

Peetambar Dahal
University of California, davis

Meghnath Dhimal
Nepal Health Research Council

Krishna Belbase
Nutrition Evaluation Program

Sundar Tiwari
Agricultural and Forestry University

Follow this and additional works at: <https://jdc.jefferson.edu/medoncfp>
John Groopman

 Part of the [University of California, Davis Nutrition Commons](#), and the [Oncology Commons](#)

[Let us know how access to this document benefits you](#)

See next page for additional authors

Recommended Citation

Dahal, Peetambar; Dhimal, Meghnath; Belbase, Krishna; Tiwari, Sundar; Groopman, John; West, Keith; Pollock, Bradley; Pyakurel, Subodh; Acharya, Ganga; Aryal, Sunil; Ghimire, Yug Nath; Neupane, Manish; Poudel, Rosina; Van Ashbrouck, Johan; Kunusoth, Keshavulu; De Saeger, Sarah; De Boevre, Marthe; Gharti-Chhetri, Gokarna; Gurung, Tek Bahadur; and Bradford, Kent J, "Improving nutrition and immunity with dry chain and integrated pest management food technologies in LMICs." (2020). *Department of Medical Oncology Faculty Papers*. Paper 119.
<https://jdc.jefferson.edu/medoncfp/119>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Medical Oncology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Peetambar Dahal, Meghnath Dhimal, Krishna Belbase, Sundar Tiwari, John Groopman, Keith West, Bradley Pollock, Subodh Pyakurel, Ganga Acharya, Sunil Aryal, Yug Nath Ghimire, Manish Neupane, Rosina Poudel, Johan Van Ashbrouck, Keshavulu Kunusoth, Sarah De Saeger, Marthe De Boevre, Gokarna Gharti-Chhetri, Tek Bahadur Gurung, and Kent J Bradford

Improving nutrition and immunity with dry chain and integrated pest management food technologies in LMICs



Preventive public health measures to cope with non-communicable and infectious diseases include improving immunity through better nutrition. Immune dysfunction is regarded as both a cause and a consequence of malnutrition. Malnutrition is a daily killer in low-income and middle income countries (LMICs), and evidence suggests that susceptibility to and severity of infections increases with malnutrition, leading to illness and death. Undernutrition interacts with repeated bouts of infectious disease, causing an estimated 3.5 million preventable maternal and child deaths annually.¹ Recognising the central role of nutrition in improving immunity and reducing morbidity and mortality, the UN is implementing its Scaling Up Nutrition initiative in 61 LMICs. Furthermore, 12 of 17 UN Sustainable Development Goals are related to nutrition. Nutrition needs have been particularly highlighted by the current economic and food supply chain disruptions due to the COVID-19 pandemic.

High-income countries have used health-focused approaches, such as micronutrient fortification, to improve nutrition, and food safety and quality are primary concerns. Although nutrition efforts are underway in LMICs, antinutrients and toxins that compromise human health are widely prevalent in food systems. Thus, many people face silent health crises while consuming both high-moisture and low-moisture food products, impeding efforts to alleviate malnutrition. For example, natural carcinogenic fungal toxins (mycotoxins, especially aflatoxins) and insect infestations develop when low moisture foods or feeds are stored at high relative humidity.^{2,3} Aflatoxins are also transferred to meat and dairy products through feeds, thus affecting human health. Alarming levels of the most potent carcinogenic aflatoxin marker were detected in blood samples of pregnant Nepalese women in 2014 and 2019.⁴

Mycotoxin management is lacking in LMICs because flooding and rainfall damage traditionally stored dry products annually and aggravate nutrition security. Mould mitigation strategies include using inputs such as tolerant or resistant cultivars, combined with good agricultural practices and biological control.⁵ Studies

have identified inhibitors of toxigenic storage moulds in feeds, and both moulds and insects did not proliferate in corn stored in Bill & Melinda Gates Foundation-promoted hermetic bags. Nutrients were maintained for 4 years in sundried and pesticide-treated grains.⁶ Long-term breeding strategies to combine mould resistance with other traits are being pursued. Meanwhile, the dry chain (drying products using natural or artificial methods to safe levels followed by moisture-proof storage) could be implemented immediately to minimise mould and insect infestations and nutrient losses.² However, such straightforward approaches to preserve food and feed quality have yet to be promoted.⁷⁻⁹ To minimise food chain disruption by disasters, including COVID-19, it is crucial for LMICs to safely preserve staple foods, and global organisations, such as the UN World Food Program and the Bill & Melinda Gates Foundation, could assist with implementation of dry chain systems.

Additionally, artificial toxins are ingested every day by people in LMICs because of improper pesticide use in high-moisture fruits and vegetables. Although the health sector is also addressing pesticide residues, a standoff occurred at the Nepal-India border due to fears of excess pesticides in imported foods. Genetically engineered crops provide options to reduce pesticide residues.¹⁰ Implementing integrated pest management by cultural, mechanical, biological, and physical methods, as well as soft chemicals, could minimise economic, health, and environmental risks to farmers and consumers.

Multidisciplinary efforts are needed to minimise food toxins by implementation of interventions like integrated pest management and cold and dry chains, followed by sensitive toxin monitoring of domestic and imported food and feed products. High-moisture foods need cold chain (refrigerated) protection to minimise nutrient losses and wastage. Implementing analogous pesticide-free dry chain protective measures for low-moisture products, particularly seeds and grains, will additionally enable food reserves for disasters such as COVID-19, improve trade ratios, and complement nutritional security, even during normal periods. Although LMICs often put emphasis on the economics of

For more on implementing integrated pest management see <https://ipmil.cired.vt.edu/>

food production rather than on its quality, the measures that have been undertaken to limit the COVID-19 pandemic and associated global economic stress exemplify that human health must be above economic concerns. Thus, LMICs should prioritise minimisation of food toxins to conform to FAO/WHO Codex standards and protect human health. Implementation of a combination of toxin mitigation strategies followed by sensitive monitoring procedures will improve food security, reduce malnutrition, enhance immunity, and minimise the effect of both non-communicable and infectious diseases. This strategy would provide a long-lasting legacy to minimise future deaths caused by malnutrition and be a tribute to human lives lost during the COVID-19 pandemic.

We declare no competing interests.

Copyright © 2020 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

**Peetambar Dahal, Meghnath Dhimal, Krishna Belbase, Sundar Tiwari, John Groopman, Keith West, Bradley Pollock, Subodh Pyakurel, Ganga Acharya, Sunil Aryal, Yug Nath Ghimire, Manish Neupane, Rosina Poudel, Johan Van Ashbrouck, Keshavulu Kunusoth, Sarah De Saeger, Marthe De Boevre, Gokarna Gharti-Chhetri, Tek Bahadur Gurung, Kent J Bradford*
peetambardahal@gmail.com

Department of Plant Sciences (PD, KJB) and Department of Public Health Sciences (BP), University of California, Davis, CA 95616, USA; Nepal Health Research Council, Ministry of Health, Kathmandu, Nepal (MD); Nutrition Evaluation Program, UNICEF, New York, NY, USA (KB); Division of Entomology, Agricultural and Forestry University, Bharatpur, Nepal (ST); Department of Environmental Health and Engineering (JG) and Department of International

Health (KW), Johns Hopkins University, Baltimore, MD, USA; Planning Commission, Province 1 Government, Biratnagar, Nepal (SP); Ministry of Agriculture and Livestock Development, Government of Nepal, Kathmandu, Nepal (GA); Division of Entomology (SA) and Division of Agricultural Economics (YNG), Nepal Agricultural Research Council, Kathmandu, Nepal; Department of Medical Oncology, Thomas Jefferson University, Pennsylvania, PA, USA (MN); Department of Internal Medicine, Leonard J Chabert Medical Center, Houma, LA, USA (RP); Rhino Research, Bangkok, Thailand (JVA); Telangana State Seed and Organic Certification Authority, Government of Telangana, Hyderabad, India (KK); Department of Bioanalysis, Ghent University, Belgium (SDS, MDB); Division of Natural Sciences, Palm Beach State College, Lake Worth, FL, USA (GG-C); and Nepal Fisheries Society, Kathmandu, Nepal (TBG)

- 1 Development Initiatives. 2018 Global Nutrition Report: shining a light to spur action on nutrition. Bristol, UK: Development Initiatives.
- 2 Bradford KJ, Dahal P, Van Asbrouck J, et al. The dry chain: reducing postharvest losses and improving food safety in humid climates. *Trends Food Sci Technol* 2018; **71**: 84–93.
- 3 Mahato DK, Lee KE, Kamle M, et al. Aflatoxins in food and feed: an overview on prevalence, detection and control strategies. *Front Microbiol* 2019; **10**: 2266.
- 4 Andrews-Trevino JY, Webb P, Shively G, et al. Dietary determinants of aflatoxin B 1-lysine adduct in pregnant women consuming a rice-dominated diet in Nepal. *Eur J Clin Nutr* 2020; **74**: 732–40.
- 5 Agbetameh D, Ortega-Beltran A, Awuah RT, et al. Potential of atoxigenic *Aspergillus flavus* vegetative compatibility groups associated with maize and groundnut in Ghana as biocontrol agents for aflatoxin management. *Front Microbiol* 2019; **10**: 2069.
- 6 Yin D, Yuan J, Guo Y, Chiba LI. Effect of storage time on the characteristics of corn and efficiency of its utilization in broiler chickens. *Anim Nutr* 2017; **3**: 252–57.
- 7 Clark H, Awa M, Coll S, et al. A future for the world's children? A WHO-UNICEF-Lancet Commission. *Lancet* 2020; **395**: 605–58.
- 8 Global Grand Challenges. Affordable, accessible, and appealing: the next generation of nutrition (round 21). March 5, 2018. <https://gcgh.grandchallenges.org/challenge/affordable-accessible-and-appealing-next-generation-nutrition-round-21> (accessed April 15, 2020).
- 9 Pyakurel S. WFP violated Paris Declaration on aid effectiveness. Republica. July 5, 2015. <http://archive.myrepublica.com/2015-16/politics/story/24062/wfp-violated-paris-declaration-on-aid-effectiveness-insec.html> (accessed April 15, 2020).
- 10 Shelton AM, Sarwer SH, Hossain MJ, Brookes G, Paranjape V. Impact of Bt brinjal cultivation in the market value chain in five districts of Bangladesh. *Front Bioeng Biotechnol* 2020; **8**: 498.