

4-11-2017

## Structure of Nascent Chromatin Is Essential for Hematopoietic Lineage Specification.

Svetlana Petruk  
*Thomas Jefferson University*

Samanta A. Mariani  
*Thomas Jefferson University; The University of Edinburgh*

Marco De Dominicis  
*Thomas Jefferson University*

Patrizia Porazzi  
*Thomas Jefferson University*

Valentina Minieri  
*Thomas Jefferson University*  
Follow this and additional works at: <https://jdc.jefferson.edu/cbfp>

 Part of the [Medicine and Health Sciences Commons](#)

*See next page for additional authors*

[Let us know how access to this document benefits you](#)

---

### Recommended Citation

Petruk, Svetlana; Mariani, Samanta A.; De Dominicis, Marco; Porazzi, Patrizia; Minieri, Valentina; Cai, Jingli; Iacovitti, Lorraine; Flomenberg, Neal; Calabretta, Bruno; and Mazo, Alexander, "Structure of Nascent Chromatin Is Essential for Hematopoietic Lineage Specification." (2017). *Department of Cancer Biology Faculty Papers*. Paper 115.  
<https://jdc.jefferson.edu/cbfp/115>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Cancer Biology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).


---

**Authors**

Svetlana Petruk, Samanta A. Mariani, Marco De Dominicis, Patrizia Porazzi, Valentina Minieri, Jingli Cai, Lorraine Iacovitti, Neal Flomenberg, Bruno Calabretta, and Alexander Mazo

## Structure of Nascent Chromatin Is Essential for Hematopoietic Lineage Specification

### Graphical Abstract



### Authors

Svetlana Petruk, Samanta A. Mariani, Marco De Dominici, ..., Neal Flomenberg, Bruno Calabretta, Alexander Mazo

### Correspondence

bruno.calabretta@jefferson.edu (B.C.), alexander.mazo@jefferson.edu (A.M.)

### In Brief

Petruk et al. find that hematopoietic progenitor cells possess a state of post-replicative chromatin globally devoid of the repressive histone mark H3K27me3. This de-condensed chromatin is required for recruitment of lineage-determining transcription factors to DNA just after replication in early stages of cytokine-induced myeloid or erythroid lineage specification.

### Highlights

- Accumulation of H3K27me3 on nascent DNA occurs rapidly in undifferentiated HPCs
- Accumulation of H3K27me3 on nascent DNA is delayed 12 hr following induction
- This delay is caused by induction of the H3K27me3 KDM UTX
- “Open” nascent chromatin facilitates recruitment of lineage-specifying TFs to DNA



# Structure of Nascent Chromatin Is Essential for Hematopoietic Lineage Specification

Svetlana Petruk,<sup>1,5</sup> Samanta A. Mariani,<sup>2,5,6</sup> Marco De Dominicis,<sup>2</sup> Patrizia Porazzi,<sup>2</sup> Valentina Minieri,<sup>2</sup> Jingli Cai,<sup>3</sup> Lorraine Iacovitti,<sup>3</sup> Neal Flomenberg,<sup>4</sup> Bruno Calabretta,<sup>2,\*</sup> and Alexander Mazo<sup>1,7,\*</sup>

<sup>1</sup>Department of Biochemistry and Molecular Biology

<sup>2</sup>Department of Cancer Biology

<sup>3</sup>Department of Neuroscience

<sup>4</sup>Department of Medical Oncology

Sidney Kimmel Medical College, Kimmel Cancer Center, Thomas Jefferson University, Philadelphia, PA 19107, USA

<sup>5</sup>Co-first author

<sup>6</sup>Present address: The Queen's Medical Research Institute, Centre for Inflammation Research, The University of Edinburgh, Edinburgh EH8 9YL, UK

<sup>7</sup>Lead Contact

\*Correspondence: [bruno.calabretta@jefferson.edu](mailto:bruno.calabretta@jefferson.edu) (B.C.), [alexander.mazo@jefferson.edu](mailto:alexander.mazo@jefferson.edu) (A.M.)  
<http://dx.doi.org/10.1016/j.celrep.2017.03.035>

## SUMMARY

The role of chromatin structure in lineage commitment of multipotent hematopoietic progenitors (HPCs) is presently unclear. We show here that CD34<sup>+</sup> HPCs possess a post-replicative chromatin globally devoid of the repressive histone mark H3K27me3. This H3K27-unmodified chromatin is required for recruitment of lineage-determining transcription factors (TFs) C/EBP $\alpha$ , PU.1, and GATA-1 to DNA just after DNA replication upon cytokine-induced myeloid or erythroid commitment. Blocking DNA replication or increasing H3K27me3 levels prevents recruitment of these TFs to DNA and suppresses cytokine-induced erythroid or myeloid differentiation. However, H3K27me3 is rapidly associated with nascent DNA in more primitive human and murine HPCs. Treatment of these cells with instructive cytokines leads to a significant delay in accumulation of H3K27me3 in nascent chromatin due to activity of the H3K27me3 demethylase UTX. Thus, HPCs utilize special mechanisms of chromatin modification for recruitment of specific TFs to DNA during early stages of lineage specification.

## INTRODUCTION

The process of hematopoiesis is regulated by a network of transcription factors (TFs) acting cooperatively or antagonistically on multipotent or lineage-restricted hematopoietic progenitors/precursors (Friedman, 2007; Orkin and Zon, 2008). A critical step in differentiation of hematopoietic stem cells (HSCs) is the commitment of multipotent progenitors to lineage-restricted progenitors. The prevailing explanation for lineage commitment is that it depends on stochastic fluctuation in the levels of

competing lineage-determining TFs that activate a lineage-specific gene expression program while suppressing those leading to alternative choices (Cantor and Orkin, 2001; Chang et al., 2008; Graf and Enver, 2009; Orkin, 2000; Till et al., 1964).

However, recent studies support an “instructive” model whereby lineage-specific cytokines (e.g., granulocyte-colony stimulating factor [G-CSF], macrophage-colony stimulating factor [M-CSF], or erythropoietin [EPO]) act on multipotent progenitors to activate lineage-specific master TFs and promote the expansion of the myeloid or the erythroid lineage (Grover et al., 2014; Mossadegh-Keller et al., 2013; Rieger et al., 2009). On the other hand, single-cell transcription profiling of primitive hematopoietic progenitors purified based on the expression of surface markers commonly used to define a “multipotent state” has revealed that truly multipotent progenitors may not exist and that these cells are highly heterogeneous and may be already committed to a specific lineage (Notta et al., 2016; Paul et al., 2015; Perié et al., 2015). Collectively, these findings raise the possibility that both “stochastic” and “instructive” mechanisms contribute to lineage specification and that “instructive” cytokines function in this process earlier than previously thought. Specification of myeloid cell fate requires the TFs PU.1 and C/EBP $\alpha$ , but macrophage versus granulocytic differentiation depends on the relative levels of PU.1 or C/EBP $\alpha$ , respectively (Dahl et al., 2003; Iwasaki et al., 2006; Laslo et al., 2006). EPO has an instructive role for erythroid commitment of stem cell-enriched cell subsets, a process that is, at least in part, GATA-1-dependent (Cui et al., 2009; Grover et al., 2014).

While C/EBP $\alpha$ , PU.1, or GATA-1 can promote unilineage differentiation by inducing new transcriptional programs, it remains unclear how these TFs cause changes in the existing epigenetic maintenance system. Upon EPO-induced erythroid differentiation of the multipotent CD133<sup>+</sup>CD34<sup>+</sup> HPCs global changes in gene expression are in general correlated with the changes in association of the H3K4me3 or the H3K27me3 positive or negative histone marks respectively, although some “bivalent” genes lost or retained both marks (Cui et al., 2009). These changes in

chromatin marks and gene expression profiles were detected in cells induced to erythroid differentiation for 9–11 days (Cui et al., 2009), suggesting that they may not reflect “commitment” to differentiation toward a specific lineage that is likely to occur much earlier. The dynamics of certain chromatin modifications was also globally examined in purified subsets representing distinct stages of murine hematopoietic differentiation (Lara-Astiaso et al., 2014). The results of this analysis are consistent with the existence of a transcription factor network that may control chromatin dynamics and lineage specification.

However, these studies are mostly correlative and do not provide a mechanistic explanation for how changes in transcriptional programs and chromatin structure occur. An essential issue is how newly induced TFs overcome the existing chromatin-based barrier when they are recruited to new target genes. It is also not clear whether chromatin marks play any role in allowing the access of lineage-determining TFs to the regulatory regions of the genes essential for differentiation and whether chromatin changes occur before or after changes in the transcriptional programs. Thus, at present we only know the end results of lineage specification, but there is insufficient information on the molecular mechanisms that allow these changes during the early stages of lineage commitment. Knowledge of these mechanisms is essential in order to control lineage specification and may be important in understanding the mechanisms responsible for the differentiation arrest of leukemic cells.

In our study, we examined the assembly of chromatin in vivo at single-cell and gene-specific levels following DNA replication. We show that CD34<sup>+</sup> HPCs possess a post-replicative chromatin globally devoid of the repressive histone mark H3K27me3. This type of chromatin is required for recruitment of C/EBP $\alpha$ , PU.1, and GATA-1 to DNA just after DNA replication upon treatment with instructive cytokines to induce erythroid or myeloid commitment. Blocking DNA replication or increasing H3K27me3 levels prevents recruitment of these TFs to DNA and suppresses cytokine-induced erythroid or myeloid differentiation. By contrast, H3K27me3 is rapidly associated with nascent DNA in primitive CD34<sup>+</sup>CD38<sup>-</sup> HPCs, and treatment of these cells with instructive cytokines leads to a period of de-condensed, H3K27me3-unmodified nascent chromatin due to increase of the H3K27me3 demethylase UTX. These studies suggest that HPCs utilize special mechanisms of chromatin alteration for recruitment of specific TFs to DNA during initial stages of differentiation.

## RESULTS

### Dynamics of H3K27me3 Chromatin during Lineage Specification of Multipotent Hematopoietic Progenitors

Our previous studies suggested that accumulation of major histone modifications, including H3K27me3, following DNA replication is delayed by at least 1 hr in cells of developing *Drosophila* embryos (Petruk et al., 2012). This suggests that modified histones do not directly carry epigenetic information and implies that post-replicative chromatin may transiently consist of nucleosomes lacking H3K27me3 marks, which were not detected previously in bulk chromatin (Petruk et al., 2012). Because the


absence of repressive H3K27me3 correlates with low density of nucleosomes (Bell et al., 2010; Shlyueva et al., 2014; Yuan et al., 2012), such unmodified histones may create a very de-condensed conformation of post-replicative chromatin. We hypothesize that this unusual H3K27me3-unmodified post-replicative chromatin may be essential during early stages of differentiation to create more favorable conditions for binding of newly induced lineage-determining TFs.

First, we examined whether chromatin lacking H3K27me3 is a common feature of human HPCs. We used the chromatin assembly assay (CAA) (Petruk et al., 2012) to examine the rate of accumulation of H3K27me3 in human cord blood and G-CSF-mobilized CD34<sup>+</sup> progenitor cells. DNA in proliferating CD34<sup>+</sup> cells was pulse-labeled with 5-ethynyl-2'-deoxyuridine (EdU), which was then conjugated with biotin, and the proximity of H3K27me3 to nascent DNA was assessed by the proximity ligation assay (PLA, Olink) using antibodies to biotin and H3K27me3. Following CAA, cells were immunostained with EdU and DAPI to assess the specificity of the assay. We found that in CD34<sup>+</sup> cells H3K27me3 accumulates significantly at nascent DNA only at around 2 hr following DNA replication (Figure 1A).

Then, we examined whether the accumulation of post-replicative H3K27me3 undergoes changes during lineage specification of HPCs. Cells were induced toward granulocytic or macrophage differentiation by treatment with G-CSF or M-CSF, respectively, and accumulation of H3K27me3 on nascent DNA was analyzed after 15 min of EdU incorporation. No H3K27me3 was detected on 15 min EdU-labeled nascent DNA at 6 hr and 12 hr after G-CSF or M-CSF treatment. However, we detected significant accumulation of H3K27me3 on 15 min EdU-labeled nascent DNA at 24 hr after induction with each cytokine (Figure 1B). Thus, irrespective of lineage choice, the H3K27me3-unmodified post-replicative chromatin structure of HPCs persists for at least 12 hr following exposure to instructive cytokines. The post-replicative structure of chromatin may become more compact 1 day into HPC differentiation.

### Lineage-Determining Transcription Factors Bind to Post-replicative H3K27me3-Unmodified Chromatin upon Cytokine-Induced Lineage Commitment

Induction with “instructive” cytokines leads to expression of key TFs that specifically change the transcriptional landscape of differentiating HPCs. It is not known whether there is a specific mechanism that allows these TFs to bind to DNA during cell differentiation. Previous EMSA analyses showed that C/EBP $\alpha$  is associated with promoters of target genes after 3 days of G-CSF-induced differentiation of myeloid precursor cells (Iida et al., 2008). To further investigate this, we used the CAA to examine the association of C/EBP $\alpha$ , PU.1, and GATA-1 with DNA following induction with G-CSF, M-CSF, and EPO, respectively, of cord blood or G-CSF-mobilized peripheral blood CD34<sup>+</sup> cells. As expected, cytokine treatment induced a decrease in CD34 positivity first detected at 72 hr and the simultaneous appearance of lineage-specific differentiation markers (Figures 2A and S1). C/EBP $\alpha$  and PU.1 are first detected on the 1 hr EdU-labeled nascent DNA at 6 hr following treatment with G-CSF and M-CSF, respectively, while GATA-1 is detected at 12 hr following induction with EPO (Figure 2B). Thus, irrespective



**Figure 1. Kinetics of H3K27me3 Accumulation following DNA Replication in Cytokine-Treated CD34<sup>+</sup> HPCs**

(A) Left: accumulation of H3K27me3 on nascent DNA of G-CSF-mobilized CD34<sup>+</sup> HPCs. DNA was labeled with EdU for 15 min and chased to 1, 2, and 4 hr. Following conjugation with biotin, CAA was performed between nascent DNA (biotin) and H3K27me3. Red, PLA; green, EdU (biotin); blue, DAPI. Lower panels show PLA signals only. Right: quantification of the results of CAA experiments shown to the left by counting the number of PLA signals per EdU-labeled nuclei in 50 cells/each of the three independent experiments.


(B) Left: accumulation of post-replicative H3K27me3 during differentiation of CD34<sup>+</sup>HPCs. Cells were induced for myeloid differentiation with either G-CSF (upper panels) or M-CSF (lower panels) for 0, 6, 12, and 24 hr. DNA was labeled with EdU for 15 min and CAA performed between EdU (biotin) and H3K27me3. Red, PLA; green, EdU (biotin); blue, DAPI. Lower panels show PLA signals only. Right: quantification of the results of CAA experiments shown to the left by counting the number of PLA signals per EdU-labeled nuclei in 50 cells/each of the three independent experiments. Error bars represent  $\pm$  SD. p Values were determined by ANOVA. ns, non-significant; \*p < 0.05.

of the induced differentiation program, specific TFs are recruited to DNA at 6–12 hr following treatment with lineage-determining cytokines, much earlier than previously thought. Importantly, the time of recruitment is within the period in which cytokine-treated HPCs carry the H3K27me3-unmodified post-replicative chromatin.

These results are consistent with our hypothesis that recruitment of key lineage-determining TFs to DNA may be facilitated by the low-nucleosome density of post-replicative chromatin devoid of H3K27me3. This state of histone H3K27 modification may result from the activities of H3K27me3-specific homologous histone demethylases (KDMs) UTX and JMJD3. To examine this, we inhibited the activities of these KDMs by treatment with GSKJ4, a small molecule inhibitor of these enzymes (Kruidenier et al., 2012). Then, cord blood CD34<sup>+</sup> cells were induced for 12 hr to myeloid or erythroid differentiation with G-CSF or M-CSF,

or EPO, respectively, and accumulation of H3K27me3 was analyzed by CAA on nascent DNA labeled with EdU for 15 min. Similar to previous results with G-CSF and M-CSF (Figure 1), we did not detect H3K27me3 on 15 min EdU-labeled nascent DNA following 12 hr induction with EPO (Figure 3A, upper panels). Treatment of cells with GSKJ4 leads to significant accumulation of H3K27me3 on nascent DNA labeled with EdU for 15 min following 12 hr induction of these cells with all three lineage-determining cytokines (Figure 3A, lower panels). Then, we asked whether faster accumulation of post-replicative H3K27me3 affects recruitment of induced TFs in these cell lineages. The strong recruitment of C/EBP $\alpha$ , PU.1, and GATA-1, detected at 12 hr following induction with the corresponding cytokine (Figures 2 and 3A, upper panels), was completely blocked by treatment with GSKJ4 (Figure 3A, lower panels). This suggests that the H3K27me3-unmodified post-replicative





**Figure 2. Kinetics of Recruitment of Lineage-Determining TFs C/EBP $\alpha$ , PU.1, and GATA-1 to DNA**

(A) In vitro differentiation of G-CSF-mobilized CD34<sup>+</sup> cells. Cells were treated with G-CSF, M-CSF, or EPO for the indicated times. Differentiation was assessed by flow cytometry for lineage-specific markers.

(B) G-CSF-mobilized CD34<sup>+</sup> HPCs were induced toward myeloid or erythroid differentiation by treatment with G-CSF, M-CSF, or EPO, respectively, for 6, 12, and 24 hr. DNA was pulse-labeled with EdU for 15 min and chased to 1 hr. CAA was performed between nascent DNA (biotin) and C/EBP $\alpha$ , PU.1, or GATA-1. red, PLA; green, EdU (biotin), blue, DAPI. Lower panel shows PLA signals only. Bottom: quantification of the results of CAA experiments shown above by counting the number of PLA signals per EdU-labeled nuclei in 50 cells/each of the three independent experiments. Error bars represent  $\pm$  SD. p Values were determined by ANOVA. \*p < 0.05.

CD15 differentiation antigens (27.5 versus 53.2%) (Figure S2B). Of interest, at the 72 hr time point chosen for the analysis of erythroid differentiation markers, CD34 positivity was essentially identical in cells treated with EPO only or with GSKJ4 and EPO (Figure S3). Thus, the H3K27me3 unmodified structure of nascent chromatin is essential for lineage specification of HPCs.

### C/EBP $\alpha$ Is Recruited to Post-replicative Chromatin in the S Phase of the Cell Cycle

It is not well established when lineage-specific TFs are recruited to DNA in undifferentiated cells; while some studies propose that this occurs during the G1 phase (Kapinas et al., 2013; Murray and Kirschner, 1989; Singh and Dalton, 2009), other studies dispute this idea (Filipczyk et al., 2007; Jonk et al., 1992; Li et al., 2012; Mummery et al., 1987).

To examine this, CD34<sup>+</sup> cells were labeled with EdU and then treated with G-CSF for 24 hr. This allows detection of C/EBP $\alpha$  on labeled DNA (Figure 4A, left). However, blocking S phase entry in these cells by treatment with thymidine (Figure 4B, left) prevented the accumulation of C/EBP $\alpha$  on EdU-labeled DNA (Figure 4A and 0 hr). Release from thymidine block for 2 and 4 hr allows significant re-initiation of DNA replication (Figure 4B, middle and right) and leads to efficient accumulation of C/EBP $\alpha$  on nascent DNA (Figure 4A, 2 and 4 hr). These findings support our hypothesis that the initial association of TFs with DNA following cytokine treatment occurs primarily during early post-replicative periods in S phase.


### Accumulation of H3K27me3-Modified Chromatin in Primitive HPCs

To assess whether the H3K27me3-unmodified chromatin also exists in more primitive HPCs, we examined the loading of

chromatin is favorable for recruitment of C/EBP $\alpha$ , PU.1, and GATA-1 to their DNA binding sites.

By contrast, maintaining H3K27me3 accumulated on nascent chromatin may prevent association of key lineage-inducing TFs to DNA, leading to decreased induction of the gene targets of these TFs, and may ultimately result in a block of lineage specification.

In line with this, expression of GATA-1 was not affected by GSKJ4 treatment of EPO-stimulated CD34<sup>+</sup> cells (Figure 3B); by contrast, inhibiting GATA-1 binding to DNA (Figure 3A, right panel) led to a marked decrease in the expression of GATA-1-regulated genes (Welch et al., 2004) and a complete block of EPO-induced differentiation revealed by loss of erythroid marker expression (Figures 3C and 3D). Likewise, treatment with GSKJ4 had no effect on G-CSF-induced C/EBP $\alpha$  expression but suppressed cytokine-induced myeloid differentiation of mobilized CD34<sup>+</sup> cells, as indicated by its ability to block the increase in the expression of several C/EBP $\alpha$ -regulated genes (Figure S2A). Moreover, compared to treatment with G-CSF only, co-treatment with GSKJ4 resulted in a less differentiated phenotype revealed by the marked decrease in the proportion of cells co-expressing the CD33 and the



**Figure 3. H3K27me3-Unmodified Post-replicative Chromatin Is Essential for DNA Binding of Lineage-Determining TFs**

(A) Top: accumulation of post-replicative H3K27me3 leads to decreased recruitment of lineage-determining TFs. Cord blood CD34<sup>+</sup> cells were induced toward myeloid differentiation with G-CSF (left) or M-CSF (middle), or erythroid differentiation with EPO (right) for 12 hr. During the induction period cells were left untreated (upper panels) or treated with the GSKJ4 inhibitor of KDMs UTX


H3K27me3 on nascent DNA in the CD90<sup>+</sup> subpopulation of CD34<sup>+</sup> cells. Surprisingly, CD90<sup>+</sup> cells exhibited high accumulation of H3K27me3 on DNA labeled with EdU for only 15 min (Figure 5A). Because most CD34<sup>+</sup>CD90<sup>+</sup> cells are contained within the CD34<sup>+</sup>CD38<sup>-</sup> subset, we confirmed this fast mode of accumulation of H3K27me3 in uninduced CD34<sup>+</sup>CD38<sup>-</sup> cells (Figure 5B). However, 12 hr induction of CD34<sup>+</sup>CD38<sup>-</sup> cells with G-CSF or M-CSF led to slow accumulation of post-replicative H3K27me3 (Figure 5B). Further treatment for 24 hr restored fast accumulation of post-replicative H3K27me3. This suggests that compared to CD34<sup>+</sup> cells, most of which carry H3K27me3-unmodified post-replicative chromatin, H3K27me3-containing chromatin forms very fast after DNA replication in the more primitive CD34<sup>+</sup>CD90<sup>+</sup> and CD34<sup>+</sup>CD38<sup>-</sup> subsets. Induction with cytokines slows post-replicative accumulation of H3K27me3, making it similar to that of bulk CD34<sup>+</sup> cells. Similar to CD34<sup>+</sup> cells, further differentiation accelerates accumulation of post-replicative H3K27me3 in both types of cells (Figures 1B and 5A). The differences in the initial rate of accumulation of post-replicative H3K27me3 between CD34<sup>+</sup>, CD34<sup>+</sup>CD38<sup>-</sup>, and CD34<sup>+</sup>CD90<sup>+</sup> cells may be explained by the fact that a significant proportion of bulk CD34<sup>+</sup> cells likely consists of progenitors with the molecular profile of cells committed to a specific lineage but unable to further differentiate in the absence of instructive cytokines. Instead, the H3K27me3-modified chromatin of more primitive CD34<sup>+</sup>CD38<sup>-</sup> and CD34<sup>+</sup>CD90<sup>+</sup> progenitors may reflect the undifferentiated status of hematopoietic stem cells (HSCs) and, upon cytokine treatment, exhibit significant changes in the rate of accumulation of H3K27me3 after DNA replication. Similar to that detected in bulk CD34<sup>+</sup> cells (Figure 2B), G-CSF treatment of CD34<sup>+</sup>CD38<sup>-</sup> cells promotes the accumulation of the lineage-determining TF C/EBP $\alpha$  on nascent DNA (Figure S4).

We also assessed the loading of H3K27me3 on nascent DNA of primitive Lin-Sca-1<sup>+</sup>Kit<sup>+</sup> mouse HPCs. Like human CD34<sup>+</sup>CD38<sup>-</sup> cells, these cells exhibited high accumulation of H3K27me3 on DNA labeled with EdU for only 15 min (Figure S5); moreover, treatment of Lin-Sca-1<sup>+</sup>Kit<sup>+</sup> cells with recombinant murine G-CSF for 4 hr induced a state of globally de-condensed,

and JMJD3 (lower panels). DNA was pulse-labeled with EdU for 15 min and chased to 1 hr. CAA was performed between nascent DNA (biotin) and C/EBP $\alpha$  (left), PU.1 (middle), or GATA-1 (right). Red, PLA; green, EdU (biotin); blue, DAPI. Lower panel shows PLA signals only. Bottom: quantification of the results of CAA experiments shown above by counting the number of PLA signals per EdU-labeled nuclei in 50 cells/each of the three independent experiments. (B and C) G-CSF-mobilized CD34<sup>+</sup> cells were cultured for 24 hr in the presence of the CC100 cytokine cocktail (CTRL), or with EPO only (EPO) or with EPO and GSKJ4 (EPO + GSKJ4). (B) Cells were immunostained with antibody to GATA-1 (red). Blue, DAPI. Lower panel shows GATA-1 signals only. (C) Western blot analysis with GATA1 and actin antibodies.

(D and E) Effect of UTX/JMJD3 inhibition on EPO-induced erythroid differentiation of CD34<sup>+</sup> HPCs. Cells were cultured for 72 hr in the presence of the CC100 cytokine cocktail (CTRL), in the presence of EPO only (EPO), or in the presence of EPO only and GSKJ4 (EPO + GSKJ4). (D) Real-time PCR analysis (performed in triplicate) of GATA-1 target genes in untreated EPO-, or EPO plus GSKJ4-treated CD34<sup>+</sup>CD38<sup>+</sup> cells. (E) Erythroid differentiation monitored by flow cytometry. Error bars represent  $\pm$  SD. p Values were determined by Student's t test. \*\*\*p < 0.001.





**Figure 4. C/EBP $\alpha$  Is Recruited to DNA Only during Post-replicative Period**

(A) Cord blood CD34<sup>+</sup> HPCs were labeled with EdU for 30 min and induced with G-CSF for 24 hr in the absence (Cont) or presence of 2.5 mM thymidine. After release from thymidine block, CAA was performed between nascent DNA (biotin) and C/EBP $\alpha$  at 0, 2, and 4 hr. Red, PLA; green, EdU (biotin); blue, DAPI. Lower panel shows PLA signals only. A graph below shows quantification of the results of CAA experiments by counting the number of PLA signals per EdU-labeled nuclei in 50 cells/each of the three independent experiments.

(B) Following DNA labeling with EdU for 30 min and induction with G-CSF, CD34<sup>+</sup> HPCs were grown for 24 hr in the presence of thymidine. Thymidine block was released for 0, 2, and 4 hr, and cells were labeled for 20 min with BrdU. Cells were double-immunostained for EdU (green) and BrdU (red). Error bars represent  $\pm$  SD. p Values were determined by ANOVA. \*p < 0.05.


delayed in regulatory regions of repressed genes that contain H3K27me3 during transcriptional interphases. To examine this, we analyzed the kinetics of accumulation of H3K27me3 on nascent DNA at the promoters of several repressed genes by re-chromatin immunoprecipitation (ChIP) assays (Francis et al., 2009; Petruk et al., 2012). Because this type of analysis requires  $\sim 10^7$  cells per each sample, it is not feasible to use purified CD34<sup>+</sup>CD38<sup>-</sup> cells in these assays. Thus, we used batches of CD34<sup>+</sup> cells which contain significant proportion of CD38<sup>-</sup> cells. DNA of uninduced CD34<sup>+</sup> cells and CD34<sup>+</sup> cells induced with EPO was labeled with BrdU for 15 min or for 15 min followed by a chase to 2 hr. Cells were induced with EPO for 24 hr because induction with this cytokine leads to a longer period of delayed accumulation of H3K27me3 on nascent DNA than after induction with G-CSF (not shown). Chromatin was first immunoprecipitated with antibody to H3K27me3, and in the next step the purified DNA was immunoprecipitated with antibody to BrdU. ChIP assays at the first immunoprecipitation step demonstrate that four examined genes contain similar amounts of H3K27me3 at their promoters before and after EPO induction (Figure 6A), confirming that they remain repressed during erythroid differentiation (Cui et al., 2009). In re-ChIP assays, we detected significant accumulation of H3K27me3 on 15 min BrdU-labeled DNA (Figure 6B), suggesting a fast mode of accumulation of H3K27me3 characteristic for early progenitor cells (Figure 5). Accumulation of H3K27me3 is significantly delayed at the promoters of all repressed genes on 15 min of BrdU-labeled DNA following induction with EPO (Figure 6B). These results confirm at a

H3K27me3-unmodified nascent chromatin. However, an 8 hr treatment of these cells with G-CSF was sufficient to induce the rapid accumulation of H3K27me3 on nascent DNA. Together, these data indicate that accumulation of H3K27me3 on nascent DNA is similar in primitive human and mouse HPCs, although in the latter cells the kinetics appears to be somewhat faster.

#### Accumulation of H3K27me3 Is Delayed on Nascent Chromatin of Repressed Genes

Our results suggest that following induction with cytokines accumulation of H3K27me3 is globally delayed following DNA replication. This implies that accumulation of H3K27me3 is

precipitated with antibody to BrdU. ChIP assays at the first immunoprecipitation step demonstrate that four examined genes contain similar amounts of H3K27me3 at their promoters before and after EPO induction (Figure 6A), confirming that they remain repressed during erythroid differentiation (Cui et al., 2009). In re-ChIP assays, we detected significant accumulation of H3K27me3 on 15 min BrdU-labeled DNA (Figure 6B), suggesting a fast mode of accumulation of H3K27me3 characteristic for early progenitor cells (Figure 5). Accumulation of H3K27me3 is significantly delayed at the promoters of all repressed genes on 15 min of BrdU-labeled DNA following induction with EPO (Figure 6B). These results confirm at a



**Figure 5. H3K27me3-Containing Nascent Chromatin Undergoes Dynamic Changes in Cytokine-Treated Primitive HPCs**

(A) CD34<sup>+</sup> HPCs were labeled with EdU for 15 min. CAA was performed for H3K27me3 antibody. Cells were then immunostained for CD90 (green) and EdU (blue).

(B) Induction of CD34<sup>+</sup>CD38<sup>-</sup> HPCs with G-CSF (top) or M-CSF (bottom) leads to de-methylation of H3K27me3. Purified CD34<sup>+</sup>CD38<sup>-</sup> HPCs were induced with G-CSF (upper panels) or M-CSF (lower panels) for 0, 12, and 24 hr. Cells were labeled with EdU for 15 min. CAA was performed between nascent DNA (biotin) and H3K27me3 antibodies. Red, PLA; green, EdU (biotin); blue, DAPI. Lower panel shows PLA signals only. Quantification of the results of CAA experiments is shown under images and was estimated by counting the number of PLA signals per EdU-labeled nuclei in 50 cells/each of the three independent experiments. Error bars represent  $\pm$  standard deviation. p Values were determined by ANOVA. \*p < 0.05.


gene level that accumulation of H3K27me3 on nascent DNA is delayed following induction of differentiation of HPCs.

### Cytokine Treatment Leads to Induction of H3K27me3 KDMs UTX and JMJD3 in Primitive HPCs

The observed differences in the rate of accumulation of H3K27me3 in CD34<sup>+</sup>CD38<sup>-</sup> cells may depend on changes in the amounts or the activities of the H3K27-specific methylating and de-methylating enzymes, such as KMT EZH2 and KDM UTX, respectively. Thus, we examined the presence of these enzymes on DNA following DNA replication. Similar amounts of the KMT EZH2 were detected in uninduced CD34<sup>+</sup>CD38<sup>-</sup> cells and in the same cells induced for 12 hr to myeloid differentiation by G-CSF or M-CSF treatment (Figure 7A). However, UTX was not detected on DNA in untreated CD34<sup>+</sup>CD38<sup>-</sup> cells, but was detected on DNA of these cells 12 hr following induction with each cytokine (Figure 7A). These results were confirmed by several independent assays. Western blot analysis revealed an increase in the UTX protein in G-CSF- or M-CSF-treated (12 hr) CD34<sup>+</sup>CD38<sup>-</sup>, whereas UTX levels did not change after cytokine treatment of the less primitive CD34<sup>+</sup>CD38<sup>+</sup> progenitors (Figure 7B); compared to high amounts of EZH2 in all CD34<sup>+</sup> cells, UTX was barely detectable by immunofluorescence (Figure 7C), or by CAA on 1 hr EdU-labeled DNA (Figure 7D) in the nuclei of more primitive CD90<sup>+</sup> cells. Likewise, using CAA, we detected low amounts of the homologous H3K27me3 KDM JMJD3 on DNA of the CD90<sup>+</sup> cells and an increase in this protein in the CD90<sup>-</sup> cells (Figure 7D). Thus, a 12-hr cytokine treatment of primitive HPCs leads to induction of the KDMs UTX and JMJD3 proteins and their further accumulation on DNA. This may explain the switch detected in these cells from the fast to the slow mode of accumulation of post-replicative H3K27me3 after DNA replication (Figure 5B).

### DISCUSSION

Cytokine-induced lineage specification of CD34<sup>+</sup> HPCs involves the recruitment of lineage-determining TFs (e.g., GATA-1 or C/EBP $\alpha$ ) to the regulatory regions of differentiation-related genes and is accompanied by complex changes in transcriptional programs and chromatin structure (e.g., patterns of H3K4me3 and H3K27me3 histone marks) (Cui et al., 2009; Lara-Astiaso et al., 2014; Mercer et al., 2011; Wei et al., 2009). However, it is not



**Figure 6. Kinetics of H3K27me3 Accumulation at Nascent DNA of Specific Genes during Erythroid Differentiation of CD34<sup>+</sup> HPCs**  
 (A) ChIP assays with the H3K27me3 antibody. Uninduced and 24 hr EPO-induced CD34<sup>+</sup> cells were labeled with BrdU for 15 min or 15 min and chased to 60 min. Chromatin was immunoprecipitated with H3K27me3 and IgG antibodies. (B) ChIP assays with the BrdU antibody. DNA from the first immunoprecipitation step with H3K27me3 antibody (A) was denatured and immunoprecipitated with the BrdU antibody. Q-PCR was performed at genes repressed before and after EPO induction (GPR85, WNT16, HOXB8, HOXB13) and at a non-repressed control gene (CD11B) (Cui et al., 2009). Percent of input for re-ChIP experiments was calculated using the amount of material eluted from the beads following immunoprecipitation with H3K27me3 as 100% input. Data are represented as mean  $\pm$  SD of technical replicates of one of the independent experiments. p Values were calculated by a one-way ANOVA with Bonferroni's post hoc test. The statistical significance was defined as  $p < 0.05$ .

known whether binding of TFs to new target genes is a regulated process and whether it is affected by histone modifications and the structure of chromatin. Thus, we have a very limited view of the molecular mechanisms regulating a switch in transcriptional and epigenetic programs in dividing HPCs as these cells undergo lineage specification.

In this study, we uncovered a chromatin-based molecular mechanism of HPCs differentiation toward a specific lineage. The H3K27me3 mark rapidly associates with nascent DNA in primitive human CD34<sup>+</sup>CD38<sup>-</sup> cells (see model in Figure 7E) and mouse Lin-Sca-1<sup>+</sup>Kit<sup>+</sup> cells. Upon induction with instructive cytokines, the global accumulation of H3K27me3 on nascent DNA is significantly delayed. In line with this, accumulation of H3K27me3 in nascent chromatin is also globally delayed in the more differentiated CD34<sup>+</sup>CD38<sup>+</sup> HPCs. Our data suggest that transition from a more compact, H3K27me3-modified, to a more de-condensed, H3K27me3-unmodified chromatin structure, may be explained by the induction of H3K27me3 KDMs UTX and JMJD3 that may efficiently de-methylate this residue.


Once established, the H3K27-unmodified chromatin exists for a short period of time at very early stages of differentiation. During this period, cytokines also induce expression of lineage-specific TFs that quickly associate with DNA (Figure 7E). The transient H3K27-unmodified chromatin state is essential for the initial association of these TFs with nascent DNA that occurs at early stages of DNA replication. Remarkably, maintaining the H3K27me3 repressive mark on nascent chromatin of CD34<sup>+</sup> HPCs by pharmacological inhibition of the H3K27me3 demethylases blocks the association of lineage-determining TFs to DNA and suppresses cytokine-induced erythroid or myeloid differentiation (Figures 3 and S2). Consistent with these findings, conditional knockout of UTX in female mice induces marked depletion of erythroid progenitors in the bone marrow and of erythrocytes

in the peripheral blood, in addition to profound thrombocytopenia and relative leukocytopenia (Thieme et al., 2013).

Fast accumulation of the H3K27me3 mark on nascent DNA resumes once CD34<sup>+</sup> HPCs becomes committed to a single lineage (Figure 7E); this process may be designed to ensure that uni-lineage committed progenitors complete their differentiation program by hindering DNA binding of unwanted TFs that may antagonize the effect of the lineage-determining TF or cooperatively promote differentiation along a closely related lineage (Friedman, 2007).

The lineage-determining TFs analyzed in this work belong to the class of pioneer factors (Iwafuchi-Doi and Zaret, 2014; Zaret and Carroll, 2011) that are able to overcome the barrier of chromatin, bind to the so-called L-type chromatin that contains nucleosomes without many activating or repressive marks including H3K27me3, and facilitate binding of other essential TFs (Iwafuchi-Doi and Zaret, 2016). The pioneer property of these TFs was demonstrated mostly during reprogramming of differentiated cells following ectopic expression of these proteins (Di Stefano et al., 2014; Feng et al., 2008; Heinz et al., 2010; Kulesa et al., 1995; Xie et al., 2004). However, actively repressed chromatin (R-type), which includes genes marked by H3K27me3, is not accessible to any TF, including pioneer factors (Iwafuchi-Doi and Zaret, 2016).

Our work uncovers a chromatin-based mechanism of recruitment to DNA of lineage-determining TFs induced at physiological levels by cytokines during differentiation of hematopoietic progenitor cell subsets. Based on this model, lineage-determining TFs do not need to overcome the chromatin barrier as proposed for pioneer factors on L-type chromatin, but gain access to their specific binding sites on a transiently "open" chromatin of nascent DNA. This mechanism of recruitment to DNA of lineage-determining TFs is not limited to HPCs, as binding to H3K27-unmethylated post-replicative chromatin is also a feature



**Figure 7. Induction of H3K27me3 KDMs UTX and JMJD3 during Lineage Specification**

(A) Kinetics of EZH2 (left panels) and UTX (right panels) accumulation during induction of CD34<sup>+</sup>CD38<sup>-</sup> HPCs with G-CSF (upper panels) or M-CSF (lower panels). Purified CD34<sup>+</sup>CD38<sup>-</sup> HPCs were induced with G-CSF (left) or M-CSF (right) for 0 and 12 hr. Cells were labeled with EdU for 15 min and chased to 1 hr. CAA was performed between nascent DNA (biotin) and EZH2 (upper panels) or UTX (lower panel) antibodies. Red, PLA; green, EdU (biotin); blue, DAPI. Lower panel shows PLA signals only. Graphs represent quantification of the results of CAA experiments shown above by counting the number of PLA signals per EdU-labeled nuclei in 50 cells/each of the three independent experiments. Error bars represent  $\pm$  SD. p Values were determined by ANOVA. \*p < 0.05.

(B) CD34<sup>+</sup>CD38<sup>-</sup> or CD34<sup>+</sup>CD38<sup>+</sup> HPCs were uninduced or induced by treatment with G-CSF or M-CSF. Cells were analyzed by western blotting with antibody to UTX protein.

(C) CD34<sup>+</sup> HPCs were immunostained with antibodies to UTX (red), CD90 (green, indicated by arrows), and EZH2 (blue).

(D) CD34<sup>+</sup> HPCs were pulse-labeled with EdU for 1 hr. CAA was performed for UTX and JMJD3 antibodies and EdU (biotin). Cells were then immunostained for CD90 (green, indicated by arrows) and EdU (blue).

(E) A model for opening of post-replicative chromatin and recruitment of TFs to nascent DNA upon induction of primitive HPCs with lineage-specific cytokines. Primitive (CD34<sup>+</sup>CD38<sup>-</sup> or CD34<sup>+</sup>CD90<sup>+</sup>) HPCs exhibit the H3K27me3 repressive mark on post-replicative chromatin; treatment with instructive cytokines induces H3K27me3 demethylase activity (UTX in the model) and expression of lineage-determining TFs. H3K27me3-unmodified post-replicative chromatin allows binding of lineage-determining TFs to nascent DNA. Upon commitment to unilineage differentiation, H3K27me3 accumulates fast on nascent DNA preventing binding of additional TFs thus restricting transcriptional program and differentiation potential of these cells.



of TFs promoting lineage-specification in mouse and human embryonic stem cells (Petruk et al., 2017 [an accompanying paper in *Molecular Cell*]). Thus, these findings of dynamic changes in chromatin conformation may provide a molecular explanation of the biological plasticity of hematopoietic and embryonic stem cells. Future studies may also provide important insights into more mechanism-based approaches in treatment of hematopoietic disorders including leukemia.

## EXPERIMENTAL PROCEDURES

### Induction of HPCs with Different Cytokines

Human CD34<sup>+</sup>CD38<sup>+</sup> cells were maintained in SFEM (Stem Cell Technologies) enriched with CC100 cocktail of cytokines (SCF, 100 ng/mL; FLT3L, 100 ng/mL; IL-3, 20 ng/mL; IL-6, 20 ng/mL; Stem Cell Technologies) for 12 hr after thawing. Cells were then washed to eliminate the CC100 cytokines and were plated at a concentration of 10<sup>6</sup> cells/mL in SFEM with the addition of either 100 ng/mL of G-CSF (Peprotech) or 10,000 U/mL of M-CSF (R&D Systems) to induce myeloid differentiation, or 10 ng/mL of EPO (Peprotech) to induce erythroid differentiation. The cytokines were re-added at every medium-change.

### Purification of CD34<sup>+</sup>CD38<sup>−</sup> HPCs

Peripheral blood human CD34<sup>+</sup> cells mobilized by G-CSF treatment were sorted 12 hr after thawing on the BD FACS Aria machine, according to the expression of CD38 surface marker. Purified CD34<sup>+</sup>CD38<sup>−</sup> were induced to myeloid lineage with G-CSF or M-CSF and used in CAA and western blotting experiments with antibodies to H3K27me3, EZH2, and UTX.

### Purification of Lin-Sca1<sup>+</sup>Kit<sup>+</sup> Cells

C57BL/6J mice (Jackson Laboratory, stock number 000664) were kept in a pathogen-free facility and euthanized in accordance with the Institutional Animal Care and Use Committee (IACUC) of Thomas Jefferson University. Femoral, tibial, and pelvic bones of 6- to 8-week-old mice were dissected from surrounding muscles and ground in PBS 2% fetal bovine serum (FBS) using a mortar and pestle. Bone marrow cells were filtered through a 40 μm nylon mesh cell strainer (Fisherbrand), and erythrocytes were lysed in 0.8% ammonium chloride (STEMCELL Technologies). Cells were washed twice before staining with PerCP-Cy 5.5 mouse lineage antibody cocktail (BD Pharmingen), PE rat anti-mouse Ly-6A/E (Sca1, BD Pharmingen), and fluorescein isothiocyanate (FITC) rat anti-mouse CD117 (Kit, BD Pharmingen). Lin-Sca1<sup>+</sup>Kit<sup>+</sup> (LSK) cells were sorted using a MoFlo AstriosEQ (Beckman Coulter) and then seeded at 1 × 10<sup>6</sup>/mL in SFEM (STEMCELL Technologies) supplemented with SCF (12.5 μg), FLT3L (12.5 μg), interleukin (IL)-3 (5 μg), and IL-6 (5 μg) (all from Gemini) for 20 hr.

### Analysis of Protein Association with Nascent DNA by CAA

CAA experiments were performed as described before (Petruk et al., 2012, 2013) with small variations. HPCs were pulse-labeled with 5 μM EdU. Approximately 20,000 cells per each condition were harvested and cytopun (750 rpm for 5 min) on polylysine-coated slides (Polyscience), fixed with 4% formaldehyde, permeabilized in 0.25% Triton, and subjected to Click-iT reaction (Invitrogen) to conjugate biotin to EdU. Cells were incubated with mouse antibody to biotin and rabbit antibodies to the protein of interest. The PLA reaction was performed according to the manufacturer (Olink) and as described before. Following PLA, EdU was detected by immunostaining with Alexa Fluor 488-conjugated anti-biotin mouse antibody.

### Re-ChIP with BrdU-Labeled DNA

For each experiment, 8–10 × 10<sup>6</sup> CD34<sup>+</sup> peripheral blood G-CSF-mobilized cells were either not induced or induced for 24 hr with EPO. Undifferentiated and induced cells were labeled with 75 μM BrdU (Sigma-Aldrich) for 15 min or 15 min followed by chase for 2 hr and fixed with 1% formaldehyde for 10 min at room temperature. Chromatin was purified and treated with micrococcal nuclease using Simple ChIP Chromatin IP Kit (Cell Signaling). After treatment, chromatin was immunoprecipitated first with antibody to H3K27me3 (Millipore) or rabbit IgG (Jackson ImmunoResearch). This material was analyzed as con-

ventional ChIP. Following purification, DNA was further immunoprecipitated with antibody to BrdU and used for PCR amplification (re-ChIP). The detailed protocol is described in Francis et al. (2009) and Petruk et al. (2012). All PCR reactions were performed with an Applied Biosystems StepOne Real-Time PCR system. Genes containing H3K27me3 at their promoters (GPR85, WNT16, HOXB8, HOXB13) and a control-activated gene (CD11B) before and after cytokine induction were selected according to (Cui et al., 2009).

### Treatment with Thymidine

To control the efficiency of the thymidine block of DNA replication, CD34<sup>+</sup>CD38<sup>+</sup> cord blood cells were labeled with EdU for 30 min, induced by G-CSF, and grown for 24 hr in the presence of 2 mM thymidine. Following release from thymidine block for 2 or 4 hr, cells were labeled with BrdU to monitor resumption of DNA synthesis. Following “Click-iT” reaction to conjugate biotin to EdU (Petruk et al., 2012), cells were fixed in 4% formaldehyde, denatured in 2N HCL for 30 min for detection of BrdU, and immunostained with antibodies to biotin and BrdU. To perform thymidine block analysis for recruitment of C/EBPα to DNA, nascent DNA of CD34<sup>+</sup>CD38<sup>+</sup> cells was labeled for 30 min with EdU, and then cells were induced with G-CSF and grown for 24 hr in the presence of 2 mM thymidine. Thymidine was removed by three washes with SFEM, cells were grown for the additional 2 or 4 hr and were cytopun on slides, fixed with 4% formaldehyde, and permeabilized in 0.25% Triton. Following “Click-iT” reaction, CAA was performed between nascent DNA (biotin) and C/EBPα as described above.

### Treatment with GSKJ4

CD34<sup>+</sup>CD38<sup>+</sup> cells (cord blood or peripheral blood G-CSF-mobilized) were induced toward myeloid or erythroid differentiation for 12–72 hr in the presence of 5 μM GSKJ4 (Tocris Bioscience) dissolved in DMSO. In control experiments, cells were incubated with the same concentration of DMSO without GSKJ4. Erythroid differentiation was monitored after 72 hr by flow cytometry analysis detection of CD36 and glycophorin A/GYPA/CD235a double-positive cells. Expression of GATA-1-regulated genes was analyzed at 48 hr by real-time PCR with the following primers: GATA-1 forward 5'-GCTACACCGA GTGAACCGGC-3', reverse 5'-CAGACTGGAGCCCCGTTTCT-3'; ALAS2 forward 5'-TTCAAGACTGTGAACCGCTG-3', reverse 5'-GCTCATGCCAGGT AATCAT-3', GYPA forward 5'-GGGTGATGGCTGGTGTATTGG-3', reverse 5'-AAAGGCACGTCTGTGTGAGG-3'; and ABCB10 forward 5'-CGGAGGC CCGGAAGCTC-3', reverse 5'-TTCCCCAGGAAGAAAGGGGC-3'.

Granulocytic differentiation was monitored by flow cytometry analysis of CD33 and CD15 levels (at 72 hr post-G-CSF treatment) and by expression of C/EBPα-regulated genes (at 72 hr post G-CSF treatment) detected by real-time PCR with the following primers: CEBPE (C/EBPε) forward 5'-CCAGCCGAGGCAGCTACAAT-3', reverse 5'-CAAAGGGCCCTTGAGAAC GC-3'; MPO (myeloperoxidase) forward 5'-TCTCCTCACCAACCGCTCAG-3', reverse 5'-CCGGTTGTGCTCCCGAAGTA-3'; ELANE (neutrophilic elastase) forward 5'-CTGGGAGCCCATAACCTCTCG-3', reverse 5'-CGTTGAGCTGG AGAATCACGA-3'; and S100A8 (calgranulin A) forward 5'-ATATCAGGAAA AAGGGTGACAGACG-3', reverse 5'-GCTGCCACGCCATCTTTAT-3'. Real-time PCR data were normalized for expression of GAPDH detected with the following primers forward 5'-CCCATCCCATCTTCCAGGAG-3' and reverse 5'-CTTCTCCATGGTGGTGAAGACG-3'.

### Antibodies

Anti-human CD34-PE (Miltenyi Biotec #130-081-002) and anti-human CD38-APC (Miltenyi Biotec #130-092-261) were used to analyze the percentage of CD34<sup>+</sup> and CD38<sup>+</sup> populations. Anti-human CD15-PE-Cy7 (BD Pharmingen #560827), CD33-PE (BD Pharmingen #555450), CD11b-APC (BD Pharmingen #561015), and CD14-PE (BD Pharmingen #555398) were used to test the percentage of cells positive for granulocytic and macrophage differentiation markers. CD235a-PE-Cy5 (BD Pharmingen #561776) and CD36-PE (BD Pharmingen #561821) were used to assess erythroid differentiation. Cells were incubated with PBS + 1% BSA to prevent antibodies' non-specific binding and then stained with fluorochrome-conjugated antibodies for 15 min at room temperature. Cytofluorimetric analysis was performed on LSRII flow cytometer (BD Bioscience, USA). For CAA and immunostaining, the following antibodies were used: anti-C/EBP (1:200, Abcam); anti-PU.1 (1:100, Santa



Cruz); anti-GATA-1 (1:400, Cell Signaling); anti-H3K27me3 (1:2500, Millipore); mouse anti-biotin (1:1,000, Jackson ImmunoResearch); UTX (1:400, Thermo Fisher Scientific); EZH2 (1:400, Cell Signaling); JMJD3 (1:200, Abgent); BrdU (1:200, BD Bioscience); and CD90 (1:100, R&D Systems).

### Statistical Analysis

For CAA experiments and re-ChIP experiment, p values were calculated by a one-way ANOVA with Bonferroni's post hoc test. The statistical significance was defined as  $p < 0.05$ .

### SUPPLEMENTAL INFORMATION

Supplemental Information includes five figures and can be found with this article online at <http://dx.doi.org/10.1016/j.celrep.2017.03.035>.

### AUTHOR CONTRIBUTIONS

A.M., S.P., and B.C. designed the study and wrote the manuscript. S.P. performed the PLA studies. S.A.M. and M.D.D. performed purification and assessed immunophenotype and gene expression of hematopoietic progenitor subsets. P.P. performed differentiation assays of GSKJ4-treated CD34<sup>+</sup> cells. V.M. and M.D.D. performed purification of mouse HPCs. N.F. provided mobilized CD34<sup>+</sup> cells. J.C. and L.I. discussed the results and designed experiments. All authors commented on the manuscript.

### ACKNOWLEDGMENTS

This work was supported by the NIH (R01GM075141 to A.M., R01HL127895 to A.M. and B.C., R01AI125650 to A.M. and L.I., and NS075839 to L.I.) and the Parkinson's Council and The Strauss Foundation (to L.I.).

Received: June 14, 2016

Revised: December 28, 2016

Accepted: March 9, 2017

Published: April 11, 2017

### REFERENCES

- Bell, O., Schwaiger, M., Oakeley, E.J., Lienert, F., Beisel, C., Stadler, M.B., and Schübeler, D. (2010). Accessibility of the *Drosophila* genome discriminates PcG repression, H4K16 acetylation and replication timing. *Nat. Struct. Mol. Biol.* **17**, 894–900.
- Cantor, A.B., and Orkin, S.H. (2001). Hematopoietic development: a balancing act. *Curr. Opin. Genet. Dev.* **11**, 513–519.
- Chang, H.H., Hemberg, M., Barahona, M., Ingber, D.E., and Huang, S. (2008). Transcriptome-wide noise controls lineage choice in mammalian progenitor cells. *Nature* **453**, 544–547.
- Cui, K., Zang, C., Roh, T.Y., Schones, D.E., Childs, R.W., Peng, W., and Zhao, K. (2009). Chromatin signatures in multipotent human hematopoietic stem cells indicate the fate of bivalent genes during differentiation. *Cell Stem Cell* **4**, 80–93.
- Dahl, R., Walsh, J.C., Lancki, D., Laslo, P., Iyer, S.R., Singh, H., and Simon, M.C. (2003). Regulation of macrophage and neutrophil cell fates by the PU.1:C/EBP $\alpha$  ratio and granulocyte colony-stimulating factor. *Nat. Immunol.* **4**, 1029–1036.
- Di Stefano, B., Sardina, J.L., van Oevelen, C., Collombet, S., Kallin, E.M., Vicent, G.P., Lu, J., Thieffry, D., Beato, M., and Graf, T. (2014). C/EBP $\alpha$  poises B cells for rapid reprogramming into induced pluripotent stem cells. *Nature* **506**, 235–239.
- Feng, R., Desbordes, S.C., Xie, H., Tillo, E.S., Pixley, F., Stanley, E.R., and Graf, T. (2008). PU.1 and C/EBP $\alpha$ /beta convert fibroblasts into macrophage-like cells. *Proc. Natl. Acad. Sci. USA* **105**, 6057–6062.
- Filipczyk, A.A., Laslett, A.L., Mummery, C., and Pera, M.F. (2007). Differentiation is coupled to changes in the cell cycle regulatory apparatus of human embryonic stem cells. *Stem Cell Res. (Amst.)* **1**, 45–60.

Francis, N.J., Follmer, N.E., Simon, M.D., Aghia, G., and Butler, J.D. (2009). Polycomb proteins remain bound to chromatin and DNA during DNA replication in vitro. *Cell* **137**, 110–122.

Friedman, A.D. (2007). Transcriptional control of granulocyte and monocyte development. *Oncogene* **26**, 6816–6828.

Graf, T., and Enver, T. (2009). Forcing cells to change lineages. *Nature* **462**, 587–594.

Grover, A., Mancini, E., Moore, S., Mead, A.J., Atkinson, D., Rasmussen, K.D., O'Carroll, D., Jacobsen, S.E., and Nerlov, C. (2014). Erythropoietin guides multipotent hematopoietic progenitor cells toward an erythroid fate. *J. Exp. Med.* **211**, 181–188.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576–589.

Iida, S., Watanabe-Fukunaga, R., Nagata, S., and Fukunaga, R. (2008). Essential role of C/EBP $\alpha$  in G-CSF-induced transcriptional activation and chromatin modification of myeloid-specific genes. *Genes Cells* **13**, 313–327.

Iwafuchi-Doi, M., and Zaret, K.S. (2014). Pioneer transcription factors in cell reprogramming. *Genes Dev.* **28**, 2679–2692.

Iwafuchi-Doi, M., and Zaret, K.S. (2016). Cell fate control by pioneer transcription factors. *Development* **143**, 1833–1837.

Iwasaki, H., Mizuno, S., Arinobu, Y., Ozawa, H., Mori, Y., Shigematsu, H., Takatsu, K., Tenen, D.G., and Akashi, K. (2006). The order of expression of transcription factors directs hierarchical specification of hematopoietic lineages. *Genes Dev.* **20**, 3010–3021.

Jonk, L.J., de Jonge, M.E., Kruyt, F.A., Mummery, C.L., van der Saag, P.T., and Kruijer, W. (1992). Aggregation and cell cycle dependent retinoic acid receptor mRNA expression in P19 embryonal carcinoma cells. *Mech. Dev.* **36**, 165–172.

Kapinas, K., Grandy, R., Ghule, P., Medina, R., Becker, K., Pardee, A., Zaidi, S.K., Lian, J., Stein, J., van Wijnen, A., and Stein, G. (2013). The abbreviated pluripotent cell cycle. *J. Cell. Physiol.* **228**, 9–20.

Kruidenier, L., Chung, C.W., Cheng, Z., Liddle, J., Che, K., Joberty, G., Bantscheff, M., Bountra, C., Bridges, A., Diallo, H., et al. (2012). A selective jumoni H3K27 demethylase inhibitor modulates the proinflammatory macrophage response. *Nature* **488**, 404–408.

Kulesa, H., Frampton, J., and Graf, T. (1995). GATA-1 reprograms avian myelomonocytic cell lines into eosinophils, thromboplasts, and erythroblasts. *Genes Dev.* **9**, 1250–1262.

Lara-Astiaso, D., Weiner, A., Lorenzo-Vivas, E., Zaretzky, I., Jaitin, D.A., David, E., Keren-Shaul, H., Mildner, A., Winter, D., Jung, S., et al. (2014). Immunogenetics. Chromatin state dynamics during blood formation. *Science* **345**, 943–949.

Laslo, P., Spooner, C.J., Warmflash, A., Lancki, D.W., Lee, H.J., Sciammas, R., Gantner, B.N., Dinner, A.R., and Singh, H. (2006). Multilineage transcriptional priming and determination of alternate hematopoietic cell fates. *Cell* **126**, 755–766.

Li, V.C., Ballabeni, A., and Kirschner, M.W. (2012). Gap 1 phase length and mouse embryonic stem cell self-renewal. *Proc. Natl. Acad. Sci. USA* **109**, 12550–12555.

Mercer, E.M., Lin, Y.C., Benner, C., Jhunjhunwala, S., Dutkowski, J., Flores, M., Sigvardsson, M., Ideker, T., Glass, C.K., and Murre, C. (2011). Multilineage priming of enhancer repertoires precedes commitment to the B and myeloid cell lineages in hematopoietic progenitors. *Immunity* **35**, 413–425.

Mossadegh-Keller, N., Sarrazin, S., Kandalla, P.K., Espinosa, L., Stanley, E.R., Nutt, S.L., Moore, J., and Sieweke, M.H. (2013). M-CSF instructs myeloid lineage fate in single haematopoietic stem cells. *Nature* **497**, 239–243.

Mummery, C.L., van den Brink, C.E., and de Laat, S.W. (1987). Commitment to differentiation induced by retinoic acid in P19 embryonal carcinoma cells is cell cycle dependent. *Dev. Biol.* **121**, 10–19.

Murray, A.W., and Kirschner, M.W. (1989). Dominoes and clocks: the union of two views of the cell cycle. *Science* **246**, 614–621.

- Notta, F., Zandi, S., Takayama, N., Dobson, S., Gan, O.I., Wilson, G., Kaufmann, K.B., McLeod, J., Laurenti, E., Dunant, C.F., et al. (2016). Distinct routes of lineage development reshape the human blood hierarchy across ontogeny. *Science* *357*, aab2116.
- Orkin, S.H. (2000). Diversification of haematopoietic stem cells to specific lineages. *Nat. Rev. Genet.* *1*, 57–64.
- Orkin, S.H., and Zon, L.I. (2008). Hematopoiesis: an evolving paradigm for stem cell biology. *Cell* *132*, 631–644.
- Paul, F., Arkin, Y., Giladi, A., Jaitin, D.A., Kenigsberg, E., Keren-Shaul, H., Winter, D., Lara-Astiaso, D., Gury, M., Weiner, A., et al. (2015). Transcriptional heterogeneity and lineage commitment in myeloid progenitors. *Cell* *163*, 1663–1677.
- Perié, L., Duffy, K.R., Kok, L., de Boer, R.J., and Schumacher, T.N. (2015). The branching point in erythro-myeloid differentiation. *Cell* *163*, 1655–1662.
- Petruk, S., Sedkov, Y., Johnston, D.M., Hodgson, J.W., Black, K.L., Kovermann, S.K., Beck, S., Canaani, E., Brock, H.W., and Mazo, A. (2012). TrxG and PcG proteins but not methylated histones remain associated with DNA through replication. *Cell* *150*, 922–933.
- Petruk, S., Black, K.L., Kovermann, S.K., Brock, H.W., and Mazo, A. (2013). Stepwise histone modifications are mediated by multiple enzymes that rapidly associate with nascent DNA during replication. *Nat. Commun.* *4*, 2841.
- Petruk, S., Cai, J., Sussman, R., Sun, G., Kovermann, S.K., Mariani, S.K., Calabretta, B., McMahon, S.B., Brock, H.W., Iacovitti, L., and Mazo, A. (2017). Delayed accumulation of H3K27me3 on nascent DNA is essential for recruitment of transcription factors at early stages of stem cell differentiation. *Mol. Cell* *66*, Published online April 20, 2016. <http://dx.doi.org/10.1016/j.molcel.2017.03.006>.
- Rieger, M.A., Hoppe, P.S., Smejkal, B.M., Eitelhuber, A.C., and Schroeder, T. (2009). Hematopoietic cytokines can instruct lineage choice. *Science* *325*, 217–218.
- Shlyueva, D., Stampfel, G., and Stark, A. (2014). Transcriptional enhancers: from properties to genome-wide predictions. *Nat. Rev. Genet.* *15*, 272–286.
- Singh, A.M., and Dalton, S. (2009). The cell cycle and Myc intersect with mechanisms that regulate pluripotency and reprogramming. *Cell Stem Cell* *5*, 141–149.
- Thieme, S., Gyárfás, T., Richter, C., Özhan, G., Fu, J., Alexopoulou, D., Muders, M.H., Michalk, I., Jakob, C., Dahl, A., et al. (2013). The histone demethylase UTX regulates stem cell migration and hematopoiesis. *Blood* *121*, 2462–2473.
- Till, J.E., McCulloch, E.A., and Siminovitch, L. (1964). A stochastic model of stem cell proliferation, based on the growth of spleen colony-forming cells. *Proc. Natl. Acad. Sci. USA* *51*, 29–36.
- Wei, G., Wei, L., Zhu, J., Zang, C., Hu-Li, J., Yao, Z., Cui, K., Kanno, Y., Roh, T.Y., Watford, W.T., et al. (2009). Global mapping of H3K4me3 and H3K27me3 reveals specificity and plasticity in lineage fate determination of differentiating CD4+ T cells. *Immunity* *30*, 155–167.
- Welch, J.J., Watts, J.A., Vakoc, C.R., Yao, Y., Wang, H., Hardison, R.C., Blobel, G.A., Chodosh, L.A., and Weiss, M.J. (2004). Global regulation of erythroid gene expression by transcription factor GATA-1. *Blood* *104*, 3136–3147.
- Xie, H., Ye, M., Feng, R., and Graf, T. (2004). Stepwise reprogramming of B cells into macrophages. *Cell* *117*, 663–676.
- Yuan, W., Wu, T., Fu, H., Dai, C., Wu, H., Liu, N., Li, X., Xu, M., Zhang, Z., Niu, T., et al. (2012). Dense chromatin activates Polycomb repressive complex 2 to regulate H3 lysine 27 methylation. *Science* *337*, 971–975.
- Zaret, K.S., and Carroll, J.S. (2011). Pioneer transcription factors: establishing competence for gene expression. *Genes Dev.* *25*, 2227–2241.