

3-19-2022


Pharmacologic Ascorbic Acid as Early Therapy for Hospitalized Patients with COVID-19: A Randomized Clinical Trial

Dagan Coppock
Thomas Jefferson University

Pierre-Christian Violet
National Institutes of Health

Gustavo Vasquez
Thomas Jefferson University

Katherine Belden
Thomas Jefferson University
Follow this and additional works at: <https://jdc.jefferson.edu/petfp>

 [Digital Commons](#), [Pharmaceutical Sciences Commons](#), [Infectious Disease Commons](#), [Integrative Medicine Commons](#), and the [Pharmacy and Pharmaceutical Sciences Commons](#) at [Thomas Jefferson University](#)

[Let us know how access to this document benefits you](#)

~~See next page for additional authors~~

Recommended Citation

Coppock, Dagan; Violet, Pierre-Christian; Vasquez, Gustavo; Belden, Katherine; Foster, Michael; Mullin, Bret; Magee, Devon; Mikell, Isabelle; Shah, Lokesh; Powers, Victoria; Curcio, Brian; Monti, Daniel A.; and Levine, Mark, "Pharmacologic Ascorbic Acid as Early Therapy for Hospitalized Patients with COVID-19: A Randomized Clinical Trial" (2022). *Department of Pharmacology and Experimental Therapeutics Faculty Papers*. Paper 143.
<https://jdc.jefferson.edu/petfp/143>


This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Pharmacology and Experimental Therapeutics Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Dagan Coppock, Pierre-Christian Violet, Gustavo Vasquez, Katherine Belden, Michael Foster, Bret Mullin, Devon Magee, Isabelle Mikell, Lokesh Shah, Victoria Powers, Brian Curcio, Daniel A. Monti, and Mark Levine

Article

Pharmacologic Ascorbic Acid as Early Therapy for Hospitalized Patients with COVID-19: A Randomized Clinical Trial

Dagan Coppock ^{1,*}, Pierre-Christian Violet ², Gustavo Vasquez ¹, Katherine Belden ¹, Michael Foster ³, Bret Mullin ³, Devon Magee ³, Isabelle Mikell ⁴, Lokesh Shah ⁴, Victoria Powers ⁴, Brian Curcio ⁵, Daniel Monti ⁶ and Mark Levine ^{2,*}

¹ Division of Infectious Diseases, Department of Medicine, Sidney Kimmel Medical College, Thomas Jefferson University, 1015 Chestnut Street, Philadelphia, PA 19107, USA; gvasquezmd@gmail.com (G.V.); katherine.belden@jefferson.edu (K.B.)

² Molecular and Clinical Nutrition Section, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, 10 Center Drive, Bethesda, MD 20892, USA; pierre-christian.violet@nih.gov

³ Jefferson Clinical Research Institute, Thomas Jefferson University, 833 Chestnut Street, Philadelphia, PA 19107, USA; mfooster610@gmail.com (M.F.); bret.mullin@jefferson.edu (B.M.); devon.magee@jefferson.edu (D.M.)

⁴ Sidney Kimmel Medical College, Thomas Jefferson University, 1015 Walnut Street, Philadelphia, PA 19107, USA; isabelle.mikell@students.jefferson.edu (I.M.); lokesh.shah@students.jefferson.edu (L.S.); tori.powers@students.jefferson.edu (V.P.)

⁵ Division of Biostatistics, Department of Pharmacology and Experimental Therapeutics, Sidney Kimmel Medical College, Thomas Jefferson University, 1015 Chestnut Street, Philadelphia, PA 19107, USA; curciob@arcadia.edu

⁶ Department of Integrative Medicine and Nutritional Sciences, Sidney Kimmel Medical College, Thomas Jefferson University, 925 Chestnut Street, Philadelphia, PA 19107, USA; daniel.monti@jefferson.edu

* Correspondence: dagan.coppock@jefferson.edu (D.C.); markl@bdg8.niddk.nih.gov (M.L.)



Citation: Coppock, D.; Violet, P.-C.; Vasquez, G.; Belden, K.; Foster, M.; Mullin, B.; Magee, D.; Mikell, I.; Shah, L.; Powers, V.; et al. Pharmacologic Ascorbic Acid as Early Therapy for Hospitalized Patients with COVID-19: A Randomized Clinical Trial. *Life* **2022**, *12*, 453. <https://doi.org/10.3390/life12030453>

Academic Editor: Ramón Cabellos

Received: 29 January 2022

Accepted: 15 March 2022

Published: 19 March 2022

Corrected: 31 August 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Despite the widespread availability of effective vaccines, new cases of infection with severe acute respiratory syndrome coronavirus-2, the cause of coronavirus disease 2019 (COVID-19), remain a concern in the settings of vaccine hesitancy and vaccine breakthrough. In this randomized, controlled, phase 2 trial, we hypothesized that high-dose ascorbic acid delivered intravenously to achieve pharmacologic concentrations may target the high viral phase of COVID-19 and thus improve early clinical outcomes. Sixty-six patients admitted with COVID-19 and requiring supplemental oxygen were randomized to receive either escalating doses of intravenous ascorbic acid plus standard of care or standard of care alone. The demographic and clinical characteristics were well-balanced between the two study arms. The primary outcome evaluated in this study was clinical improvement at 72 h after randomization. While the primary outcome was not achieved, point estimates for the composite outcome and its individual components of decreased use of supplemental oxygen, decreased use of bronchodilators, and the time to discharge were all favorable for the treatment arm. Possible favorable effects of ascorbic acid were most apparent during the first 72 h of hospitalization, although these effects disappeared over the course of the entire hospitalization. Future larger trials of intravenous ascorbic acid should be based on our current understanding of COVID-19 with a focus on the potential early benefits of ascorbic in hospitalized patients.

Keywords: SARS-CoV-2; COVID-19; vitamin C; ascorbic acid

1. Introduction

In the United States, infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has led to an estimated 6.2 million hospitalizations and 767,000 deaths due to coronavirus disease 2019 (COVID-19) [1]. Despite the widespread availability of effective messenger ribonucleic acid vaccines, new infections remain a concern in the settings of vaccine hesitancy and vaccine breakthrough [2,3]. Furthermore, as new variants of SARS-CoV-2 emerge, such as the Delta variant, there is the risk of infection despite successful

vaccination [4,5]. In these contexts, the development of new treatments for COVID-19 remains a priority.

COVID-19 pneumonia appears to be a two-phase disease, in which the first phase is dominated by a high viral load in the airways, followed by a later phase of low or undetectable virus and progressive organizing fibrosis [6]. We hypothesized that high-dose ascorbic acid (AA) delivered intravenously to achieve pharmacologic concentrations may target that early high viral phase through two distinct potential mechanisms: Lymphocyte activation and/or inhibition of early viral replication [7–12]. Both mechanisms are linked to hydrogen peroxide formation, for which pharmacologic AA concentrations serve as a prodrug [13–16]. When given intravenously, but not orally, pharmacologic AA concentrations increase the steady-state formation of hydrogen peroxide in vivo over hours in extracellular fluid but not in the blood [14,15].

To translate potential pharmacologic AA efficacy to acute COVID-19 treatment, we chose primary and secondary clinical endpoints that reflected these postulated mechanisms of action. These endpoints included clinical improvement within 72 h of pharmacologic AA treatment, including hospital discharge, the need for oxygen, the use of bronchodilators, days of fever, and clinical decline or serious adverse events. Here we present an exploratory, single-center 2:1 randomized open-label clinical study evaluating these endpoints in 66 people with acute COVID-19 admitted to one hospital system in the United States.

2. Methods

2.1. Study Design and Patients

This study was approved by the Thomas Jefferson University Office of Human Research Institutional Review Board (Protocol #20D.431). It was conducted in accordance with Federal-Wide Assurance #00002109 to the U.S. Department of Health and Human Services. Written informed consent was obtained from all patients. The study was registered at ClinicalTrials.gov (Identifier: NCT04363216). This prospective, randomized, open-label, phase II clinical trial was conducted at Thomas Jefferson University Hospital and Jefferson Methodist Hospital from 8 May 2020 through 31 December 2020. Potential study participants were identified based upon hospital admission with a new diagnosis of SARS-CoV-2. Subjects were screened by study personnel as soon after admission as feasible. Subjects were considered eligible for inclusion if they were male or a non-pregnant female, were greater than 18 years of age, were confirmed positive for SARS-CoV-2 infection by nasal swab, and required supplemental oxygen without an anticipated need for mechanical ventilation within 24 h. Patients were excluded if they had an estimated glomerular filtration rate less than 50 at the time of screening, known glucose-6-phosphate dehydrogenase deficiency, an anticipated need for mechanical ventilation within 24 h, a home oxygen requirement, known allergy to AA, or if they were enrolled in another COVID-19 treatment study at the time of enrollment. Patients who met screening requirements were approached for the study after discussion with their primary inpatient team (see Table S1 for CONSORT checklist).

2.2. Randomization

Once written consent was obtained, patients were randomized, using the REDCap Randomization Module, in a 2:1 ratio into either a treatment arm that received standard of care (SoC) plus AA or a control arm that received SoC alone. Randomization was stratified by risk of complications (high vs. low). Patients were considered high risk if they were older than 60 or had any of the following comorbidities: Hypertension, structural lung disease, cardiovascular disease, diabetes, immunocompromising conditions, or conditions requiring immunosuppressing medications.

2.3. Interventions

If randomized to the treatment arm, patients received routine clinical care plus up to 6 escalating doses of daily AA infusions in magnesium chloride to reduce the sensation of

burning on injection. On day zero (the randomization day), patients received AA 0.3 g/kg intravenously once via a continuous 2-h infusion. On day one, patients received AA 0.6 g/kg intravenously once via continuous infusion. Thereafter, patients received 0.9 g/kg intravenously once daily through day five. Patients received these infusions while they were admitted. If patients were discharged prior to the final (day five) infusion, AA was discontinued at the time of discharge. If randomized to the control arm, patients received routine clinical care.

2.4. Procedures

While enrolled in the study, patients were continuously monitored by the study coordinator and investigators for adverse and serious adverse events. Patients were withdrawn if they progressed to a requirement of high-flow oxygen, were transferred to the intensive care unit, were intubated prior to completion of the study procedures, or if they developed any clinical condition that raised concerns for safety. An interim safety analysis was performed after the first 21 patients who were randomized to the treatment arm completed the course of AA.

2.5. Outcome Measures

The primary endpoint was a composite outcome—clinical improvement at 72 h post-randomization, defined as a 50% reduction in the highest flow rate of oxygen during the 72-h period, a 50% reduction in the most frequent use of bronchodilators within a 12-h window within the 72-h period, or hospital discharge (whichever came first).

The secondary endpoints analyzed included clinical decline within 36 h of randomization, defined as requiring ICU-level care; the rate of oxygen supplementation given, defined by the days to a 50% reduction in supplementation during the hospital course; any day of fever as defined by a temperature of greater than 100.4 °F; the time to discharge; and serious adverse events specific to the treatment. Adverse and serious adverse events were defined based upon guidelines set forth on [clinicaltrials.gov](https://www.clinicaltrials.gov) [17].

2.6. Statistical Analyses

All analyses were prespecified in the trial protocol and followed the intention-to-treat principle, i.e., included all subjects as randomized. The protocol specified interim analyses based on the proportion of clinical decline among the first 21 patients in the AA group. The stopping rule specified that the trial would continue only if fewer than 5 of the first 21 patients experienced clinical decline requiring high-flow oxygen and/or ICU-level care; this corresponds to the one-sided upper 95% exact binomial confidence limit excluding the unacceptably high rate of 40%.

The main endpoint (clinical improvement within 72 h from randomization) was analyzed with a stratified Mantel–Haenszel procedure, with the risk of complication as the stratification factor. The comparison of the two trial groups was based on a 1-sided exact test with alpha 0.05; the corresponding 90% exact confidence interval was also computed. The dichotomous secondary endpoints of any fever and adverse events were analyzed similarly. The time to a 50% decrease in oxygen supplementation and time to discharge were analyzed with the Kaplan–Meier method and Cox proportional hazards regression. A secondary analysis of the time to discharge was performed, also on an intention-to-treat basis, with adjustments made for admission-to-randomization time.

2.7. Sample Size and Power

With a target sample size of 66 patients (44 in the AA group and 22 in the SoC group), the trial had 82% power to detect a difference of 60% vs. 25% clinical improvement in the AA vs. the SoC arm, using the one-sided Fisher's exact test with alpha 0.05.

3. Results

3.1. Patients

Of the 101 patients approached, seventy consented to the study and were randomized. Of those seventy, four were excluded prior to analysis (Figure 1). This included one patient withdrawal and three exclusions due to clinical deterioration between consent and randomization. Of the remaining sixty-six patients, forty-four were randomized to receive AA plus SoC and twenty-two were randomized to receive SoC alone.




Figure 1. CONSORT flow diagram.

In regard to demographics and baseline comorbidities, patients were generally balanced between the treatment and control arms (Table 1). The mean age in the treatment arm was 60 years compared to 61 years in the control arm. Fifty percent of the population

was female in both groups. At the time of randomization, the mean body mass index was 33.5 in the treatment arm and 35 in the SoC arm.

Table 1. Demographic and clinical characteristics of patients in the treatment and control groups.

	Number (%)	
	Standard of Care Alone (N = 22)	Ascorbic Acid Plus Standard of Care (N = 44)
Characteristic		
Mean age (SD), years	61 (11)	60 (17)
Age groups, years		
<50	3 (13.6)	13 (29.5)
50–59	5 (22.7)	7 (15.9)
60–69	10 (45.5)	13 (29.5)
70+	4 (18.2)	11 (25.0)
Sex		
Female	11 (50.0)	22 (50.0)
Male	11 (50.0)	22 (50.0)
Race/Ethnicity		
Caucasian/White, Non-Hispanic	14 (63.6)	23 (52.3)
African American/Black, Non-Hispanic	4 (18.2)	17 (38.6)
Asian, Non-Hispanic	1 (4.5)	2 (4.5)
Hispanic	3 (13.6)	2 (4.5)
Mean BMI (SD), kg/m ²	35.0 (8.7)	33.5 (8.8)
BMI groups, kg/m ²		
Normal weight (18.5–24.9)	2 (9.1)	6 (13.6)
Overweight (25–29.9)	5 (22.7)	11 (25.0)
Obese (30+)	15 (68.2)	27 (61.4)
Smoking status		
Never smoker	12 (57.1)	23 (52.3)
Former smoker	9 (42.9)	17 (38.6)
Current smoker	0 (0.0)	4 (9.1)
Comorbid conditions		
Diabetes	8 (36.4)	15 (34.9)
Cardiovascular disease	12 (54.5)	29 (65.9)
Chronic obstructive pulmonary disease	5 (22.7)	8 (18.6)
Organ transplant recipient	1 (4.5)	0 (0.0)
Risk of complications		
Low	4 (18.2)	9 (20.5)
High	18 (81.8)	35 (79.5)
Mean time since the onset of symptoms (SD), days	7.2 (3.9)	8.1 (3.6)
Medications administered after randomization		
Antibiotics	2 (9.0)	9 (20.5)
Remdesivir	20 (90.9)	41 (93.2)
Glucocorticoids	18 (81.8)	33 (75.0)
NSAIDs	11 (50.0)	8 (18.1)

BMI, body mass index; kg/m², kilogram per meter squared; NSAIDs, non-steroidal anti-inflammatory drugs; SD, standard deviation.

The mean time from COVID-19-related symptom onset to randomization was 8.1 days (standard deviation 3.6) in the treatment arm and 7.2 days (standard deviation 3.9) in the SoC arm. The baseline oxygen requirement was higher in the SoC arm compared to

the treatment arm (3 L/min vs. 2.5 L/min, respectively). During the peri-randomization period, six patients were febrile in the SoC arm, while only one patient was febrile in the treatment arm. Although baseline clinical characteristics were defined by randomization, the time from admission to randomization was noted to be longer in the AA group—1.9 days vs. 1.3 days, respectively. Though there were notable differences in antibiotic and non-steroidal anti-inflammatory, the current SoC for COVID-19, which includes remdesivir and corticosteroids, was balanced between the two study arms.

3.2. Interim Analyses

Among the first 21 patients in the AA group, only 2 (9.5%) needed high-flow oxygen and ICU-level care (upper 95% confidence limit: 27.1%). Therefore, the trial continued to its full accrual.

3.3. Primary Outcome

The study's primary outcome was clinical improvement within 72 h of randomization, based on three components: 50% reduction in oxygen flow rate, 50% reduction in bronchodilator use, or discharge. The results are summarized in Table 2. The point estimate for overall clinical improvement, as defined by the primary outcome, was larger in the AA arm compared to the SoC arm (86% vs. 73%; odds ratio, OR = 2.36; 90% confidence interval, CI: 0.66 to 8.07), though the difference was not statistically significant (1-sided $p = 0.158$). When adjusted for admission-to-randomization time, the odds ratio for clinical improvement with treatment was 2.00 (90% CI, 0.57 to 7.00; 1-sided $p = 0.219$). Furthermore, time of initiation of the AA treatment did not substantially impact the results; clinical improvement within 72 h was noted in 19 of the 23 (83%) treatment-arm patients who received their first infusion within the first 3 h vs. 19 of the 21 (90%) who received their first infusion later.

Table 2. Primary endpoint.

	Standard of Care Alone (<i>N</i> = 22)	Ascorbic Acid Plus Standard of Care (<i>N</i> = 44)	Odds Ratio (90% Confidence Interval)	<i>p</i> -Value
Number of patients who achieved clinical improvement within 72 h of randomization (<i>N</i>)	16 (72.7%)	38 (86.4%)	2.36 (0.66, 8.07)	0.158
Number of patients who achieved a 50% reduction in supplemental oxygen (<i>N</i>)	15 (68.2%)	34 (77.3%)		
Number of patients who achieved a 50% reduction in bronchodilator use (<i>N</i>)	1 (4.5%)	3 (6.8%)		
Number of patients discharged (<i>N</i>)	2 (9.1%)	13 (29.5%)		

Regarding the composite endpoint's components, a 50% decrease in supplemental oxygen flow rate was observed in 77.3% of individuals in the AA arm compared to 68.2% of those in the SoC arm. Less than a third of the patients in both groups were using bronchodilators; among those who were, a 50% reduction in use was noted in 6.8% of the AA arm vs. 4.5% of the SoC arm. Finally, discharge within 72 h of randomization was 29.5% in the AA arm vs. 9.1% in the SoC arm.

3.4. Secondary Outcomes

Table 3 summarizes the results for the secondary endpoints of fever and adverse events. Fever was less common in the AA group compared to the SoC group (14% vs. 36%; $p = 0.038$). However, more patients in the SoC group than in the AA group had fever at enrollment; when those patients were excluded, the difference between the groups was attenuated (14% vs. 25%; $p = 0.263$). Adverse event rates were higher in the AA arm than in the SoC arm, but the difference was not significant (any adverse event: 55% vs. 32%; $p = 0.133$; serious adverse events: 18% vs. 9%; $p = 0.293$). The serious adverse events included death (one SoC), hypothermia (one AA), hypoxia (six AA, two SoC), leukocytosis (one AA), pulmonary edema (two AA), and sepsis (two AA). There were no reported study-related deaths in either arm.

Table 3. Secondary endpoints.

	Standard of Care Alone (N = 22)	Ascorbic Acid Plus Standard of Care (N = 44)	Hazard Ratio (90% Confidence Interval)	p-Value
Time to 50% reduction in supplemental oxygen (days)	2.24	1.87	1.13 (0.65, 1.97)	0.334
Time to discharge (days)	4.65	4.3	1.03 (0.61, 1.76)	0.453
Number of patients with any fever (N)	8 (36.4%)	6 (13.6%)	0.28 (0.09, 0.93)	0.038
Number of patients with any fever, excluding patients with fever at randomization (N)	4 (25.0%)	6 (14.0%)	0.50 (0.13, 2.15)	0.263
Any serious adverse event (N)	2 (9.1%)	8 (18.2%)	2.23 (0.46, 15.5)	0.293
Clinical decline within 36 h of randomization (N)	2 (9.1%)	4 (9.1%)	0.95 (0.16, 7.84)	0.643

For the overall study period, the time to a 50% reduction in supplemental oxygenation was not statistically significant between the AA and SoC arms (median = 1.9 days vs. 2.2 days; hazard ratio (HR) 1.13 [90% CI, 0.65 to 1.97]; $p = 0.334$) (Figure 2). The time to discharge was also not statistically different between the two groups (median = 4.3 days vs. 4.6 days; HR 1.03 [CI 90%, 0.61 to 1.76]; $p = 0.453$) (Figure 2). Adjustment for the admission-to-randomization time did not materially change these results; the estimated hazard ratios were 1.05 (90% CI, 0.60 to 1.85; 1-sided $p = 0.432$) for the time to a 50% reduction in oxygen supplementation and 1.03 (90% CI: 0.61 to 1.76; 1-sided $p = 0.453$) for the time to discharge. Differences in the cumulative proportion curves for a reduction in supplemental oxygen and discharge were more pronounced in the first 72 h of the study (Figure 2, insets). Regarding protocol deviations, as noted in Figure 1, three patients (1 AA, 2 SoC) were incorrectly excluded from the study shortly after consent and randomization due to clinical deterioration. Notes to file were included in trial documentation.




Figure 2. Time to a 50% reduction in oxygen supplementation and time to discharge in the study arms.

4. Discussion

In this unblinded, randomized exploratory trial of hospitalized adults with COVID-19, intravenous AA plus SoC vs. SoC alone was provided to individuals admitted to a general medical ward with positive COVID-19 tests who required supplemental oxygen. Based on our primary outcomes, there was no statistical difference between those patients who received AA compared to those who received SoC for the primary outcome of clinical improvement within 72 h of randomization.

The scientific premise of this trial is that AA administered intravenously to produce pharmacologic concentrations might be an effective treatment agent if given early in the clinical course of COVID-19 and at sufficient doses. As has been previously described, AA may play a role in lymphocyte activation and proliferation [7,8,10]. This physiologic observation may be harnessed and augmented with pharmacologic AA concentrations. When given intravenously, pharmacologic AA concentrations promote the steady-state formation of hydrogen peroxide in vivo over hours [14,15]. Hydrogen peroxide, in turn,

may serve as an intracellular second messenger in the antiviral activity of lymphocytes [12]. Independent of lymphocyte activation, pharmacologic AA also has direct inhibitory effects on the replication of some viruses *in vitro*, presumably via hydrogen peroxide formation [9]. Together, these properties of pharmacologic AA could be harnessed in the early treatment of COVID-19 pneumonia.

Several trials of AA in COVID-19 patients are in progress or under analysis [18–20], in addition to a clinical trial of oral AA and zinc in patients with COVID-19 that was stopped early due to futility [21]. The trial presented here is unique in that it not only examined intravenous as opposed to oral AA but is also based on the hypothesis-driven mechanism where pharmacologic ascorbic acid produces hydrogen peroxide in the extracellular fluid. The data supporting this mechanism of action and clinical dosing application are primarily from the use of pharmacologic ascorbic acid in cancer treatment. Via hydrogen peroxide formation, pharmacologic ascorbic acid concentrations selectively target cancer cell death without harm to normal tissue [14–16,22–25]. The use of the intravenous route was essential to our study design as oral AA does not produce pharmacologic plasma concentrations of the vitamin nor does it increase baseline hydrogen peroxide concentrations in extracellular fluid [13,23]. The dose selected was based on its ability to safely generate maximal hydrogen peroxide concentrations *in vivo* in extracellular fluid, utilizing pre-clinical data and data obtained from patients with malignancies who received intravenous AA treatment.

Intravenous AA has been studied as a sepsis or acute respiratory distress syndrome (ARDS) therapeutic agent in clinical trials with placebo control and randomization [26–34]. AA was used either as a single agent or in combination with hydrocortisone and/or thiamine. The majority of evidence from these and other studies does not show benefits from supraphysiologic AA concentrations achieved from the doses used [35,36]. In sepsis studies, the doses used generally were approximately 25 mg/kg every 6 h, much less than the dose used here of 0.9 g/kg (900 mg/kg) once daily. Similarly, in sepsis studies, estimated peak plasma concentrations are predicted to be less than 5% of the peak concentrations predicted from the dose in our COVID trial [15,23,37]. Doses used in sepsis studies were initially selected to rapidly reverse AA deficiency unrelated to the generation of hydrogen peroxide, although the use of even these comparatively lower AA doses in a sepsis population might confer increased risk because of concurrent acute kidney injury in sepsis [38,39].

Based on our hypotheses, we predicted that the benefit of AA would be most apparent in the early study period. For this reason, we constructed a primary endpoint of clinical improvement within 72 h of randomization. The endpoint included three components—a reduction in supplemental oxygen, a reduction in bronchodilator use, and discharge. At this stage in our understanding of COVID-19, a reduction in supplemental oxygen use and discharge remain viable measures of a patient’s clinical course. The Center for Disease Control and Prevention continues to monitor COVID-19-related ventilatory and discharge patterns as a component of its National Hospital Care Survey (NHCS) [40]. Furthermore, in its guideline “COVID-19: Developing Drugs and Biological Products for Treatment and Prevention”, the United States Food and Drug Administration (FDA) recommends the need for hospitalization and the reduction in supplemental oxygen as important clinical outcomes to measure in pharmaceutical trials [41]. However, the reduction of bronchodilator use is not featured in either NHCS data or in the FDA trial recommendations. This likely reflects our current understanding of the pathophysiology of COVID-19. Bronchodilator use does not appear to improve clinical outcomes in COVID-19, possibly due to its predilection for parenchymal and vascular pathology as opposed to airway disease [42].

There was no statistically significant difference between AA and SoC for the primary endpoint of early clinical improvement. Despite the lack of composite statistical significance, point estimates of the endpoint’s individual components were all supportive of the treatment arm. As shown in analyses of the secondary endpoints (Table 3, Figure 2), there were early splits in the cumulative proportion curves for the time to a 50% reduction in oxygenation and the time to discharge. These early potential benefits disappeared over time. Although these are post-hoc analyses, they hint at a possible advantage to the

outcomes during the 72-h window after randomization, consistent with our hypotheses. Further consistent with our hypotheses, these trends were nullified as the curves crossed later in the hospitalization. As secondary endpoints, both the times to discharge and a 50% reduction in oxygenation were statistically non-significant when measured over the entire hospitalization. This pattern was not wholly unexpected. As noted above, an early benefit was consistent with our proposed mechanism of action.

Regarding safety, the proportion of patients who experienced a clinical decline in the treatment arm did not meet stopping rules and the trial was continued to its full accrual. Though the point estimates were higher for serious adverse events and clinical decline in the first 36 h, the study was not powered to detect these differences, which were not statistically significant.

Our trial had limitations. In retrospect, the study was small with optimistic outcome assumptions. Outcomes in this trial were defined in the early part of the pandemic, and do not reflect our current understanding of COVID-19 as a disease process. Furthermore, our patient risk stratification, while focused on age and comorbidities, may be improved in future studies. An emphasis on more severe underlying metabolic disease (i.e., diabetes), as opposed to any pulmonary or cardiovascular disease, may better improve patient selection and risk analyses. Viral load and specific analyses of lymphocyte populations were not feasible for practical and funding limitations, and these data may guide potential future trials of pharmacologic AA. As has become important both in clinical trials and in clinical practice, more consideration for symptom onset and baseline inflammatory markers, such as D-dimer and C-reactive protein, may lead to better correlates for clinical practice. Such considerations can strengthen future study designs for AA in the treatment of COVID-19.

Although the primary endpoint was not met in this study, trends in supplemental oxygen use and patient discharge suggest possible clinical signals consistent with the action of AA at pharmacologic doses. Even at these high doses, there was no statistical difference in safety between the two study arms. There is a continuing need for early treatment of COVID-19 in hospitalized patients, and this need will only increase as viral variants evolve. We believe that future larger studies exploring the use of pharmacologic AA are warranted and should account for its early disease effects and our evolving understanding of COVID-19 infection.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/life12030453/s1>, Table S1: CONSORT 2010 checklist of information to include when reporting a randomised trial.

Author Contributions: P.-C.V., D.M. (Daniel Monti) and M.L. conceived the study; D.C., D.M. (Daniel Monti), M.L. and M.F. developed the trial methodology; D.C., G.V., K.B. and B.M. reviewed and enrolled subjects; D.C., D.M. (Daniel Monti), G.V. and K.B. provided clinical oversight and monitored safety; B.M., D.M. (Devon Magee), I.M., L.S. and V.P. were responsible for preparing and curating data; B.C. and D.C. performed formal analyses; D.C. and M.L. prepared the original manuscript draft; all authors reviewed and edited the manuscript; D.M. (Daniel Monti) acquired funding; D.C., D.M. (Daniel Monti) and M.L. supervised overall efforts. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the Marcus Foundation and the Intramural program National Institute of Diabetes and Digestive and Kidney Disease; DK053212-16.

Institutional Review Board Statement: This study was approved by the Thomas Jefferson University Office of Human Research Institutional Review Board (Protocol #20D.431). It was conducted in accordance with Federal-Wide Assurance #00002109 to the U.S. Department of Health and Human Services.

Informed Consent Statement: Informed consent was obtained from all individual participants included in the study.

Data Availability Statement: The data presented in this study are available on request from the corresponding author. The data are not publicly available due to our institution's patient privacy policies.

Acknowledgments: The authors thank the Marcus Foundation, Atlanta, GA, for funding and support of this project. Mark Levine and Pierre-Christian Violet were supported by the Intramural Research Program, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD (grant Z01 DK053212-16). The authors thank the Jefferson Clinical Research Institute for its support throughout this project.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

AA	ascorbic acid, vitamin C
COVID-19	coronavirus disease 2019
SARS-CoV-2	severe acute respiratory syndrome coronavirus 2
SoC	standard of care

References

- Centers for Disease Control and Prevention. Estimated COVID-19 Burden. Available online: <https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/burden.html> (accessed on 9 September 2021).
- Solis Arce, J.S.; Warren, S.S.; Meriggi, N.F.; Scacco, A.; McMurry, N.; Voors, M.; Syunyaev, G.; Malik, A.A.; Aboutajdine, S.; Adejo, O.; et al. COVID-19 vaccine acceptance and hesitancy in low- and middle-income countries. *Nat. Med.* **2021**, *27*, 1385–1394. [[CrossRef](#)]
- Bergwerk, M.; Gonen, T.; Lustig, Y.; Amit, S.; Lipsitch, M.; Cohen, C.; Mandelboim, M.; Gal Levin, E.; Rubin, C.; Indenbaum, V.; et al. COVID-19 Breakthrough Infections in Vaccinated Health Care Workers. *N. Engl. J. Med.* **2021**, *385*, 1474–1484. [[CrossRef](#)] [[PubMed](#)]
- Hacisuleyman, E.; Hale, C.; Saito, Y.; Blachere, N.E.; Bergh, M.; Conlon, E.G.; Schaefer-Babajew, D.J.; DaSilva, J.; Muecksch, F.; Gaebler, C.; et al. Vaccine Breakthrough Infections with SARS-CoV-2 Variants. *N. Engl. J. Med.* **2021**, *384*, 2212–2218. [[CrossRef](#)] [[PubMed](#)]
- Baj, A.; Novazzi, F.; Pasciuta, R.; Genoni, A.; Ferrante, F.D.; Valli, M.; Partenope, M.; Tripiciano, R.; Ciserchia, A.; Catanoso, G.; et al. Breakthrough Infections of E484K-Harboring SARS-CoV-2 Delta Variant, Lombardy, Italy. *Emerg. Infect. Dis.* **2021**, *27*, 3180–3182. [[CrossRef](#)] [[PubMed](#)]
- Desai, N.; Neyaz, A.; Szabolcs, A.; Shih, A.R.; Chen, J.H.; Thapar, V.; Nieman, L.T.; Solovyov, A.; Mehta, A.; Lieb, D.J.; et al. Temporal and spatial heterogeneity of host response to SARS-CoV-2 pulmonary infection. *Nat. Commun.* **2020**, *11*, 1–15. [[CrossRef](#)] [[PubMed](#)]
- Siegel, B.V.; Morton, J.I. Vitamin C and the immune response. *Experientia* **1977**, *33*, 393–395. [[CrossRef](#)] [[PubMed](#)]
- Jeong, Y.-J.; Hong, S.-W.; Kim, J.-H.; Jin, D.-H.; Kang, J.S.; Lee, W.J.; Hwang, Y.-I. Vitamin C-treated murine bone marrow-derived dendritic cells preferentially drive naïve T cells into Th1 cells by increased IL-12 secretions. *Cell. Immunol.* **2011**, *266*, 192–199. [[CrossRef](#)] [[PubMed](#)]
- Harakeh, S.; Jariwalla, R.J.; Pauling, L. Suppression of human immunodeficiency virus replication by ascorbate in chronically and acutely infected cells. *Proc. Natl. Acad. Sci. USA* **1990**, *87*, 7245–7249. [[CrossRef](#)] [[PubMed](#)]
- Van Gorkom, G.N.Y.; Wolterink, R.G.J.K.; Van Elssen, C.H.M.J.; Wieten, L.; Germeraad, W.T.V.; Bos, G.M.J. Influence of Vitamin C on Lymphocytes: An Overview. *Antioxidants* **2018**, *7*, 41. [[CrossRef](#)]
- Biancatelli, R.M.L.C.; Berrill, M.; Marik, P.E. The antiviral properties of vitamin C. *Expert Rev. Anti-Infect. Ther.* **2019**, *18*, 99–101. [[CrossRef](#)] [[PubMed](#)]
- Reth, M. Hydrogen peroxide as second messenger in lymphocyte activation. *Nat. Immunol.* **2002**, *3*, 1129–1134. [[CrossRef](#)] [[PubMed](#)]
- Padayatty, S.J.; Sun, H.; Wang, Y.; Riordan, H.D.; Hewitt, S.M.; Katz, A.; Wesley, R.A.; Levine, M. Vitamin C Pharmacokinetics: Implications for Oral and Intravenous Use. *Ann. Intern. Med.* **2004**, *140*, 533–537. [[CrossRef](#)] [[PubMed](#)]
- Chen, Q.; Espey, M.G.; Sun, A.Y.; Lee, J.-H.; Krishna, M.C.; Shacter, E.; Choyke, P.L.; Pooput, C.; Kirk, K.L.; Buettner, G.R.; et al. Ascorbate in pharmacologic concentrations selectively generates ascorbate radical and hydrogen peroxide in extracellular fluid in vivo. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 8749–8754. [[CrossRef](#)]
- Chen, Q.; Espey, M.G.; Sun, A.Y.; Pooput, C.; Kirk, K.L.; Krishna, M.C.; Khosh, D.B.; Drisko, J.; Levine, M. Pharmacologic doses of ascorbate act as a prooxidant and decrease growth of aggressive tumor xenografts in mice. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 11105–11109. [[CrossRef](#)] [[PubMed](#)]
- Levine, M.; Violet, P.-C. Data Triumph at C. *Cancer Cell* **2017**, *31*, 467–469. [[CrossRef](#)] [[PubMed](#)]
- Centers for Disease Control and Prevention. Adverse Events. Available online: https://clinicaltrials.gov/ct2/help/adverse_events_desc (accessed on 8 March 2022).
- Sisli Hamidiye Etfal Training and Research Hospital. High-Dose Vitamin C Treatment in Critically Ill COVID-19 Patients. Available online: <https://ClinicalTrials.gov/show/NCT04710329> (accessed on 8 March 2022).

19. Lahore, U.O. Role of Mega Dose of Vitamin C in Critical COVID-19 Patients. Available online: <https://ClinicalTrials.gov/show/NCT04682574> (accessed on 8 March 2022).
20. Hunter Holmes Mcguire Veteran Affairs Medical Center; Mcguire Research Institute, Administration of Intravenous Vitamin C in Novel Coronavirus Infection (COVID-19) and Decreased Oxygenation. Available online: <https://ClinicalTrials.gov/show/NCT04357782> (accessed on 8 March 2022).
21. Thomas, S.; Patel, D.; Bittel, B.; Wolski, K.; Wang, Q.; Kumar, A.; Il'Giovine, Z.J.; Mehra, R.; McWilliams, C.; Nissen, S.E.; et al. Effect of High-Dose Zinc and Ascorbic Acid Supplementation vs Usual Care on Symptom Length and Reduction among Ambulatory Patients with SARS-CoV-2 Infection: The COVID A to Z Randomized Clinical Trial. *JAMA Netw. Open* **2021**, *4*, e210369. [[CrossRef](#)] [[PubMed](#)]
22. Chen, Q.; Espey, M.G.; Krishna, M.C.; Mitchell, J.B.; Corpe, C.P.; Buettner, G.R.; Shacter, E.; Levine, M. Pharmacologic ascorbic acid concentrations selectively kill cancer cells: Action as a pro-drug to deliver hydrogen peroxide to tissues. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 13604–13609. [[CrossRef](#)] [[PubMed](#)]
23. Hoffer, L.J.; Levine, M.; Assouline, S.; Melnychuk, D.; Padayatty, S.J.; Rosadiuk, K.; Rousseau, C.; Robitaille, L.; Miller, W.H. Phase I clinical trial of i.v. ascorbic acid in advanced malignancy. *Ann. Oncol.* **2008**, *19*, 1969–1974. [[CrossRef](#)]
24. Nauman, G.; Gray, J.C.; Parkinson, R.; Levine, M.; Paller, C.J. Systematic Review of Intravenous Ascorbate in Cancer Clinical Trials. *Antioxidants* **2018**, *7*, 89. [[CrossRef](#)] [[PubMed](#)]
25. Schoenfeld, J.D.; Sibenaller, Z.A.; Mapuskar, K.A.; Wagner, B.A.; Cramer-Morales, K.L.; Furqan, M.; Sandhu, S.; Carlisle, T.L.; Smith, M.C.; Abu Hejleh, T.; et al. O₂– and H₂O₂-Mediated Disruption of Fe Metabolism Causes the Differential Susceptibility of NSCLC and GBM Cancer Cells to Pharmacological Ascorbate. *Cancer Cell* **2017**, *32*, 268. [[CrossRef](#)] [[PubMed](#)]
26. Fujii, T.; Luethi, N.; Young, P.; Frei, D.R.; Eastwood, G.M.; French, C.J.; Deane, A.; Shehabi, Y.; Hajjar, L.A.; Oliveira, G.; et al. Effect of Vitamin C, Hydrocortisone, and Thiamine vs Hydrocortisone Alone on Time Alive and Free of Vasopressor Support Among Patients With Septic Shock. *JAMA* **2020**, *323*, 423–431. [[CrossRef](#)] [[PubMed](#)]
27. Iglesias, J.; Vassallo, A.V.; Patel, V.; Sullivan, J.B.; Cavanaugh, J.; Elbaga, Y. Outcomes of Metabolic Resuscitation Using Ascorbic Acid, Thiamine, and Glucocorticoids in the Early Treatment of Sepsis: The ORANGES Trial. *Chest* **2020**, *158*, 164–173. [[CrossRef](#)] [[PubMed](#)]
28. Wani, S.J.; A Mufti, S.; Jan, R.A.; Shah, S.U.; Qadri, S.M.; Khan, U.H.; Bagdadi, F.; Mehfooz, N.; Koul, P.A. Combination of vitamin C, thiamine and hydrocortisone added to standard treatment in the management of sepsis: Results from an open label randomised controlled clinical trial and a review of the literature. *Infect. Dis.* **2020**, *52*, 271–278. [[CrossRef](#)] [[PubMed](#)]
29. Chang, P.; Liao, Y.; Guan, J.; Guo, Y.; Zhao, M.; Hu, J.; Zhou, J.; Wang, H.; Cen, Z.; Tang, Y.; et al. Combined Treatment With Hydrocortisone, Vitamin C, and Thiamine for Sepsis and Septic Shock: A Randomized Controlled Trial. *Chest* **2020**, *158*, 174–182. [[CrossRef](#)]
30. Moskowitz, A.; Huang, D.T.; Hou, P.C.; Gong, J.; Doshi, P.B.; Grossestreuer, A.V.; Andersen, L.W.; Ngo, L.; Sherwin, R.L.; Berg, K.M.; et al. Effect of Ascorbic Acid, Corticosteroids, and Thiamine on Organ Injury in Septic Shock: The ACTS Randomized Clinical Trial. *JAMA* **2020**, *324*, 642–650. [[CrossRef](#)] [[PubMed](#)]
31. Hwang, S.Y.; Ryoo, S.M.; Park, J.E.; Jo, Y.H.; Jang, D.-H.; Suh, G.J.; Kim, T.; Kim, Y.-J.; Kim, S.; Cho, H.; et al. Combination therapy of vitamin C and thiamine for septic shock: A multi-centre, double-blinded randomized, controlled study. *Intensiv. Care Med.* **2020**, *46*, 2015–2025. [[CrossRef](#)] [[PubMed](#)]
32. Menon, V.; Mohamed, Z.U.; Prasanna, P.; Moni, M.; Edathadathil, F.; Prasanna, P.; Menon, A.; Nair, S.; Greeshma, C.; Sathyapalan, D.T.; et al. Vitamin C Therapy for Routine Care in Septic Shock (ViCTOR) Trial: Effect of Intravenous Vitamin C, Thiamine, and Hydrocortisone Administration on Inpatient Mortality among Patients with Septic Shock. *Indian J. Crit. Care Med.* **2020**, *24*, 653–661. [[CrossRef](#)] [[PubMed](#)]
33. Sevransky, J.E.; Rothman, R.E.; Hager, D.N.; Bernard, G.R.; Brown, S.M.; Buchman, T.G.; Busse, L.W.; Coopersmith, C.M.; DeWilde, C.; Ely, E.W.; et al. Effect of Vitamin C, Thiamine, and Hydrocortisone on Ventilator- and Vasopressor-Free Days in Patients With Sepsis: The VICTAS Randomized Clinical Trial. *JAMA* **2021**, *325*, 742–750. [[CrossRef](#)]
34. Fowler, A.A., 3rd; Truitt, J.D.; Hite, R.D.; Morris, P.E.; Dewilde, C.; Priday, A.; Fisher, B.; Thacker, L.R., 2nd; Natarajan, R.; Brophy, D.F.; et al. Effect of Vitamin C Infusion on Organ Failure and Biomarkers of Inflammation and Vascular Injury in Patients with Sepsis and Severe Acute Respiratory Failure. *JAMA* **2019**, *322*, 1261–1270. [[CrossRef](#)] [[PubMed](#)]
35. Fujii, T.; Salanti, G.; Belletti, A.; Bellomo, R.; Carr, A.; Furukawa, T.A.; Luethi, N.; Luo, Y.; Putzu, A.; Sartini, C.; et al. Effect of adjunctive vitamin C, glucocorticoids, and vitamin B1 on longer-term mortality in adults with sepsis or septic shock: A systematic review and a component network meta-analysis. *Intensiv. Care Med.* **2021**, *48*, 16–24. [[CrossRef](#)]
36. Agarwal, A.; Hager, D.N.; Sevransky, J.E. Any Role of High-Dose Vitamin C for Septic Shock in 2021? *Semin. Respir. Crit. Care Med.* **2021**, *42*, 672–682. [[CrossRef](#)]
37. Rosengrave, P.; Spencer, E.; Williman, J.; Mehrtens, J.; Morgan, S.; Doyle, T.; Van Der Heyden, K.; Morris, A.; Shaw, G.; Carr, A.C. Intravenous vitamin C administration to patients with septic shock: A pilot randomised controlled trial. *Crit. Care* **2022**, *26*, 1–10. [[CrossRef](#)] [[PubMed](#)]
38. Marik, P.E.; Khangoora, V.; Rivera, R.; Hooper, M.H.; Catravas, J. Hydrocortisone, Vitamin C, and Thiamine for the Treatment of Severe Sepsis and Septic Shock. *Chest* **2017**, *151*, 1229–1238. [[CrossRef](#)] [[PubMed](#)]
39. McCune, T.R.; Toepp, A.J.; Sheehan, B.E.; Sherani, M.S.K.; Petr, S.T.; Dodani, S. High dose intravenous vitamin C treatment in sepsis: Associations with acute kidney injury and mortality. *BMC Nephrol.* **2021**, *22*, 387. [[CrossRef](#)] [[PubMed](#)]

40. Centers for Disease Control and Prevention. National Hospital Care Survey. Available online: <https://www.cdc.gov/nchs/nhcs/index.htm> (accessed on 8 March 2022).
41. United States Food and Drug Administration. COVID-19: Developing Drugs and Biological Products for Treatment or Prevention Guidance for Industry. Available online: <https://www.fda.gov/media/137926/download> (accessed on 8 March 2022).
42. Villamañán, E.; Sobrino, C.; Carpio, C.; Moreno, M.; Arancón, A.; Lara, C.; Pérez, E.; Jiménez, C.; Zamarrón, E.; Jiménez-Nácher, I.; et al. Inhaled bronchodilators use and clinical course of adult inpatients with Covid-19 pneumonia in Spain: A retrospective cohort study. *Pulm. Pharmacol. Ther.* **2021**, *69*, 102007. [[CrossRef](#)] [[PubMed](#)]