

1-1-2018

Sex Differences in Outcomes after Stroke in Patients with Diabetes in Ontario, Canada.

Mandip S. Dhamoon
Icahn School of Medicine at Mount Sinai

John W. Liang
Thomas Jefferson University

Limei Zhou
Institute for Clinical Evaluative Sciences

Melissa Stamplecoski
Institute for Clinical Evaluative Sciences

Moira K. Kapral
Institute for Clinical Evaluative Sciences; University of Toronto

Follow this and additional works at: <https://jdc.jefferson.edu/neurologyfp>



Part of the [Department of Neurology Faculty Papers](#), [Diabetes, and Metabolism Commons](#), and the [Neurology Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Dhamoon, Mandip S.; Liang, John W.; Zhou, Limei; Stamplecoski, Melissa; Kapral, Moira K.; and Shah, Baiju R., "Sex Differences in Outcomes after Stroke in Patients with Diabetes in Ontario, Canada." (2018). *Department of Neurology Faculty Papers*. Paper 148.
<https://jdc.jefferson.edu/neurologyfp/148>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Neurology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Mandip S. Dhamoon, John W. Liang, Limei Zhou, Melissa Stamplecoski, Moira K. Kapral, and Baiju R. Shah

Sex differences in outcomes after stroke in patients with diabetes in Ontario, Canada

Mandip S. Dhamoon, MD,DrPH*,¹ John W. Liang*,MD,^{1,2} Limei Zhou,PhD,⁴ Melissa
Stamplecoski,BSc,⁴ Moira K. Kapral,MD,MSc,^{3,4} Baiju R. Shah,MD,PhD^{3,4}

¹Department of Neurology, Icahn School of Medicine at Mount Sinai, New York, NY.

²Department of Neurology, Thomas Jefferson University, Philadelphia, PA

³Department of Medicine, University of Toronto, Toronto,ON

⁴Institute for Clinical Evaluative Sciences, Toronto,ON

*co-first authors

Correspondence address:

John W. Liang MD

Thomas Jefferson University Hospital

Department of Neurology

909 Walnut Street, 4th floor

Philadelphia, PA 19107

Email: john.liang@jefferson.edu

Telephone: (718) 734-7760

Abstract word count: 250

Text word count: 3734

Abstract

Background: Outcomes after stroke in those with diabetes are not well characterized, especially by sex and age. We sought to calculate the sex- and age-specific risk of cardiovascular outcomes after ischemic stroke among those with diabetes.

Methods: Using population-based demographic and administrative health care databases in Ontario, Canada, all patients with diabetes hospitalized with index ischemic stroke between April 1, 2002 and March 31, 2012 were followed for death, stroke, and myocardial infarction (MI). Kaplan-Meier survival analysis and Fine-Gray competing risk models estimated hazards of outcomes by sex and age, unadjusted and adjusted for demographics and vascular risk factors.

Results: Among 25495 diabetic patients with index ischemic stroke, incidence of death was higher in women than in men (14.08 per 100 person-years [95% CI 13.73-14.44] vs. 11.89 [11.60-12.19]), but was lower after adjustment for age and other risk factors (adjusted hazard ratio [HR] 0.95 [0.92-0.99]). Recurrent stroke incidence was similar by sex, but men were more likely to be readmitted for MI (1.99 per 100 person-years [1.89-2.10] vs 1.58 [1.49-1.68] among females). In multivariable models, females had a lower risk of readmission for any event (HR 0.96 [95% CI 0.93-0.99]).

Conclusions: In this large, population-based, retrospective study among diabetic patients with index stroke, women had higher unadjusted death rate but lower unadjusted incidence of MI. In adjusted models, females had a lower death rate compared to males, although the increased risk of MI among males persisted. These findings confirm and quantify sex differences in outcomes after stroke in patients with diabetes.

1 INTRODUCTION

2 There are sex differences in the risk of cardiovascular disease in people with diabetes. Compared
3 to men, women with diabetes have a 40% higher risk of incident coronary heart disease¹ and
4 27% higher risk of stroke.² However, sex differences in outcomes in diabetic patients following
5 an incident event are unclear, with conflicting findings in previous studies.^{3,4} Sex differences
6 have been demonstrated for myocardial infarction (MI)^{5,6} and other cardiovascular disease,⁷ but
7 data on sex differences in outcomes among people with diabetes after incident stroke are less
8 consistent. Relatively small studies have reported higher in-hospital mortality⁸ and long-term
9 mortality⁹ for diabetic females, but others have shown no association of sex and outcomes^{10,11}
10 Furthermore, prior studies mostly examined mortality and did not measure readmission rates.
11 Studies to date have not adequately assessed for socioeconomic status and medication usage,
12 which may confound the relationship between sex and outcomes. There is a lack of reliable
13 population-based data on the effect of sex on mortality and readmissions among diabetic patients
14 following an incident stroke.

15 The objective of this analysis was to examine differences in cardiovascular events and
16 mortality by sex and age among those with diabetes after ischemic stroke in Ontario. We
17 hypothesized that women had higher mortality compared to men and that the readmission risk for
18 cardiovascular events differed by sex.

19

20 METHODS

21 We conducted a retrospective analysis of a population-based sample using linked administrative
22 databases in Ontario, Canada's most populous province. Because of government-funded health
23 insurance for all permanent residents of Ontario, data were available on the entire population.

1 The Ontario Registered Persons Database (RPDB) provided data on mortality after stroke, and
2 the Canadian Institute for Health Information Discharge Abstract Database (CIHI-DAD)
3 identified readmissions for stroke and myocardial infarction (MI). CIHI-DAD contains ≤ 25
4 diagnosis fields for admissions to Ontario hospitals and uses the International Classification of
5 Diseases, 10th revision coding system (ICD-10) for the year 2002 onwards. In previous studies
6 of CIHI-DAD in Canadian hospitals, there was high positive predictive value (85% for ischemic
7 stroke, 98% for intracerebral hemorrhage, and 91% for subarachnoid hemorrhage) and Kappa of
8 0.89 for agreement between coder and researcher using ICD-10 codes.¹² The Ontario Drug
9 Benefits (ODB) database provided information on prescriptions filled by all residents aged ≥ 65
10 years. These databases were linked via a unique, encoded identifier and analyzed at the Institute
11 for Clinical Evaluative Sciences (ICES). The institutional ethics review board of Sunnybrook
12 Health Sciences Centre approved this study.

13

14 **Sample selection**

15 We included those with an index ischemic stroke admission during the study period in CIHI-
16 DAD, identified with any of the following ICD-10 codes: I63 (excluding I63.6), I64, H34.0, or
17 H34.1 in the “most responsible diagnosis” field, which has been shown to have 92% accuracy for
18 stroke diagnosis.¹³ We identified diagnosis of diabetes prior to or at the time of the index
19 ischemic stroke admission by linking to the Ontario Diabetes Database (ODD), which has a
20 sensitivity of 91% and specificity of 99%.¹⁴ We limited the sample to those with ischemic stroke
21 and diabetes who were ≥ 18 years of age at the time of admission. Index stroke admissions from
22 4/1/2002 to 3/31/2012 were included, with maximum follow-up to 3/31/2013.

23

1 **Baseline assessment**

2 Age was calculated as age at admission for index ischemic stroke. Income was estimated using
3 neighborhood-level household income and was categorized into quintiles. Duration of diabetes
4 was calculated by using the diagnosis date in the ODD and was categorized into: 0 to <3, 3 to <6,
5 and ≥ 6 years. Duration of Ontario residence was inferred from duration of having a health card
6 and was categorized into: 0 to <5, 5 to <10 and ≥ 10 years. Using standard ICD-9 (prior to 2002)
7 and ICD-10 (2002 onwards) code clusters, we identified history of stroke or transient ischemic
8 attack (TIA), atrial fibrillation, hypertension, MI, coronary artery disease, and peripheral
9 vascular disease (PVD).

10 The Charlson comorbidity index (CCI) was calculated using all diagnosis codes and types
11 from all hospitalizations during the two year period prior to and including the index admission,
12 using ≤ 25 available ICD-9 and ICD-10 codes for each hospitalization. Since all participants in
13 this analysis had diabetes, the diabetes indicators were excluded from our CCI calculation. The
14 CCI was dichotomized into < 2 versus ≥ 2 , as in previous research.¹⁵

15 Only patients ≥ 65 years of age had complete information on prescription medication use.
16 Baseline medication was defined as any prescription medication use within a 120-day window
17 after the index stroke discharge, and medication classes included diabetic, statin, and anti-
18 hypertensive medications as well as warfarin. Aspirin, which is available over-the-counter, had
19 incomplete capture; hence, antiplatelet medication use was not adjusted for in the sensitivity
20 analyses.

21

22 **Longitudinal follow-up**

1 Outcomes were: death, any-cause readmission, stroke/TIA readmission, MI readmission, stroke
2 or MI readmission, and a composite of death or any-cause readmission. To create these
3 outcomes, hospital readmissions for the following were assessed: recurrent stroke/TIA (ICD-10
4 code I63 [excluding I63.6], I64, H34.0, H34.1, G45 [excluding G45.4], H34.0), intracerebral
5 hemorrhage (I61), recurrent CAD including MI (acute MI, codes I21, I22; unstable angina, code
6 I20) and cardiac procedures (coronary artery bypass graft and percutaneous cardiac intervention).

7 8 **Statistical analysis**

9 Baseline characteristics were reported in the overall sample and stratified by sex. We calculated
10 proportions for categorical variables and means and medians for continuous variables. Incidence
11 was calculated as the incidence of each outcome per 100 person-years (with 95% confidence
12 intervals), reported by age and sex subgroups. Two significance tests were performed using a
13 Poisson regression model: one for the group comparison within each stratum, and the other for
14 the overall test of significance of that stratum.

15 For all readmission outcomes (excluding death), a competing risk model proposed by
16 Fine and Gray¹⁶ was used to estimate the hazard ratio and 95% CI of outcomes, with death
17 defined as the competing risk. For the outcome of death, Cox proportional hazards models were
18 used to estimate the hazard ratio and 95% CI. Models included demographic variables (age, sex,
19 and income) and vascular risk factors (hypertension, atrial fibrillation, stroke or TIA, MI, CAD,
20 PVD, and CCI). A sensitivity analysis was performed among those aged ≥ 65 years, among
21 whom medication use was adjusted for and categorized into anti-hypertensive medication use,
22 diabetes medication use, statin use, and warfarin use. All analyses were performed with SAS
23 version 9.3 (SAS Institute Inc, Cary, NC).

1 RESULTS

2 Out of 84,731 index ischemic stroke admissions during the study time period, 29,752 had
3 diabetes prior to index admission and were included. After applying exclusions (age <18 years;
4 death during index admission; death after index discharge but before any readmissions), the final
5 sample consisted of 25,495 individuals. Median follow-up time was 3.2 years.

6 Table 1 shows baseline characteristics in the entire sample (n=25,495) and by sex (11,902
7 females and 13,593 males). Compared to males, females were older and more frequently from
8 lower neighborhood income quintiles. Females more often had history of prior stroke or TIA,
9 atrial fibrillation, and hypertension but less often had history of MI, CAD or PVD; females had
10 lower CCI scores but longer average length of hospital stay. In those aged ≥ 65 years, females at
11 baseline were significantly less often taking diabetic or statin medications.

12 There were 12,435 deaths during follow-up. The overall frequencies of outcomes by
13 time interval are listed in Table 1. The unadjusted incidence of death was higher among females
14 (14.08 per 100 person-years, 95% CI 13.73-14.44 vs. 11.89, 95% CI 11.60-12.19 among males)
15 and there were higher rates of death among higher age groups (Table 2). Kaplan-Meier curves
16 showed lower survival probability for females compared to males ($p < 0.0001$) (Figure 1A). After
17 adjusting for age, income, vascular risk factors, and CCI, women had lower risk of death
18 compared to men (HR 0.95, 95% CI 0.92-0.99, Table 3). Among those ≥ 65 years old with or
19 without adjustment for medications, a similar finding of lower risk of death among females was
20 seen (HR 0.93, 95% CI 0.89-0.97 and HR 0.93, 95% CI 0.90-0.97, respectively).

21 There were 17,406 any-cause readmissions during follow-up, of which 3,794 were for
22 stroke and 2,512 for MI. The unadjusted incidence rate of readmission for any cause or for stroke
23 was similar by sex (Table 2). There was higher unadjusted incidence among males for

1 readmission due to MI (1.99, 95% CI 1.89-2.10 vs. 1.58, 95% CI 1.49-1.68 among females).
2 There was a pattern of increased incidence of death and lower incidence of readmissions for
3 stroke, MI, and stroke or MI among successively higher age categories, probably reflecting the
4 competing risk of death for stroke and MI (Table 2). Kaplan-Meier curves showed a similar
5 survival probability for males and females of any readmission (Figure 1B) and stroke
6 readmission (Figure 2A), but a higher risk of MI readmission among males (Figure 2B).

7 In multivariable competing risk models, females had a lower risk of readmission for any
8 event (0.96, 95% CI 0.93-0.99, Table 3). There was no sex difference in risk of readmission for
9 stroke, but females had a lower risk of MI readmission compared to males (0.88, 95% CI 0.81-
10 0.95). Among those age ≥ 65 years with and without adjustment for medication usage, risk of
11 readmission for any event remained lower in females but risk of MI readmission was no longer
12 different by sex.

13

14 **DISCUSSION**

15 In this population-based, retrospective study using administrative linkage with full population
16 coverage for the province of Ontario, we found that women with diabetes, compared to men, had
17 higher unadjusted mortality and risk of readmission for any cause or death following an incident
18 stroke but lower risk of readmission for MI. Unadjusted readmission rates for any cause and for
19 stroke were similar by sex. Diabetic female patients with incident stroke were older and from a
20 lower socioeconomic status. They were more likely to have hypertension and atrial fibrillation
21 but less likely to have prior MI, CAD, or PVD. They were also less likely to be taking diabetic
22 and statin medications at baseline and had a longer average length of hospital stay during the
23 incident stroke, suggesting either differences in disease severity or disparities in optimal

1 treatment by sex. These differences likely accounted for the unadjusted mortality difference seen
2 by sex, because in adjusted models females had lower risk of death and lower risk of any-cause
3 readmissions compared to males, although the increased risk of MI among males persisted after
4 adjustment. Readmission rates for stroke remained similar between males and females in
5 adjusted models. Also, as expected, there was a higher risk of mortality and lower risk of
6 readmission with increasing age, likely due to the competing risk of death.

7 The impact of traditional cardiovascular risk factors varies by sex, especially for smoking
8 (which carries a 25% greater risk for coronary heart disease among women than men^{17, 18}) and
9 diabetes.^{1, 2, 17, 18} In addition to traditional risk factors, there are risk factors specific to women,
10 including gestational hypertension and pre-eclampsia, gestational diabetes, and placental
11 disorders such as intrauterine growth restriction and stillbirth.^{17, 19-21} Polycystic ovarian
12 syndrome, the most common female endocrine disorder, results in insulin resistance and
13 development of metabolic syndrome.²² Oral contraceptive pills, used by 82% of sexually ever-
14 active women, are associated with elevated risk of venous thrombosis, MI and ischemic stroke
15 from a presumed pro-coagulant effect.¹⁸ Systemic autoimmune collagen vascular diseases, such
16 as systemic lupus erythematosus and rheumatoid arthritis, are more common in females and lead
17 to accelerated atherosclerosis and progression to heart disease.^{20, 21} Depression is twice as
18 common in women and associated with a 70% risk for heart disease; it can lead to non-adherence
19 with diet, medications, and follow-up.²⁰

20 Social support and self-reported quality of life have also been reported to be lower in
21 diabetic women.^{23, 24} Prior studies show that women have lower socioeconomic status and lower
22 access to preventative measures and treatments for diabetes.^{7, 25} There are also treatment
23 disparities – women are less likely to be prescribed medications for modifiable risk factors and,

1 even when undergoing treatment, they are treated less effectively.^{20, 22, 26, 27} Our findings also
2 reflect this female socioeconomic disadvantage as well as lower medication usage, despite
3 universal health coverage.

4 Prior studies of sex differences in outcomes among diabetic stroke patients have been in
5 smaller samples, focused on mortality, and with limited control of socioeconomic status and
6 medication usage.⁸⁻¹⁰ In a Spanish prospective single center stroke registry of 561 diabetic stroke
7 patients, there was higher mortality among women but similar stroke recurrence rates by sex.⁸
8 Elevated female mortality and similar stroke recurrence rates by sex were also reported in a
9 single-center Chinese study of 2360 diabetic stroke patients.¹⁰ In a Swedish population-based
10 study involving 2549 diabetic stroke patients under age 75 years, there was also higher mortality
11 noted in females; however readmissions were not assessed.⁹

12 Our study included over 25,000 diabetic stroke patients and provides reliable evidence
13 that women have higher mortality after stroke. We demonstrated that this difference is not
14 present after adequate adjustment, and women actually have a lower age-adjusted risk for
15 mortality. This finding is in agreement with a recent meta-analysis using 16,957 pooled
16 individual participant data from 13 population-based stroke incidence studies from Europe,
17 Australia, South America, and the Caribbean; it reported a lower crude survival rate in women at
18 1 and 5 years, which was reversed after adjustment.²⁸ The 5-year pooled estimates had
19 significant heterogeneity because few studies had follow-up beyond 1 year and there was
20 missing data across studies, particularly on stroke risk factors such as diabetes (only 5 out of 13
21 studies reported diabetes status, n=667). Our current report consist of more patients than all 13
22 studies combined and provides confirmation that female sex, by itself, is not responsible for
23 increased mortality after stroke in those with diabetes. Beyond mortality outcomes, we also

1 demonstrate that there is no sex difference in stroke readmission risk, but males are at higher risk
2 of readmission for MI. To our knowledge, there have been no prior studies among diabetic
3 patients with incident stroke reporting the effect of sex and MI vs stroke readmissions.

4 We found that men had higher risk of MI readmission, suggesting a possible sex-specific
5 sensitivity to different diabetes-related complications that has been previously demonstrated.²⁷
6 Men, compared to women, have been reported to have higher coronary atheroma burden, more
7 diffuse endothelial dysfunction, more severe structural abnormalities in the epicardial coronary
8 arteries, and more vulnerable plaques.²⁹ Young women are at lower risk of cardiovascular death,
9 MI, and stroke compared to males, presumably from the cardiovascular protective effects of
10 estrogen.^{20, 29, 30} However, the risk profile reverses after menopause with a 10-fold rise in
11 cardiovascular disease in women compared to a 4.5-fold rise in men of similar age.^{20, 29, 30} The
12 protective effects of estrogen likely explains the disappearance of the elevated MI readmission
13 risk when the sample is restricted to those above age 65, with or without adjustment for
14 medication usage. Interestingly, this same protective effect was not seen for stroke readmission
15 in analyses of the unadjusted, adjusted, and age >65 subgroup models. This suggests that
16 diabetes in pre-menopausal women may blunt the protective effects of estrogen to varying
17 degrees which may be organ specific.^{17, 20, 30, 31}

18 This study attempts to overcome several limitations of prior studies in the area of sex
19 differences after ischemic stroke. Due to universal health coverage in Ontario, the sample is
20 unprecedented in that it includes all adults with diabetes and index ischemic stroke in a large
21 Canadian province, not just patients from a single center or a population sample. Hence, this
22 results in extensive population coverage over a long follow-up period with limited selection bias.
23 Lack of power to detect sex differences is not a concern. Also, due to the unique linking among

1 different databases, adjustment for the important confounders of socioeconomic status and
2 medication use was possible.

3 There are limitations associated with the usage of administrative and claims-based data,
4 which may be prone to misclassification and inaccuracy. For example, sample selection using
5 ICD code I64, “stroke, not specified as hemorrhagic or infarct”, could possibly capture stroke-
6 types other than ischemic stroke. However, the sensitivity and specificity of diagnosis based on
7 ICD-10-based codes has been shown to be excellent. We only have data on events that resulted
8 in readmission; therefore some events such as TIA may be missed if patients were not admitted
9 to the hospital. Data on stroke characteristics, such as subtype, severity, location, size, and
10 discharge handicap were not available. Similarly, data was available on duration of diabetes but
11 not on severity of diabetes as assessed by hemoglobin A1c levels. We were able to control for
12 diabetic, statin and anti-hypertensive medication but could not assess for aspirin usage as it is
13 available over the counter and therefore could not be reliably controlled. Aspirin’s role in
14 preventing cardiovascular events in diabetic women is uncertain.²² There is insufficient evidence
15 that aspirin has a sex-specific cardiovascular impact, and further study is indicated.
16 Cardiovascular prevention may need to be tailored according to sex, and some studies have
17 suggested differential effectiveness of interventions by sex.^{32, 33} A structured personalized
18 approach may be more effective for women compared to men, but more research is needed.²³

19 In summary, we demonstrated that diabetic female patients have higher mortality after
20 incident stroke, but female sex was not an independent risk factor. Contrary to previous studies,
21 female sex was associated with lower mortality after adjustment for vascular risk factors,
22 demographics, socioeconomic status, and medication usage.

23

Acknowledgements: none

Funding Sources: Mandip Dhamoon was supported by the National Institute of Neurological Disorders And Stroke of the National Institutes of Health under Award Number K23NS079422. Moira Kapral was supported by a Career Investigator Award from the Heart and Stroke Foundation, Ontario Provincial Office. This study was funded by an operating grant from the Canadian Stroke Network. The Institute for Clinical Evaluative Sciences is supported by an operating grant from the Ministry of Health and Long-Term Care. Parts of this material are based on data and information compiled and provided by the Canadian Institute of Health Information. We thank IMS Brogan Inc. for use of their Drug Information Database. The opinions, results and conclusions are those of the authors and should not be attributed to any supporting or sponsoring agencies. No endorsement by the above is intended or should be inferred.

Disclosures

None of the authors has a potential conflict of interest related to the manuscript.

References

1. Peters SA, Huxley RR, Woodward M. Diabetes as risk factor for incident coronary heart disease in women compared with men: a systematic review and meta-analysis of 64 cohorts including 858,507 individuals and 28,203 coronary events. *Diabetologia* 2014;57:1542-1551.
2. Peters SA, Huxley RR, Woodward M. Diabetes as a risk factor for stroke in women compared with men: a systematic review and meta-analysis of 64 cohorts, including 775,385 individuals and 12,539 strokes. *Lancet* 2014;383:1973-1980.
3. Shi GM, Zhang YD, Geng C, et al. Profile and 1-Year Outcome of Ischemic Stroke in East China: Nanjing First Hospital Stroke Registry. *J Stroke Cerebrovasc Dis* 2016;25:49-56.
4. Mathisen SM, Dalen I, Larsen JP, Kurz M. Long-Term Mortality and Its Risk Factors in Stroke Survivors. *J Stroke Cerebrovasc Dis* 2016;25:635-641.
5. Meisinger C, Heier M, von Scheidt W, Kirchberger I, Hormann A, Kuch B. Gender-Specific short and long-term mortality in diabetic versus nondiabetic patients with incident acute myocardial infarction in the reperfusion era (the MONICA/KORA Myocardial Infarction Registry). *The American journal of cardiology* 2010;106:1680-1684.
6. Blondal M, Ainla T, Marandi T, Baburin A, Eha J. Sex-specific outcomes of diabetic patients with acute myocardial infarction who have undergone percutaneous coronary intervention: a register linkage study. *Cardiovascular diabetology* 2012;11:96.
7. Flink L, Mochari-Greenberger H, Mosca L. Gender differences in clinical outcomes among diabetic patients hospitalized for cardiovascular disease. *American heart journal* 2013;165:972-978.
8. Arboix A, Milian M, Oliveres M, Garcia-Eroles L, Massons J. Impact of female gender on prognosis in type 2 diabetic patients with ischemic stroke. *European neurology* 2006;56:6-12.
9. Eriksson M, Carlberg B, Eliasson M. The disparity in long-term survival after a first stroke in patients with and without diabetes persists: the Northern Sweden MONICA study. *Cerebrovascular diseases (Basel, Switzerland)* 2012;34:153-160.
10. Zhao W, An Z, Hong Y, et al. Sex differences in long-term outcomes among acute ischemic stroke patients with diabetes in China. *Biology of sex differences* 2015;6:29.
11. Li S, Zhao X, Wang C, et al. Risk factors for poor outcome and mortality at 3 months after the ischemic stroke in patients with atrial fibrillation. *J Stroke Cerebrovasc Dis* 2013;22:e419-425.
12. Kamal N, Lindsay MP, Cote R, Fang J, Kapral MK, Hill MD. Ten-year trends in stroke admissions and outcomes in Canada. *The Canadian journal of neurological sciences Le journal canadien des sciences neurologiques* 2015;42:168-175.
13. Kokotailo RA, Hill MD. Coding of stroke and stroke risk factors using international classification of diseases, revisions 9 and 10. *Stroke* 2005;36:1776-1781.
14. Hux JE, Ivis F, Flintoft V, Bica A. Diabetes in Ontario: determination of prevalence and incidence using a validated administrative data algorithm. *Diabetes Care* 2002;25:512-516.
15. Goldstein LB, Samsa GP, Matchar DB, Horner RD. Charlson Index comorbidity adjustment for ischemic stroke outcome studies. *Stroke* 2004;35:1941-1945.
16. Fine JP, Gray RJ. A Proportional Hazards Model for the Subdistribution of a Competing Risk. *Journal of the American Statistical Association* 1999;94:496-509.
17. Appelman Y, van Rijn BB, Ten Haaf ME, Boersma E, Peters SA. Sex differences in cardiovascular risk factors and disease prevention. *Atherosclerosis* 2015;241:211-218.

18. Samai AA, Martin-Schild S. Sex differences in predictors of ischemic stroke: current perspectives. *Vascular health and risk management* 2015;11:427-436.
19. Mosca L, Benjamin EJ, Berra K, et al. Effectiveness-based guidelines for the prevention of cardiovascular disease in women--2011 update: a guideline from the American Heart Association. *Circulation* 2011;123:1243-1262.
20. Rosen SE, Henry S, Bond R, Pearte C, Mieres JH. Sex-Specific Disparities in Risk Factors for Coronary Heart Disease. *Current atherosclerosis reports* 2015;17:49.
21. Manson JE, Bassuk SS. Biomarkers of cardiovascular disease risk in women. *Metabolism: clinical and experimental* 2015;64:S33-39.
22. Regensteiner JG, Golden S, Huebschmann AG, et al. Sex Differences in the Cardiovascular Consequences of Diabetes Mellitus: A Scientific Statement From the American Heart Association. *Circulation* 2015;132:2424-2447.
23. Nielsen AB, de Fine Olivarius N, Gannik D, Hindsberger C, Hollnagel H. Structured personal diabetes care in primary health care affects only women's HbA1c. *Diabetes care* 2006;29:963-969.
24. Schunk M, Reitmeir P, Schipf S, et al. Health-related quality of life in subjects with and without Type 2 diabetes: pooled analysis of five population-based surveys in Germany. *Diabetic medicine : a journal of the British Diabetic Association* 2012;29:646-653.
25. Strom Williams JL, Lynch CP, Winchester R, Thomas L, Keith B, Egede LE. Gender differences in composite control of cardiovascular risk factors among patients with type 2 diabetes. *Diabetes technology & therapeutics* 2014;16:421-427.
26. Wexler DJ, Grant RW, Meigs JB, Nathan DM, Cagliero E. Sex disparities in treatment of cardiac risk factors in patients with type 2 diabetes. *Diabetes care* 2005;28:514-520.
27. Seghieri C, Policardo L, Francesconi P, Seghieri G. Gender differences in the relationship between diabetes process of care indicators and cardiovascular outcomes. *European journal of public health* 2016;26:219-224.
28. Phan HT, Blizzard CL, Reeves MJ, et al. Sex Differences in Long-Term Mortality After Stroke in the INSTRUCT (INternational STROKE oUtcomes sTudy). A Meta-Analysis of Individual Participant Data 2017;10.
29. Mathur P, Ostadal B, Romeo F, Mehta JL. Gender-Related Differences in Atherosclerosis. *Cardiovascular drugs and therapy / sponsored by the International Society of Cardiovascular Pharmacotherapy* 2015;29:319-327.
30. Recarti C, Sep SJ, Stehouwer CD, Unger T. Excess cardiovascular risk in diabetic women: a case for intensive treatment. *Current hypertension reports* 2015;17:554.
31. Dantas AP, Fortes ZB, de Carvalho MH. Vascular disease in diabetic women: Why do they miss the female protection? *Experimental diabetes research* 2012;2012:570598.
32. Li G, Zhang P, Wang J, et al. Cardiovascular mortality, all-cause mortality, and diabetes incidence after lifestyle intervention for people with impaired glucose tolerance in the Da Qing Diabetes Prevention Study: a 23-year follow-up study. *The lancet Diabetes & endocrinology* 2014;2:474-480.
33. Krag MO, Hasselbalch L, Siersma V, et al. The impact of gender on the long-term morbidity and mortality of patients with type 2 diabetes receiving structured personal care: a 13 year follow-up study. *Diabetologia* 2016;59:275-285.

Table 1. Baseline characteristics of sample and numbers of outcomes*

Variable	Entire Sample	Female	Male	p-value
Number of participants	25495	11902 (46.7)	13593 (53.3)	
Baseline characteristics:				
Age, mean (SD)	73.4 (11.6)	75.6 (11.5)	71.5 (11.3)	<0.001
Neighborhood income:				<0.001
1 st quintile (lowest)	6438 (25.3)	3152 (26.5)	3286 (24.2)	
2 nd quintile	5841 (22.9)	2791 (23.4)	3050 (22.4)	
3 rd quintile	4896 (19.2)	2277 (19.1)	2619 (19.3)	
4 th quintile	4449 (17.5)	1940 (16.3)	2509 (18.5)	
5 th quintile (highest)	3871 (15.2)	1742 (14.6)	2129 (15.7)	
Diabetes duration:				0.337
0 to <3 years	5983 (23.5)	2780 (23.4)	3203 (23.6)	
3 to <6 years	3683 (14.4)	1683 (14.1)	2000 (14.7)	
≥6 years	15829 (62.1)	7439 (62.5)	8390 (61.7)	
Duration of residence in Ontario:				0.477
0 to <5 years	462 (1.8)	203 (1.7)	259 (1.9)	
5 to <10 years	629 (2.5)	291 (2.4)	338 (2.5)	
≥10 years	24404 (95.7)	11408 (95.8)	12996 (95.6)	
History of stroke or TIA	2031 (8.0)	984 (8.3)	1044 (7.7)	0.072
History of atrial fibrillation	4614 (18.1)	2501 (21.0)	2113 (15.5)	<0.001
History of hypertension	16161 (63.4)	7848 (65.9)	8313 (61.2)	<0.001
History of MI	2575 (10.1)	1105 (9.3)	1470 (10.8)	<0.001
History of CAD	5619 (22.0)	2424 (20.4)	3195 (23.5)	<0.001
History of PVD	1257 (4.9)	476 (4.0)	781 (5.7)	<0.001
Charlson index score:				0.008
0-1	6801 (26.7)	3268 (27.5)	3533 (26.0)	
≥2	18694 (73.3)	8634 (72.5)	10060 (74.0)	
Length of hospital stay, mean (SD)	16.6 (26.7)	17.8 (27.6)	15.5 (25.8)	<0.001
Overall frequency of outcomes:				
Death	12435 (48.8)	6115 (51.4)	6320 (46.5)	<0.001
Death within:				
7 days	108 (0.4)	63 (0.5)	45 (0.3)	0.015
30 days	805 (3.2)	467 (3.9)	338 (2.5)	<0.001
1 year	4517 (17.7)	2324 (19.5)	2193 (16.1)	<0.001
5 years	10916 (42.8)	5439 (45.7)	5477 (40.3)	<0.001
Readmission for any cause	17406 (68.3)	8051 (67.6)	9355 (68.8)	0.044
Readmission for any cause within:				
30 days	1706 (6.7)	767 (6.4)	939 (6.9)	0.139
1 year	9871 (38.7)	4614 (38.8)	5257 (38.7)	0.88
5 years	16402 (64.3)	7621 (64)	8781 (64.6)	0.345
Readmission for any cause or death	20135 (79.0)	9553 (80.3)	10582 (77.8)	<0.001
Readmission for any cause or death within:				
30 days	2381 (9.3)	1172 (9.8)	1209 (8.9)	0.009
1 year	11799 (46.3)	5681 (47.7)	6118 (45)	<0.001
Readmission for stroke or MI	5876 (23.0)	2647 (22.2)	3229 (23.8)	0.004
Readmission for stroke or MI within:				

30 days	538 (2.1)	245 (2.1)	293 (2.2)	0.591
1 year	2591 (10.2)	1196 (10)	1395 (10.3)	0.573
5 years	5236 (20.5)	2372 (19.9)	2864 (21.1)	0.025
Readmission for stroke	3794 (14.9)	1791 (15)	2003 (14.7)	0.484
Readmission for stroke within:				
30 days	472 (1.9)	212 (1.8)	260 (1.9)	0.437
1 year	1808 (7.1)	867 (7.3)	941 (6.9)	0.262
5 years	3429 (13.4)	1620 (13.6)	1809 (13.3)	0.48
Readmission for MI	2512 (9.9)	1036 (8.7)	1476 (10.9)	<0.001
Readmission for MI within:				
30 days	66 (0.3)	33 (0.3)	33 (0.2)	0.589
1 year	844 (3.3)	360 (3)	484 (3.6)	0.017
5 years	2149 (8.4)	899 (7.6)	1250 (9.2)	<0.001
CABG or PCI	1076 (4.2)	333 (2.8)	743 (5.5)	<0.001
Baseline medications for those aged ≥65 yr:				
Number of participants aged ≥65 yr	19619 (76.9)	9788 (82.2)	9831 (72.3)	
Diabetic medications	11656 (59.4)	5655 (57.8)	6001 (61)	<0.001
Statin medications	12724 (64.9)	6096 (62.3)	6628 (67.4)	<0.001
Warfarin	4248 (21.7)	2189 (22.4)	2059 (20.9)	0.016
Anti-hypertensive medications	15778 (80.4)	7923 (80.9)	7855 (79.9)	0.065

*values are number and column percentages in parentheses unless otherwise indicated; TIA=transient ischemic attack; MI=myocardial infarction; CAD=coronary artery disease; PVD=peripheral vascular disease; CABG=coronary artery bypass graft surgery; PCI=percutaneous coronary intervention; NA=specific values not reported due to identifiability with small cell sizes; IQR=interquartile range

Table 2. Incidence rate of outcomes per 100 person-years, by subgroups of sex and age

Outcome	Overall rate (95% CI)	Rate among females (95% CI)		Rate among males (95% CI)	p-value
By sex:					
Death	12.88 (12.65-13.11)	14.08 (13.73-14.44)		11.89 (11.60-12.19)	<0.0001
Readmission for any cause	26.60 (26.00-27.22)	26.20 (25.33-27.10)		26.95 (26.12-27.81)	0.2
Readmission for any cause or death	39.30 (38.49-40.13)	41.71 (40.47, 43.00)		37.35 (36.29-38.44)	<0.0001
Readmission for stroke	2.86 (2.77-2.95)	2.90 (2.76-3.03)		2.82 (2.70-2.95)	0.4
Readmission for MI	1.80 (1.73-1.87)	1.58 (1.49-1.68)		1.99 (1.89-2.10)	<0.0001
Readmission for stroke or MI	4.77 (4.65-4.90)	4.59 (4.42-4.77)		4.94 (4.77-5.11)	0.006
By age group:					
	Overall rate (95% CI)	Rate among 18-64 yr olds (95% CI)	Rate among 65-79 yr olds (95% CI)	Rate among 80+ yr olds (95% CI)	p-value
Death	12.88 (12.65-13.11)	4.56 (4.31-4.82)	11.20 (10.89-11.51)	24.67 (24.06-25.29)	<0.0001
Readmission for any cause	26.60 (26.00-27.22)	21.02 (19.94-22.15)	28.85 (27.88-29.85)	27.81 (26.77-28.89)	<0.0001
Readmission for any cause or death	39.30 (38.49-40.13)	23.39 (22.27-24.57)	37.73 (36.57-38.93)	59.57 (57.67-61.53)	<0.0001
Readmission for stroke	2.86 (2.77-2.95)	3.12 (2.92-3.34)	2.95 (2.82-3.09)	2.58 (2.44-2.73)	<0.0001
Readmission for MI	1.80 (1.73-1.87)	2.17 (2.00-2.34)	2.02 (1.91-2.14)	1.32 (1.22-1.42)	<0.0001
Readmission for stroke or MI	4.77 (4.65-4.90)	5.34 (5.07-5.63)	5.13 (4.94-5.32)	4.03 (3.84-4.22)	<0.0001

*MI=myocardial infarction; CI=confidence interval

Table 3. Multivariable models of outcomes

Outcome Variable	Competing risk models with death as competing risk								Cox regression	
	Any event		Stroke		MI		Stroke or MI		Death	
	HR (95% CI)	p-value	HR (95% CI)	p-value	HR (95% CI)	p-value	HR (95% CI)	p-value	HR (95% CI)	p-value
Whole sample**										
Age	1.01 (1.00-1.01)	<0.0001	0.99 (0.99-0.99)	<0.0001	0.99 (0.98-0.99)	<0.0001	0.99 (0.99-0.99)	<0.0001	1.06 (1.06-1.06)	<0.0001
Female sex†	0.96 (0.93- 0.99)	0.004	1.05 (0.98-1.12)	0.2	0.88 (0.81-0.95)	0.001	0.98 (0.93-1.03)	0.4	0.95 (0.92-0.99)	0.01
Among those ≥65 yr without adjustment for medications**										
Age	1.00 (0.99-1.00)	0.02	0.98 (0.98-0.99)	<0.0001	0.97 (0.96-0.98)	<0.0001	0.98 (0.98-0.98)	<0.0001	1.07 (1.06-1.07)	<0.0001
Female sex†	0.94 (0.91-0.97)	0.0006	1.06 (0.98-1.14)	0.1	0.93 (0.84-1.02)	0.1	1.01 (0.95-1.07)	0.7	0.93 (0.90-0.97)	0.0005
Among those ≥65 yr with adjustment for medications[@]										
Age	1.00 (1.00-1.00)	0.6	0.99 (0.98-0.99)	<0.0001	0.98 (0.97-0.98)	<0.0001	0.98 (0.98-0.99)	<0.0001	1.06 (1.06-1.06)	0.0001
Female sex†	0.94 (0.91-0.97)	0.0004	1.06 (0.98-1.14)	0.1	0.92 (0.84-1.02)	0.1	1.01 (0.95-1.07)	0.8	0.93 (0.89-0.97)	0.0002

MI=myocardial infarction; HTN=hypertension; CAD=coronary artery disease; PVD=peripheral vascular disease; †male sex as referent;**models are adjusted for: income, hypertension, atrial fibrillation, stroke or TIA, MI, CAD, PVD, and Charlson score; [@]models are adjusted for: income, hypertension, atrial fibrillation, stroke or TIA, MI, CAD, PVD, Charlson score, anti-hypertensive medication use, diabetes medication use, statin use, and warfarin use

Figure 1A: Kaplan-Meier survival curve stratified by sex




Figure 1B: Kaplan-Meier curves of probability of survival free of readmission, stratified by sex




Figure 1C: Kaplan-Meier curves of probability of survival free of readmission or death, stratified by sex




Figure 2A: Kaplan-Meier curves of probability of survival free of stroke readmission, stratified by sex




Figure 2B: Kaplan-Meier curves of probability of survival free of myocardial infarction readmission, stratified by sex




Figure 2C: Kaplan-Meier curves of probability of survival free of stroke or myocardial infarction readmission, stratified by sex

