

11-11-2021

## Surgical Treatment of Femoroacetabular Impingement: Minimum 10-Year Outcome and Risk Factors for Failure

Hamed Vahedi

Steven Yacovelli

Claudio Diaz

Javad Parvizi

Follow this and additional works at: [https://jdc.jefferson.edu/rothman\\_institute](https://jdc.jefferson.edu/rothman_institute)



Part of the [Orthopedics Commons](#), and the [Surgery Commons](#)

**[Let us know how access to this document benefits you](#)**

---

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Rothman Institute Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

# Surgical Treatment of Femoroacetabular Impingement

## Minimum 10-Year Outcome and Risk Factors for Failure

Hamed Vahedi, MD, Steven Yacovelli, Claudio Diaz, MD, and Javad Parvizi, MD, FRCS

*Investigation performed at the Rothman Orthopaedic Institute at Thomas Jefferson University, Philadelphia, Pennsylvania*

**Background:** Femoroacetabular impingement (FAI) is a well-known cause of hip pain and dysfunction in young adults. Surgical treatment has been widely popularized during the past decade. However, most reported results have been limited to short-term and intermediate-term follow-up. The long-term success rate and risk factors for failure are largely unknown. This study aimed to report the long-term (minimum, 10 years) clinical outcomes of surgical treatment of FAI and to describe the clinical and radiographic parameters associated with the failure of treatment involving femoroacetabular osteoplasty (FAO) and labral repair.

**Methods:** Using our prospective hip preservation database, 164 patients (178 hips) who had undergone FAO between January 2005 and April 2009 were identified. Patient demographic characteristics, clinical history, duration of preoperative symptoms, radiographic parameters (preoperative and postoperative alpha angles, hip dysplasia and retroversion, Tönnis grade for osteoarthritis), and intraoperative findings were reviewed and compared between the success and failure groups. At a minimum 10-year follow-up, clinical functional outcomes (modified Harris hip score [mHHS] and Short Form-36 [SF-36] at 6 weeks, 6 months, 1 year, and 10 years) and failure rates (conversion to total hip arthroplasty [THA]) were collected.

**Results:** The mean patient age (and standard deviation) was  $34.3 \pm 8.4$  years, and 65 patients (40%) were female. After the surgical procedure, there was significant improvement in the mean mHHS ( $59.3 \pm 7.3$  points preoperatively to  $88.4 \pm 7.3$  points postoperatively) and the mean SF-36 ( $61.3 \pm 8.4$  points preoperatively to  $89.1 \pm 7.2$  points postoperatively). At a mean follow-up of 12.5 years, 12% (22 hips) required conversion to THA, with a mean time to THA of  $5.3 \pm 2.0$  years. Older age, longer preoperative symptomatic period, higher preoperative and postoperative alpha angles, presence of hip dysplasia, a higher Tönnis grade, joint space narrowing, and a full-thickness acetabular chondral lesion at the time of the FAO were identified as risk factors for failure and conversion to THA.

**Conclusions:** Patients with symptomatic FAI who undergo a surgical procedure experience pain relief and functional improvement that appear to endure over a decade in the majority of patients. This study on a relatively large cohort with a long-term follow-up has also identified patients who are at a higher risk for treatment failure.

**Level of Evidence:** Therapeutic Level IV. See Instructions for Authors for a complete description of levels of evidence.

Femoroacetabular impingement (FAI) and its well-described underlying structural hip abnormality are an established source of hip pain and dysfunction in active young adults that can eventually lead to hip joint arthritis<sup>1-3</sup>. Although the exact etiology of FAI has yet to be elucidated, multiple risk factors, both genetic and environmental, have been reported and likely play a role<sup>3,4</sup>. Previous studies have shown that early-stage surgical intervention in patients with symptomatic FAI with no or minimal evidence of degenerative hip disease may help to increase the longevity of the articular cartilage<sup>2,5-7</sup>. For these reasons, coupled with the improvement in diagnosis and understanding of

the disease pathology, the surgical treatment of FAI has gained increasing popularity over the past decade.

The first described technique used for femoroacetabular osteoplasty (FAO) and labral repair was surgical hip dislocation<sup>8</sup>. However, because of its relatively high complication rate, lengthy learning curve, and complexity, multiple other surgical approaches have since been described<sup>9,10</sup>. Still, the goal of surgical intervention remains the same: to address the femoral head-neck junction and acetabular abnormality based on the source and etiology of the impingement. Eliminating the source of impingement may prevent or delay further degenerative

**Disclosure:** The **Disclosure of Potential Conflicts of Interest** forms are provided with the online version of the article (<http://links.lww.com/JBJSOA/A346>).

Copyright © 2021 The Authors. Published by The Journal of Bone and Joint Surgery, Incorporated. All rights reserved. This is an open-access article distributed under the terms of the [Creative Commons Attribution-Non Commercial-No Derivatives License 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/) (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

changes and increase the symptom-free, prearthritic period in these patients<sup>1,2,6-8,11-16</sup>.

The short-term and intermediate-term clinical studies of arthroscopic FAO and risk factors for treatment failure have been reported previously, with favorable outcomes<sup>13,16-20</sup>. The senior author of this study (J.P.) has similarly reported excellent short-term to intermediate-term outcomes with his preferred technique in the past<sup>21,22</sup>. Some of the identified risk factors for treatment failure described thus far include preexisting arthritis, older age at the time of the surgical procedure, longer duration of preoperative symptoms, higher body mass index (BMI), and joint space narrowing<sup>12,13,17,18,21-27</sup>.

The primary aim of this study was to report the long-term (minimum 10-year) clinical outcomes of the surgical treatment of FAI. The secondary purpose was to describe the clinical and radiographic parameters associated with failure of treatment involving FAO and labral repair.

## Materials and Methods

Following institutional review board approval, our prospective hip preservation database was utilized to identify patients who had undergone FAO between January 2005 and April 2009 and had a minimum 10-year follow-up. Patients in the initial cohort of 197 hips who had a definitive diagnosis of hip dysplasia (8 hips), history of slipped capital femoral epiphysis (5 hips), osteonecrosis of the femoral head (3 hips), or previous femoral head or neck or acetabular fracture (3 hips) were excluded. The final cohort consisted of 164 patients (178 hips). None of the patients underwent labral debridement as part of the procedure. The mean patient age (and standard deviation) was  $34.3 \pm 8.4$  years, and 65 patients (40%) were female. Patient demographic characteristics, duration of preoperative symptoms, radiographic parameters (preoperative and postoperative alpha angles, hip dysplasia and retroversion parameters, joint space, Tönnis grade for osteoarthritis), and intraoperative findings were collected and reviewed. Borderline hip dysplasia was described as a lateral central-edge angle (LCEA) or an anterior center-edge angle (ACEA) between  $15^\circ$  and  $20^\circ$  and Tönnis angle between  $10^\circ$  and  $15^\circ$ . A false-profile view of the involved hip was used for the measurement of the ACEA. Acetabular retroversion was deemed to be present when the crossover sign, posterior wall sign, and prominent ischial spine sign were present in the anteroposterior pelvic radiographic view.

Definite failure of FAO was defined as the need for a conversion to a total hip arthroplasty (THA) at any time. Preoperative and postoperative clinical and radiographic findings were compared between the success and failure groups. Also, at a minimum 10-year follow-up, functional outcome scores (modified Harris hip score [mHHS] and Short Form-36 [SF-36]) were collected before the FAO, before the THA, and at 6 weeks, 6 months, 1 year, and 10 years after the index procedure. A single surgeon performed all of the index FAOs (using a mini-open technique) and all conversions to THAs in symptomatic hips in the failure group.

## Surgical Technique

All patients were positioned supine on a regular operating table. A 3-cm-long incision was made distal to the anterior

superior iliac spine over the tensor fasciae latae muscle. After retracting the tensor muscle laterally, the rectus muscle was reflected laterally to expose the hip capsule. To expose the medial capsule, the sartorius muscle needed to be retracted medially. Then the cam deformity was visualized through an I-shaped capsulotomy. The labrum was checked for evidence of the tear. To visualize the weight-bearing surface of the joint for any evidence of chondral lesions, minimal traction was applied. Most of the articular surface can be evaluated through this maneuver. A blunt-tipped hook was utilized to feel the posteroinferior surface of the acetabulum. Microfracture was performed for full-thickness chondral lesions, and partial-thickness lesions were excised. In patients with excessive coverage, acetabuloplasty was employed. However, in patients with evidence of acetabular dysplasia and undercoverage, very minimal removal of the acetabular rim was done to refresh the bed for labral repair. Then femoral neck osteoplasty was performed using a burr and small osteotome until achieving an impingement-free hip range of motion. After femoral neck osteoplasty, labral tear repair was done by the placement of anchor sutures in the acetabular rim. After a final check of the impingement-free range of motion, the joint capsule was closed using a running suture. Postoperatively, patients were allowed partial weight-bearing for 6 weeks, followed by full weight-bearing. Return to full activity and sports took 4 to 6 months for most cases<sup>28</sup>.

## Statistical Analysis

A bivariate analysis of the data was completed to assess the relationship of survival and failure with variables of interest. All continuous parametric data are presented as the mean and the standard deviation, and all categorical data are presented as the frequency and percentage. The p values for continuous data were calculated using the t test, and those for categorical data were calculated using the chi-square test. Following the bivariate analysis, a Cox regression was computed to analyze the relationship between key variables and time to failure. This regression helped us to identify key variables that relate to treatment failure. All statistical analyses were performed using RStudio, version 3.5.1 (The R Foundation for Statistical Computing).

## Source of Funding

None.

## Results

At the latest follow-up at a mean of 12.5 years (range, 10 to 14 years), 22 hips (12%) required conversion to THA, with a mean time to THA of  $5.3 \pm 2.0$  years. The remaining hips demonstrated favorable outcomes and did not require a THA (Fig. 1). Patients who underwent failed treatment were significantly older at  $41.0 \pm 7.11$  years of age at the time of the FAO compared with  $30.8 \pm 8.39$  years of age for patients who underwent successful treatment ( $p < 0.001$ ). No significant difference was detected in terms of sex and BMI between the groups. The failure cohort had a longer symptomatic period before the FAO procedure ( $28.2 \pm 11.4$  compared with  $18.8 \pm 7.41$  months;  $p = 0.001$ ). Borderline dysplasia was present in 7

hips (32%) in the failure group and 10 hips (6%) in the success group ( $p = 0.001$ ). A total of 8 patients had evidence of acetabular retroversion, of whom 3 underwent conversion to THA ( $p = 0.061$ ). A higher rate of full-thickness acetabular chondral lesions (32% compared with 3%) was identified in the failure cohort ( $p < 0.001$ ). Of the other radiographic parameters measured, joint space narrowing, higher Tönnis angle, lower ACEA, higher Tönnis grade of arthritis, and higher preoperative and postoperative alpha angles were found to be associated with failure. The details of the demographic characteristics and perioperative patient-related data and outcome scores are listed in Table I.

After adjusting for covariates (Table II), by using the Cox proportional hazard model, the following variables were independently associated with a higher incidence of conversion to THA after the index FAO: older age at the time of the index surgical procedure, prolonged preoperative symptomatic period, borderline dysplasia of the hip, joint space narrowing before the FAO, presence of a full-thickness chondral lesion, and higher preoperative and postoperative alpha angles.

Significant improvements in clinical functional outcomes were observed in the cohort at the 6-month, 1-year, and 10-year follow-ups after the FAO. At the latest follow-up (minimum, 10 years), the mean mHHS was  $88.4 \pm 7.34$  points in the survival group. The mean mHHS after the THA was  $92.3 \pm 5.72$  points at the latest follow-up. None of the hips underwent revision in the first 2 years after the FAO. At the 2-year follow-up, the mean mHHS was  $90.3 \pm 6.9$  points and the mean SF-36 was  $89.6 \pm 7.4$  points.

## Discussion

The current study shows the long-term clinical results, survivorship, and risk factors for the failure of surgical treatment of FAI. The surgical outcome of FAI was deemed to be acceptable, as a relatively low number of patients (12%) eventually required a THA, with the other patients appearing to have excellent clinical functional outcomes at the latest follow-

up. As expected, older age at the time of the index surgical procedure, prolonged preoperative symptomatic period, presence of borderline dysplasia of the hip, joint space narrowing, presence of full-thickness chondral lesions, and higher preoperative and postoperative alpha angles were identified as independent variables associated with higher rates of failure, as defined by the need for conversion to THA.

Hip preservation surgery has gained popularity in the past decade, mostly because of its effectiveness in alleviating symptoms and delaying the need for a THA in symptomatic young patients. Favorable intermediate-term results of FAO and labral repair using various open or arthroscopic techniques have been reported in previous studies<sup>13,16-19</sup>. However, there has been a lack of evidence available with regard to long-term functional outcomes and failure rates of FAO in the literature. Menge et al.<sup>28</sup> reported survivorship and outcomes at 10 years after arthroscopic FAO in a retrospective study of 145 patients. The study found a much higher conversion rate at 34% (50 of 145 hips). Although it is difficult to determine the exact reason for the lower treatment failure rate in our cohort, the presence of a higher rate of preoperative joint space narrowing in the cohort reported by Menge et al. may have contributed to the higher rate of treatment failure. The risk factors for failure were also identified by Menge et al. and were found to be older age, joint space of  $<2$  mm, and the need for microfracture during osteoplasty. Our study corroborates the findings of the study by Menge et al. and identifies a few more risk factors for failure.

Clinical outcomes and predictive factors for failure of hip arthroscopy in patients with FAI have been investigated in other studies as well, but mostly at short-term and intermediate-term follow-up<sup>12,13,18,19,22-27,29,30</sup>. Furthermore, the definition of failure (functional outcome, conversion to THA, revision FAO), the sample size, and the type of procedure (open, arthroscopic) differed vastly among these studies, leading to a wide range of results<sup>31</sup>.

The goals of any type of hip preservation surgery are to prevent or delay the need for THA and to provide patients with symptom-free function with the native hip. For this reason,




Fig. 1  
Kaplan-Meier survivorship of patients treated with FAO. The shaded areas indicate the 95% confidence interval. Conversion to THA was used as the definition of treatment failure.

TABLE I Patient Demographic Characteristics, Perioperative Radiographic Measurements, and Functional Outcome Scores\*

	Survival (N = 156)	Failure (N = 22)	P Value
Demographic characteristics			
Age† (yr)	30.8 ± 8.39	41.0 ± 7.11	<0.001‡
Sex§			0.488
Male	101 (65%)	12 (55%)	
Female	55 (35%)	10 (45%)	
BMI† (kg/m <sup>2</sup> )	25.9 ± 2.62	25.8 ± 2.47	0.763
Preoperative symptomatic period† (mo)	18.8 ± 7.41	28.2 ± 11.4	0.001‡
Time to THA† (yr)	NA	5.30 ± 1.99	
Radiographic measurements			
Borderline dysplasia§	10 (6%)	7 (32%)	0.001‡
Retroversion§	5 (3%)	3 (14%)	0.061
Full-thickness chondral lesion§	5 (3%)	7 (32%)	<0.001‡
Preoperative alpha angle† (deg)	68.9 ± 8.44	84.7 ± 10.3	<0.001‡
Postoperative alpha angle† (deg)	56.6 ± 5.43	59.5 ± 6.52	0.056
Tönnis grade§			<0.001‡
0	110 (71%)	3 (14%)	
1	41 (26%)	12 (55%)	
2	5 (3%)	7 (32%)	
LCEA† (deg)	27.8 ± 5.29	25.5 ± 7.51	0.183
ACEA† (deg)	27.4 ± 4.08	24.5 ± 5.62	0.028‡
Tönnis angle† (deg)	2.35 ± 2.71	4.64 ± 3.81	0.012‡
Medial joint space† (mm)	4.87 ± 0.80	4.36 ± 1.26	0.081‡
Superior joint space† (mm)	4.83 ± 0.68	3.59 ± 0.67	<0.001‡
Functional outcome score† (points)			
mHHS			
Before FAO	59.3 ± 7.29	53.1 ± 8.50	0.003‡
Before THA	NA	42.3 ± 12.3	
Latest follow-up	88.4 ± 7.34	92.3 ± 5.72	0.007‡
SF-36			
Before FAO	61.3 ± 8.39	52.4 ± 9.55	<0.001‡
Before THA	NA	47.6 ± 11.2	
Latest follow-up	89.1 ± 7.15	87.7 ± 8.21	0.470

\*NA = not applicable. †The values are given as the mean and the standard deviation. ‡Significant. §The values are given as the number of hips, with the percentage in parentheses.

similar to many other previous hip preservation studies, we decided to use conversion to THA as the definition of treatment failure<sup>12,14,23-26,32-35</sup>. With regard to risk factors for failure, previous short-term to intermediate-term follow-up studies have identified older age, female sex, higher BMI, longer duration of preoperative symptoms, presence of chondral problems, joint space narrowing, and higher Tönnis grade to be associated with poor outcomes<sup>12-14,23-26,32-36</sup>. In a recently published study by Nho et al.<sup>36</sup>, Tönnis grade of ≥1, Workers' Compensation, and higher BMI were identified as risk factors for a less optimal outcome in a large multicenter cohort of 935 patients who underwent hip arthroscopy and had a minimum 2-year follow-up. The rate of conversion to THA was 1.2% (11 cases), and 2.5% (23 cases) underwent revision arthroscopy. Also, 27.4% (256 cases) of their

cohort reported inferior clinical outcomes and did not reach a minimal clinically important difference in outcome scores postoperatively. Their cohort consisted of a larger number of patients but a shorter follow-up after the surgical procedure compared with our patient population. Also, they excluded patients with borderline dysplasia, those with a Tönnis grade of >1, and patients with a history of contralateral hip surgery.

Similarly, as reported in other studies, older age at the time of the FAO was associated with a higher rate of conversion to THA in our study<sup>12,14,16-20,26,30,31,34,37</sup>. This might be due to less optimal quality of the cartilage in older patients and a longer period of abnormal hip biomechanics before the onset of symptoms. Peters et al.<sup>37</sup> concluded that the incidence of failure of hip preservation procedures increases with a rate of 4% per year.

TABLE II Cox Proportional Hazard Model for Outcome of Failure

Covariate	Hazard Ratio*	P Value
Age in yr	0.85 (0.73 to 1.00)	0.048†
Female sex	5.69 (0.08 to 41.04)	0.426
Preoperative symptomatic period in yr	1.22 (1.07 to 1.40)	0.004†
Borderline dysplasia	$2.698 \times 10^5$ (2.50 to $2.9 \times 10^{10}$ )	0.034†
Retroversion	94.02 (0.11 to 7,717)	0.184
Chondral lesion	1,761 (8.73 to $36 \times 10^7$ )	0.011†
Preoperative alpha angle in degrees	1.37 (1.07 to 1.74)	0.011†
Postoperative alpha angle in degrees	1.28 (1.01 to 1.62)	0.045†
Tönnis grade	0.59 (0.04 to 9.08)	0.708
ACEA in degrees	1.67 (0.95 to 2.93)	0.073
Tönnis angle in degrees	0.37 (0.09 to 1.47)	0.158
Medial joint space in mm	0.13 (0.03 to 0.45)	0.002†
Superior joint space in mm	377.7 (12.99 to $1.1 \times 10^6$ )	0.004†

\*The values are given as the hazard ratio, with the 95% confidence interval in parentheses. †Significant.

Measures of the quality of the hip-joint bearing surface, including the status of articular cartilage, the amount of initial cartilage loss, the presence of degenerative changes, and joint space narrowing, were other important risk factors for long-term treatment failure in our patients. The presence of full-thickness delamination of articular cartilage has been shown to adversely affect the outcome of FAO<sup>10,11,30</sup>. In a systematic review by Kemp et al.<sup>20</sup>, a strong association was reported between the presence of severe articular cartilage damage and conversion to THA following FAI procedures. Also, preexisting degenerative joint changes categorized by Tönnis grade were another strong predictor of failure<sup>5,12,24-26,32,35</sup>. Based on current evidence, it is not clear that performing FAO on hips with evidence of arthritis should be attempted before proceeding to a discussion of THA with patients. Based on the current body of evidence, we do not believe that FAO and labral repair are indicated in patients with advanced hip arthritis.

Poor outcomes after hip preservation surgery in patients with high BMI have been reported in some studies. Gupta et al.<sup>16</sup> demonstrated that obese patients in their cohort (680 patients) started with lower absolute scores preoperatively and ended with lower absolute postoperative scores at the 2-year follow-up. Our findings differed in that the failure and survival groups had similar mean BMIs, possibly due to the fact that our patient population had a low mean BMI overall.

Another important identified risk factor for treatment failure was a longer preoperative symptomatic period. Some previous studies also mentioned a less optimal outcome in patients with a longer symptomatic period before undergoing an FAO<sup>25,26</sup>. In a study by Kunze et al.<sup>38</sup>, superior postoperative outcomes were observed in patients with early surgical intervention, specifically within 6 months from the onset of symptoms. In multivariate regression analysis, symptom duration was predictive of the visual

analog scales for pain and satisfaction. We agree with this notion and belief that early surgical intervention after failed nonoperative treatment (physical therapy, nonsteroidal anti-inflammatory drugs [NSAIDs], activity modification) can improve the outcome. Theoretically, a shorter duration of abnormal hip biomechanics may cause less permanent damage to the joint surfaces and may slow down the progression of irreversible changes.

Among the various radiographic parameters measured in our patients, borderline dysplasia, lower ACEA, joint space narrowing, higher Tönnis angle, and higher preoperative and postoperative alpha angles were detected more commonly in our failure group. We recently published our results of FAO in patients with hip dysplasia<sup>39</sup>. At the latest follow-up, 28.8% of 73 dysplastic hips eventually underwent a THA compared with 2.5% in the control group with no evidence of hip dysplasia (n = 550). More extensive preoperative discussion should be conducted with patients who have evidence of hip dysplasia with regard to the expected outcome (symptom and survival) of the native hip postoperatively. In this scenario, one may consider periacetabular osteotomy in isolation (as a definitive treatment) or in combination with FAO in patients with evidence of hip dysplasia.

Although a higher incidence of risk factors for treatment failure was detected in patients who underwent THA, a relatively high number of patients at moderate to high risk, with preexisting chondral damage, mild to moderate arthritis with some degree of joint space narrowing, or evidence of borderline hip dysplasia, are still enjoying excellent function after a minimum of 10 years after the FAO.

Our study had some limitations. Regardless of the fact that we collect data prospectively on patients undergoing hip preservation at our institution, some limitations related to this study's retrospective nature remained. In addition, we did not



investigate some other variables that may have been related to poor outcomes, such as socioeconomic status, smoking, race, activity level, and Workers' Compensation. Also, we did not use poor functional and patient-reported outcomes as one of our failure criteria. To our knowledge, there is no gold standard for the definition of a poor or less-optimal clinical outcome score. Therefore, we decided to use conversion to THA as an absolute definition of treatment failure. Also, the low number of hips in the failure group may decrease the external validity of reported risk factors for treatment failure.

However, it appears that the majority of patients with symptomatic FAI who undergo a surgical procedure experience pain relief and functional improvement that lasts over a decade. This study also identified patients who are at higher risk for treatment failure. A majority of patients experienced favorable outcomes and did not undergo THA at the latest follow-up. We believe that the findings will be of value to hip preservation

surgeons in careful patient selection and more informative preoperative discussions with patients and their families with regard to the prognosis and outcome of surgical intervention. ■

Hamed Vahedi, MD<sup>1</sup>  
Steven Yacovelli<sup>2</sup>  
Claudio Diaz, MD<sup>2</sup>  
Javad Parvizi, MD, FRCS<sup>2</sup>

<sup>1</sup>West Virginia University Medicine, Morgantown, West Virginia

<sup>2</sup>Rothman Orthopaedic Institute at Thomas Jefferson University, Philadelphia, Pennsylvania

Email for corresponding author: javadparvizi@gmail.com

## References

- Clohisy JC, St John LC, Schutz AL. Surgical treatment of femoroacetabular impingement: a systematic review of the literature. *Clin Orthop Relat Res*. 2010 Feb; 468(2):555-64.
- Ganz R, Leunig M, Leunig-Ganz K, Harris WH. The etiology of osteoarthritis of the hip: an integrated mechanical concept. *Clin Orthop Relat Res*. 2008 Feb;466(2): 264-72. Epub 2008 Jan 10.
- Wenger DR, Kishan S, Pring ME. Impingement and childhood hip disease. *J Pediatr Orthop B*. 2006 Jul;15(4):233-43.
- Philippon MJ, Schenker ML. Arthroscopy for the treatment of femoroacetabular impingement in the athlete. *Clin Sports Med*. 2006 Apr;25(2):299-308: ix.
- Beck M, Kalhor M, Leunig M, Ganz R. Hip morphology influences the pattern of damage to the acetabular cartilage: femoroacetabular impingement as a cause of early osteoarthritis of the hip. *J Bone Joint Surg Br*. 2005 Jul;87(7): 1012-8.
- Ng VY, Arora N, Best TM, Pan X, Ellis TJ. Efficacy of surgery for femoroacetabular impingement: a systematic review. *Am J Sports Med*. 2010 Nov;38(11):2337-45. Epub 2010 May 20.
- Ganz R, Parvizi J, Beck M, Leunig M, Nötzli H, Siebenrock KA. Femoroacetabular impingement: a cause for osteoarthritis of the hip. *Clin Orthop Relat Res*. 2003 Dec; 417:112-20.
- Ganz R, Gill TJ, Gautier E, Ganz K, Krügel N, Berlemann U. Surgical dislocation of the adult hip a technique with full access to the femoral head and acetabulum without the risk of avascular necrosis. *J Bone Joint Surg Br*. 2001 Nov;83(8):1119-24.
- Matsuda DK, Carlisle JC, Arthurs SC, Wierks CH, Philippon MJ. Comparative systematic review of the open dislocation, mini-open, and arthroscopic surgeries for femoroacetabular impingement. *Arthroscopy*. 2011 Feb;27(2):252-69.
- Nwachukwu BU, Rebolledo BJ, McCormick F, Rosas S, Harris JD, Kelly BT. Arthroscopic versus open treatment of femoroacetabular impingement: a systematic review of medium- to long-term outcomes. *Am J Sports Med*. 2016 Apr;44(4):1062-8. Epub 2015 Jun 9.
- Beaulé PE, Le Duff MJ, Zaragoza E. Quality of life following femoral head-neck osteochondroplasty for femoroacetabular impingement. *J Bone Joint Surg Am*. 2007 Apr;89(4):773-9.
- Beck M, Leunig M, Parvizi J, Boutier V, Wyss D, Ganz R. Anterior femoroacetabular impingement: part II. Midterm results of surgical treatment. *Clin Orthop Relat Res*. 2004 Jan;418:67-73.
- Laude F, Soriali E, Nogier A. Femoroacetabular impingement treatment using arthroscopy and anterior approach. *Clin Orthop Relat Res*. 2009 Mar;467(3):747-52. Epub 2008 Dec 16.
- Philippon MJ, Ejnisman L, Ellis HB, Briggs KK. Outcomes 2 to 5 years following hip arthroscopy for femoroacetabular impingement in the patient aged 11 to 16 years. *Arthroscopy*. 2012 Sep;28(9):1255-61. Epub 2012 May 4.
- Philippon MJ, Schroder E Souza BG, Briggs KK. Hip arthroscopy for femoroacetabular impingement in patients aged 50 years or older. *Arthroscopy*. 2012 Jan;28(1):59-65. Epub 2011 Oct 7.
- Gupta A, Redmond JM, Hammarstedt JE, Lindner D, Stake CE, Domb BG. Does obesity affect outcomes after hip arthroscopy? A cohort analysis. *J Bone Joint Surg Am*. 2015 Jan 7;97(1):16-23.
- Stappacher SD, Huemmer C, Schwab JM, Tannast M, Siebenrock KA. Surgical hip dislocation for treatment of femoroacetabular impingement: factors predicting 5-year survivorship. *Clin Orthop Relat Res*. 2014 Jan;472(1):337-48. Epub 2013 Sep 8.
- Boyer T, Dorfmann H. Arthroscopy in primary synovial chondromatosis of the hip: description and outcome of treatment. *J Bone Joint Surg Br*. 2008 Mar;90(3):314-8.
- McCarthy JC, Jarrett BT, Ojeifo O, Lee JA, Bragdon CR. What factors influence long-term survivorship after hip arthroscopy? *Clin Orthop Relat Res*. 2011 Feb; 469(2):362-71.
- Kemp JL, MacDonald D, Collins NJ, Hatton AL, Crossley KM. Hip arthroscopy in the setting of hip osteoarthritis: systematic review of outcomes and progression to hip arthroplasty. *Clin Orthop Relat Res*. 2015 Mar;473(3):1055-73. Epub 2014 Sep 18.
- Ceylan HH, Vahedi H, Azboy I, Aali Rezaie A, Parvizi J. Mini-open femoroacetabular osteoplasty: risk factors for failure and conversion to hip arthroplasty. *J Bone Joint Surg Am*. 2020 Jun 17;102(12):e59.
- Beck M, Leunig M, Parvizi J, Boutier V, Wyss D, Ganz R. Anterior femoroacetabular impingement: part II. Midterm results of surgical treatment. *Clin Orthop Relat Res*. 2004 Jan;418:67-73.
- Gedouin JE, May O, Bonin N, Nogier A, Boyer T, Sadri H, Villar RN, Laude F; French Arthroscopy Society. Assessment of arthroscopic management of femoroacetabular impingement. A prospective multicenter study. *Orthop Traumatol Surg Res*. 2010 Dec;96(8)(Suppl):S59-67. Epub 2010 Oct 28.
- Horisberger M, Brunner A, Herzog RF. Arthroscopic treatment of femoral acetabular impingement in patients with preoperative generalized degenerative changes. *Arthroscopy*. 2010 May;26(5):623-9. Epub 2010 Feb 11.
- Larson CM, Giveans MR, Stone RM. Arthroscopic debridement versus refixation of the acetabular labrum associated with femoroacetabular impingement: mean 3.5-year follow-up. *Am J Sports Med*. 2012 May;40(5):1015-21. Epub 2012 Feb 3.
- Philippon MJ, Briggs KK, Yen YM, Kuppertschmidt DA. Outcomes following hip arthroscopy for femoroacetabular impingement with associated chondrolabral dysfunction: minimum two-year follow-up. *J Bone Joint Surg Br*. 2009 Jan;91(1):16-23.
- Byrd JWT, Jones KS. Prospective analysis of hip arthroscopy with 10-year followup. *Clin Orthop Relat Res*. 2010 Mar;468(3):741-6. Epub 2009 Apr 21.
- Menge TJ, Briggs KK, Dorman GJ, McNamara SC, Philippon MJ. Survivorship and outcomes 10 years following hip arthroscopy for femoroacetabular impingement: labral debridement compared with labral repair. *J Bone Joint Surg Am*. 2017 Jun 21; 99(12):997-1004.
- Farjo LA, Glick JM, Sampson TG. Hip arthroscopy for acetabular labral tears. *Arthroscopy*. 1999 Mar;15(2):132-7.
- Margheritini F, Villar RN. The efficacy of arthroscopy in the treatment of hip osteoarthritis. *Chir Organi Mov*. 1999 Jul-Sep;84(3):257-61.
- Saadat E, Martin SD, Thornhill TS, Brownlee SA, Losina E, Katz JN. Factors associated with the failure of surgical treatment for femoroacetabular impingement: review of the literature. *Am J Sports Med*. 2014 Jun;42(6):1487-95. Epub 2013 Aug 30.
- Larson CM, Giveans MR, Taylor M. Does arthroscopic FAI correction improve function with radiographic arthritis? *Clin Orthop Relat Res*. 2011 Jun;469(6):1667-76. Epub 2010 Dec 22.

- 33.** Naal FD, Miozzari HH, Schär M, Hesper T, Nötzli HP. Midterm results of surgical hip dislocation for the treatment of femoroacetabular impingement. *Am J Sports Med.* 2012 Jul;40(7):1501-10. Epub 2012 May 3.
- 34.** Peters CL, Schabel K, Anderson L, Erickson J. Open treatment of femoroacetabular impingement is associated with clinical improvement and low complication rate at short-term followup. *Clin Orthop Relat Res.* 2010 Feb;468(2):504-10.
- 35.** Ribas M, Ledesma R, Cardenas C, Marín-Peña O, Toro J, Caceres E. Clinical results after anterior mini-open approach for femoroacetabular impingement in early degenerative stage. *Hip Int.* 2010;20(Suppl 7):S36-42. Epub 2010 May 27.
- 36.** Nho SJ, Beck EC, Nwachukwu BU, Cvetanovich GL, Neal WH, Harris JD, Weber AE, Mather RC. Survivorship and outcome of hip arthroscopy for femoroacetabular impingement syndrome performed with modern surgical techniques. *Am J Sports Med.* 2019 Jun;47(7):1662-9.
- 37.** Peters CL, Anderson LA, Diaz-Ledezma C, Anderson MB, Parvizi J. Does the nature of chondrolabral injury affect the results of open surgery for femoroacetabular impingement? *Clin Orthop Relat Res.* 2015 Apr;473(4):1342-8.
- 38.** Kunze KN, Beck EC, Nwachukwu BU, Ahn J, Nho SJ. Early hip arthroscopy for femoroacetabular impingement syndrome provides superior outcomes when compared with delaying surgical treatment beyond 6 months. *Am J Sports Med.* 2019 Jul;47(9):2038-44. Epub 2019 Jul 15.
- 39.** Vahedi H, Aalirezaie A, Rolo G, Parvizi J. Hip dysplasia compromises the outcome of femoroacetabular impingement surgery. *J Arthroplasty.* 2019 May;34(5):852-6. Epub 2019 Jan 18.