

6-1-2022

Beach Chair Versus Lateral Decubitus Surgical Positioning for Arthroscopic Anterior Shoulder Stabilization: A Retrospective Comparison of Clinical and Patient-Reported Outcomes

Ryan W Paul
Rothman Orthopaedic Institute

Sydney Streicher
Thomas Jefferson University

Alim Osman
Eastern Virginia Medical School

Chuka Ukekwe
Eastern Virginia Medical School

Follow this and additional works at: https://jdc.jefferson.edu/rothman_institute

Usman Zareef
Part of the [Orthopedics Commons](#), and the [Sports Medicine Commons](#)

[Let us know how access to this document benefits you](#)

See next page for additional authors

Recommended Citation

Paul, Ryan W; Streicher, Sydney; Osman, Alim; Ukekwe, Chuka; Zareef, Usman; Freedman, Kevin B; Erickson, Brandon J; Hammoud, Sommer; and Bishop, Meghan E, "Beach Chair Versus Lateral Decubitus Surgical Positioning for Arthroscopic Anterior Shoulder Stabilization: A Retrospective Comparison of Clinical and Patient-Reported Outcomes" (2022). *Rothman Institute Faculty Papers*. Paper 175.
https://jdc.jefferson.edu/rothman_institute/175

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Rothman Institute Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Ryan W Paul, Sydney Streicher, Alim Osman, Chuka Ukekwe, Usman Zareef, Kevin B Freedman, Brandon J Erickson, Sommer Hammoud, and Meghan E Bishop

Beach Chair Versus Lateral Decubitus Surgical Positioning for Arthroscopic Anterior Shoulder Stabilization

A Retrospective Comparison of Clinical and Patient-Reported Outcomes

Ryan W. Paul,* BS, Sydney Streicher,[†] BA, Alim Osman,[‡] MS, Chuka Ukekwe,[‡] MS, Usman Zareef,[§] BA, Kevin B. Freedman,* MD, Brandon J. Erickson,^{||} MD, Sommer Hammoud,* MD, and Meghan E. Bishop,^{||¶} MD

Investigation performed at Rothman Orthopaedic Institute, New York, New York, USA

Background: Surgical positioning can affect both perioperative and postoperative complication rates. It is unclear whether beach-chair versus lateral decubitus positioning affects outcomes in patients undergoing arthroscopic anterior shoulder stabilization surgery.

Purpose : The purpose of this study was to compare recurrent instability, complications, and patient-reported outcomes between patients who underwent arthroscopic anterior shoulder stabilization in the beach-chair versus the lateral decubitus positions. It was hypothesized that recurrent instability, complications, and patient-reported outcomes would not be affected by surgical positioning.

Study Design: Cohort study; Level of evidence, 3.

Methods: The authors reviewed the medical records of patients who underwent shoulder stabilization (*Current Procedural Terminology* codes 29806 and 29807) from 2015 to 2019. Patients were included only if anterior instability was confirmed, arthroscopic surgery was performed in response to shoulder instability, and a minimum of 2 years of follow-up data were available. Data collected for eligible patients included perioperative and postoperative complications, recurrent instability, reoperation, and revision. Patients also completed surveys for the American Shoulder and Elbow Surgeons (ASES) score, Single Assessment Numeric Evaluation (SANE) score, Oxford Shoulder Instability (OSI) score, and a return to any level of sport (RTS) questionnaire.

Results: Overall, 294 patients (162 lateral decubitus and 132 beach-chair positions) were included, with an average follow-up of 2.4 ± 1.6 years. There were no significant differences in demographics between groups, nor were there differences in the rates of postoperative dislocations, subjective instability, reoperations, revisions, or complications. There was a trend toward a higher revision rate in the beach-chair group (beach-chair, 6.1% vs lateral decubitus, 1.9%; $P = .069$). There was no significant difference between groups regarding RTS rates or postoperative ASES, SANE, and OSI scores at 3.3 ± 1.1 years postoperatively.

Conclusion: Surgical positioning for arthroscopic anterior shoulder stabilization did not significantly affect recurrent instability, complications, and patient-reported outcomes. Both beach-chair and lateral decubitus positioning provided good outcomes for anterior shoulder stabilization, with an overall recurrent dislocation rate of 7.8% at a mean of 3.3 years after surgery.

Keywords: shoulder instability; dislocation; stabilization; beach-chair; lateral decubitus; arthroscopy

The rate of shoulder dislocations is approximately 23 to 24 dislocations per 100,000 person-years, with anterior shoulder instability significantly more common than posterior shoulder instability.^{7,8,17} Patients who sustain an initial

anterior shoulder dislocation, as a general average, have a 39% chance of developing recurrent instability. The odds are 13 times greater for patients younger than 40 years and 3 times greater for men.¹³ Arthroscopic shoulder stabilization has been shown to effectively treat anterior shoulder instability, decreasing recurrence rates to between 10% and 17%.^{5,16} However, significant risk factors such as younger patient age, participation in competitive sports, off-

The Orthopaedic Journal of Sports Medicine, 10(6), 23259671221106474
DOI: 10.1177/23259671221106474
© The Author(s) 2022

This open-access article is published and distributed under the Creative Commons Attribution - NonCommercial - No Derivatives License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits the noncommercial use, distribution, and reproduction of the article in any medium, provided the original author and source are credited. You may not alter, transform, or build upon this article without the permission of the Author(s). For article reuse guidelines, please visit SAGE's website at <http://www.sagepub.com/journals-permissions>.

track Hill-Sachs lesions, and glenoid bone loss increase the chances of recurrent instability after surgery.¹⁶

Surgical positioning can affect perioperative and/or postoperative complication rates for many reasons, including proximity to nerves, issues with positioning, and surgical access.^{11,14} The beach-chair and lateral decubitus positions are the 2 most common positions for shoulder arthroscopy, and each offers its own unique advantages and disadvantages for surgeons.^{6,9,10,14,15} Beach-chair positioning allows for an easier transition to open surgery if necessary, has a quicker setup time, and also has a lower incidence rate of neuropathies, relative to lateral decubitus.^{3,6} However, lateral decubitus positioning has a similar surgery duration, easier visualization of the labrum, and greater circumferential access to the joint.^{3,6,10} A recent study found that patients who underwent Bankart repair arthroscopic anterior shoulder stabilization in the lateral decubitus position more frequently received ≥ 4 anchors and had more anchors placed in the inferior glenoid than patients in the beach-chair position. Despite these intraoperative differences, the type of surgical position is often selected according to each individual surgeon's preference, with no clear superiority in postoperative outcomes between positions.

Several review articles have been published comparing the beach-chair and lateral decubitus positions for shoulder arthroscopy,^{2,3,9,12,14,15} with 1 systematic review comparing these positions specifically for anterior shoulder stabilization surgery.³ However, the data from that review come from pooling of noncomparative retrospective studies. No large retrospective cohort study has been published to compare recurrent instability, complications, and patient-reported outcomes between patients who undergo arthroscopic anterior shoulder stabilization in beach-chair versus lateral decubitus positioning.

The primary purpose of this study was to compare recurrent instability, complications, and patient-reported outcomes between patients who underwent arthroscopic anterior shoulder stabilization in the beach-chair versus the lateral decubitus positions. We hypothesized that recurrent instability, complications, and patient-reported outcomes would be similar between the 2 positions.

METHODS

This study was exempt from institutional review board approval. We reviewed a list of patients aged 18 to 55 years who underwent arthroscopic shoulder stabilization (*Current Procedural Terminology codes 29806 and 29807*) from 2015 to 2019 at a single multicenter institution. Patients were included in the study if anterior instability was confirmed, arthroscopic surgery was performed in response to shoulder instability, and a minimum of 2 years of follow-up data were available. Patients were excluded for multidirectional instability, revision shoulder stabilization surgery, open surgery, significant glenohumeral osteoarthritis, concomitant rotator cuff repair, concomitant subacromial decompression, or isolated superior labral anterior-posterior (SLAP) repair. Patients with engaging Hill-Sachs lesions were indicated for concomitant remplissage and were also excluded from this study. Patients were excluded if surgery was not performed because of shoulder instability or if patient surgical positioning could not be confirmed.

Patient positioning was decided by the surgeon and was not randomized. Overall, 15 surgeons preferred to utilize the lateral decubitus position for all shoulder stabilization procedures, and 10 surgeons preferred the beach-chair position. All surgeons were fellowship trained in either shoulder and elbow or sport medicine. Routine venous thromboembolism prophylaxis was not provided to patients in this study cohort.

Data Collection

Patients were screened for eligibility while collecting pre- and perioperative data from the patient clinic notes and operative reports. Pre- and perioperative data collected included age, sex, body mass index, laterality of surgery, hand dominance, mechanism of injury, history of shoulder dislocation, concomitant procedures, number of anchors used, surgery duration (time from first incision to close), and perioperative complications.

Postoperative complications collected by screening postoperative patient encounters were subjective instability, shoulder dislocation, infection, nerve damage/injury, deep vein thrombosis/pulmonary embolism, significant shoulder

*Address correspondence to Meghan E. Bishop, MD, Rothman Orthopaedic Institute, 645 Madison Avenue, 3rd and 4th Floors, New York, NY 10022, USA (email: meghan.bishop@rothmanortho.com) (Twitter: @MegBishopMD).

*Rothman Orthopaedic Institute, Philadelphia, Pennsylvania, USA.

†Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, Pennsylvania, USA.

‡Eastern Virginia Medical School, Norfolk, Virginia, USA.

§Rutgers Robert Wood Johnson Medical School, New Brunswick, New Jersey, USA.

||Rothman Orthopaedic Institute, New York, New York, USA.

Presented at the AOSSM Annual Meeting, Colorado Springs, Colorado, July 2022.

Final revision submitted February 7, 2022; accepted February 25, 2022.

One or more of the authors has declared the following potential conflict of interest or source of funding: K.B.F. has received education payments from Liberty Surgical, consulting fees from Medical Device Business Services, and speaking fees and honoraria from Vericel. B.J.E. has received research support from Arthrex, DePuy, Linvatec, Smith & Nephew, and Stryker; education payments from Arthrex, DePuy, and Smith & Nephew; and consulting fees from Arthrex. S.H. has received education payments from Liberty Surgical and hospitality payments from Arthrex and Smith & Nephew and has stock/stock options in Johnson & Johnson. M.E.B. has received grant payments from Arthrex, education payments from Arthrex and Smith & Nephew, and hospitality payments from Stryker. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.

Ethical approval for this study was waived by Thomas Jefferson University (study No. 21E.514).

TABLE 1
Comparison of Preoperative Data Between Patients Who Underwent Shoulder Stabilization Surgery
in the Beach-Chair Versus Lateral Decubitus Position^a

Preoperative Variable	Lateral Decubitus, n = 162	Beach-Chair, n = 132	P
Age, y	29.2 ± 9.4	27.2 ± 9.3	.072
Male sex	124 (76.5)	103 (78.0)	.871
BMI	26.0 ± 4.8	26.3 ± 5.0	.542
Surgery on dominant side	60 (37.0)	55 (41.7)	.836
Mechanism of injury			.511
Direct impact	87 (53.7)	70 (53.0)	
Reaching	20 (12.3)	11 (8.3)	
Lifting	11 (6.8)	7 (5.3)	
Other	44 (27.2)	44 (33.3)	
Acute injury	91 (56.2)	72 (54.5)	≥.999
Preoperative ASES score	65.4 ± 23.0	58.0 ± 24.8	.162
Preoperative SANE score	51.5 ± 24.4	49.1 ± 26.9	.657

^aCategorical data are presented as No. (%), and continuous data are presented as mean ± SD. ASES, American Shoulder and Elbow Surgeons; BMI, body mass index; SANE, Single Assessment Numeric Evaluation.

pain and swelling, and reoperation/revision. Shoulder pain or swelling was considered a postoperative complication only if a subsequent arthroscopy or manipulation under anesthesia was performed. Revision was considered any shoulder stabilization reoperation, such as revision Bankart repair, capsulorrhaphy, or Latarjet procedure. Subjective instability was considered patient-reported apprehension, a subluxation event, or a dislocation event.

Patients were also contacted via REDCap (Vanderbilt University) to complete patient-reported outcome surveys, including postoperative complications and reoperations, the American Shoulder and Elbow Surgeons (ASES) score, Single Assessment Numeric Evaluation (SANE) score, Oxford Shoulder Instability (OSI) score, and a return to any level of sport (RTS) questionnaire. ASES and SANE scores range from 0 (poor) to 100 (healthy), while OSI scores range from 0 (poor) to 48 (healthy). No minimal clinically important differences are available for these scores regarding patients with anterior shoulder stabilization at a minimum 2-year follow-up. A general survey asking about postoperative complications and future ipsilateral shoulder surgery was also included to confirm the findings from postoperative encounter screening. Only patients who responded “yes” to the question, “Were you participating in a sport prior to your shoulder injury?” were included in the RTS analysis. These patients were subsequently asked, “Were you able to return to sport after surgery?” Possible answers were “yes, same/higher level”; “yes, lower level”; “no, not interested”; “no, could not return due to shoulder pain/instability”; “no, the opportunity was no longer available”; and “no, but returned to play a different sport.”

Statistical Analysis

Demographics, perioperative data, and postoperative outcomes were compared between patients who underwent surgery in beach-chair versus lateral decubitus positioning. Similar analyses were also performed comparing male versus female patients and younger (<25 years) versus older

(≥25 years) patients. The age of 25 years was selected as a cutoff because age younger than 25 years is commonly clinically used as a risk factor for failed nonoperative treatment of shoulder instability, and this age cutoff provided sufficient sample sizes for both age groups in the current study. Mann-Whitney tests were used to calculate *P* values for nonparametric data. The chi-square or Fisher exact tests were used to calculate *P* values for categorical data. *P* values less than .05 were deemed significant. A multivariate regression analysis was performed with surgical position (beach-chair vs lateral decubitus), age, sex, number of anchors placed, and history of shoulder dislocations (acute vs chronic) as independent variables and recurrent instability as the dependent variable. All statistical analyses were done using RStudio (Version 3.6.3).

RESULTS

There were 294 patients included in this study, 162 patients who underwent surgery in the lateral decubitus position and 132 in beach-chair position, with an average follow-up of 2.4 ± 1.6 years. Patients were on average 28.3 ± 9.4 years of age, and there were 227 (77.2%) men. Of the 294 patients, 215 (73%) had more than 2 years of clinical follow-up, and 175 (60%) patients completed surveys for the RTS questionnaire and ASES, SANE, and OSI scores at least 2 years postoperatively. There were no significant demographic differences between groups, and 2-year follow-up rates also did not differ (lateral decubitus, 72% vs beach-chair, 75%; *P* = .514) (Table 1).

Beach-chair positioning was associated with an increased number of anchors used (4.0 vs 3.3 for lateral decubitus; *P* = .001) (Table 2). All other perioperative variables, such as concomitant SLAP repair, surgery duration, and perioperative complications, were similar between groups. The only perioperative complication occurred in a 41-year-old man who underwent surgery in the lateral decubitus position. He experienced venous oozing during the removal of arthroscopic equipment from the anterior-

TABLE 2
Comparison of Perioperative Data Between Patients Who Underwent Shoulder Stabilization Surgery
in the Beach-Chair Versus Lateral Decubitus Position^a

Perioperative Variable	Lateral Decubitus, n = 162	Beach-Chair, n = 132	P
Concomitant SLAP repair	13 (8.0)	12 (9.1)	.744
Concomitant surgery	26 (16.0)	31 (23.5)	.109
No. of anchors placed	3.3 ± 1.1	4.0 ± 1.7	.001
Surgery duration, min	69.4 ± 21.4	66.8 ± 21.3	.637

^aCategorical data are presented as No. (%), and continuous data are presented as mean ± SD. Boldface P value indicates a statistically significant difference between groups ($P < .05$). SLAP, superior labral anterior-posterior.

TABLE 3
Comparison of Postoperative Outcomes Between Patients Who Underwent Shoulder Stabilization Surgery
in the Beach-Chair Versus Lateral Decubitus Position^a

Postoperative Outcome	Lateral Decubitus, n = 162	Beach-Chair, n = 132	P
Redislocation	12 (7.4)	11 (8.3)	.940
Subjective instability	25 (15.4)	23 (17.4)	.763
Reoperation	5 (3.1)	9 (6.8)	.223
Revision	3 (1.9)	8 (6.1)	.069
Postoperative neuropathy	7 (4.3)	5 (3.8)	.818
All complications	32 (19.8)	33 (25.0)	.349
Neuropathy	7 (4.3)	5 (3.8)	≥.999
Patient-Reported Outcome	Lateral Decubitus, n = 97	Beach-Chair, n = 78	P
Ability to RTS	47 (69.1); n = 68	34 (63.0); n = 54	.602
RTS for contact athletes	26 (66.7); n = 39	21 (65.6); n = 32	.926
RTS for overhead athletes	18 (72.0); n = 25	8 (53.3); n = 15	.231
ASES score	87.5 ± 17.1	88.3 ± 13.9	.731
SANE score	77.1 ± 23.5	77.3 ± 20.2	.959
OSI score	41.1 ± 9.3	42.1 ± 6.7	.400

^aCategorical data are presented as No. (%), and continuous data are presented as mean ± SD. ASES, American Shoulder and Elbow Surgeons; OSI, Oxford Shoulder Instability; RTS, return to any level of sport; SANE, Single Assessment Numeric Evaluation.

inferior portal, and hemostasis occurred with electrocautery and placement of sutures.

Postoperatively, no patients in either study cohort developed an infection or a venous thromboembolism. Beach-chair positioning had an increased revision rate but this trend was not significant (6.1% beach-chair vs 1.9% lateral decubitus position; $P = .069$) (Table 3). Rates of postoperative subjective instability, reoperations, revisions, neuropathy, and all complications also did not differ between groups (all $P > .05$). Overall, 23 (7.8%) included patients experienced a redislocation. In the lateral decubitus cohort, 3 of 12 patients who experienced a redislocation had experienced only 1 acute redislocation, while 5 patients had experienced several (2-5) redislocations and 4 patients had experienced frequent (≥ 6) redislocations. In the beach-chair cohort, 5 of 11 patients who experienced a redislocation had experienced only 1 acute redislocation, while 4 patients had experienced several (2-5) redislocations and 2 patients had experienced frequent (≥ 6) redislocations.

Revision shoulder stabilization procedures in the lateral decubitus cohort included 1 Latarjet procedure and 2 revision Bankart repairs, while revision procedures in the

beach-chair cohort included 4 Latarjet procedures and 4 revision Bankart repairs. Further details about each patient who required a revision shoulder stabilization procedure are available in Appendix Table A1.

There were no significant differences between groups regarding rates of returning to any level of sport as well as postoperative ASES, SANE, and OSI scores at an average of 3.3 ± 1.1 years of follow-up (all $P > .05$). RTS rates for contact athletes and overhead athletes also did not significantly differ between lateral decubitus and beach-chair positions.

Comparing 227 male and 67 female patients, all peri- and postoperative outcomes were similar except that men had an increased number of anchors placed (3.8 in men vs 3.3 in women; $P = .039$) (Appendix Tables A2 and A3). When comparing patients aged <25 versus ≥ 25 years, the younger patients had significantly shorter surgery durations (62.5 vs 72.4 minutes, respectively; $P = .048$) and higher rates of postoperative redislocations (12.8% vs 3.3%, respectively; $P = .005$), while all other peri- and postoperative outcomes were similar between age groups (Appendix Tables A4 and A5).

TABLE 4
Multivariate Regression Evaluating Independent Risk Factors for Subjective Shoulder Instability
After Arthroscopic Anterior Shoulder Stabilization

Independent Variable	Subjective Instability, Odds Ratio (95% CI)	P
Surgical position: beach-chair	1.22 (0.60-2.47)	.588
Age	1.00 (0.96-1.04)	.888
Female sex	1.44 (0.63-3.18)	.371
No. of anchors placed	0.96 (0.73-1.22)	.771
History of recurrent dislocations before surgery	0.61 (0.27-1.27)	.199

Multivariate regression analysis of surgical position, age, sex, number of anchors placed, and shoulder dislocation history showed that none of these variables were independently related to recurrent instability in the 48 included patients who developed recurrent anterior shoulder instability (Table 4).

DISCUSSION

Our hypothesis was confirmed, as recurrent instability rates, complication rates, and patient-reported outcomes were similar between groups. Patients who underwent anterior shoulder stabilization in the beach-chair position had a trend toward increased revision rates compared with patients in the lateral decubitus position, but this difference did not reach statistical significance.

A large systematic review pooled the recurrence rates of 3668 patients with arthroscopic anterior shoulder instability from 64 studies.³ They found that the lateral decubitus position provided a significantly lower recurrent instability rate for patients with anterior shoulder instability than the beach-chair position (8.5% vs 14.6%, respectively; $P = .004$).³ However, no comparative studies were included in that systematic review, which significantly limited the strength of the conclusions drawn. In comparison, the present comparative study found no difference between beach-chair and lateral decubitus positions for redislocation and subjective instability rate. The previous systematic review may have found higher recurrence rates than the redislocation rates observed in the current study because the severity of recurrent instability could not be evaluated in the systematic review, so it is unclear whether the differences in recurrent instability observed were because of apprehension, subluxation, or Frank dislocation.³ Whenever possible, subjective instability and redislocation should be evaluated in isolation, as recurrent instability as a whole may include a spectrum of instability from apprehension to dislocation.^{3,4} The present comparative study provided an evaluation of redislocation rates separate from subjective instability to clarify the severity of postoperative instability. Evaluating redislocations and subjective instability separately showed that younger patients were more likely to redislocate their shoulder after surgery. However, the frequency of subjective instability did not differ between younger and older

patients, and age was not an independent risk factor for subjective instability.

Lateral decubitus positioning is believed to allow better access to the glenohumeral joint, easier visualization of the labrum, and greater circumferential access to the joint.^{3,6,10} This is supported by the findings of a recent cross-sectional study comparing the number and location of anchors used between beach-chair and lateral decubitus positions for arthroscopic anterior shoulder stabilization.¹ The authors found that anchors were placed in the 6-o'clock portion of the glenoid more than 2 times as frequently when patients were in the lateral decubitus position, and a larger number of anchors were placed with lateral decubitus positioning as well.¹ Despite the differences in anchor utilization, the current study found no differences in redislocation or subjective instability rates between beach-chair and lateral decubitus positions. However, beach-chair positioning was associated with a greater number of anchors placed than lateral decubitus in this cohort. Although our data set on number of anchors used is much smaller than the data presented by Baron et al,¹ the drastic difference in results shows that there may be significant surgeon bias when evaluating perioperative factors such as number and location of anchors. A randomized study design would be optimal to clarify perioperative differences between beach-chair and lateral decubitus positioning; however, surgeon preferences and total surgical volume within each position significantly limit the feasibility and validity of such a clinical trial.

This is the first comparative study to our knowledge to evaluate patient-reported outcomes between patients who underwent arthroscopic anterior shoulder stabilization in the beach-chair versus lateral decubitus position. The current study found no significant differences in ASES, SANE, or OSI scores between beach-chair and lateral decubitus positions. Similarly, Frank et al³ performed a systematic review evaluating Rowe and Constant-Murley scores across 39 and 22 studies, respectively, and found that patient positioning did not affect these scores in patients with anterior shoulder stabilization. Overall, it appears that surgeons do not need to alter their preferred patient positioning for anterior shoulder stabilization, as patients can expect similar clinical and patient-reported outcomes whether undergoing surgery in the beach-chair or lateral decubitus position.

Limitations

There are several limitations to this study. First, this study was not randomized, and thus surgeon bias may have affected perioperative variables and postoperative outcomes. Second, the prevalence of significant Hill-Sachs and bony Bankart lesions, as well as the amount of glenoid or humeral head bone loss, was not evaluated. However, significant glenoid bone loss is often considered a contraindication for arthroscopic treatment of anterior shoulder instability,⁴ and patients with 15% to 20% bone loss were excluded from arthroscopic stabilization surgery in our cohort. Third, the RTS and patient-reported outcome analyses were limited by a 60% patient follow-up rate; however, a large sample size of responses was still observed. Fourth, we provided various answer options evaluating why athletes did not RTS, which may have lowered our RTS rates relative to similar studies since patients may not have returned based on choice rather than because of their shoulder. Fifth, the postoperative outcomes in this study may have been affected by 79 (26.9%) patients being lost to 2-year follow-up and only 175 (60%) patients completing patient-reported outcome measures. Sixth, this study may have been underpowered to evaluate differences in recurrent instability. A post hoc power analysis was performed, finding that a sample of 28,480 patients (14,240 per group) would be required to detect a significant difference in redislocation rates, and a sample of 724 patients (362 per group) would be needed to observe a significant difference in revision rate. However, the limited number of patients with revisions (n = 11) may limit the validity of the post hoc power analysis findings. Seventh, we did not have data on preoperative sport participation for all included patients because of the nature of retrospective medical-record review, and thus did not include information or analyses on sport participation. Last, the location of anchor placement was not evaluated because of infrequent reporting and the lack of validity in self-reported anchor placement location from the operative report. The validity of anchor data in the operative reports may also have affected our assessment of the number of anchors used, which could explain why our findings are contrary to prior studies.² A prospective study design that confirms data collection on anchor placement would strengthen the present analysis.

CONCLUSION

Surgical positioning for arthroscopic anterior shoulder stabilization did not significantly affect recurrent instability, complications, and patient-reported outcomes. Both beach-chair and lateral decubitus positioning provided good outcomes for anterior shoulder stabilization, with an overall recurrent dislocation rate of 7.8% at a mean of 3.3 years after surgery.

REFERENCES

1. Baron JE, Duchman KR, Hettrich CM, et al. Beach chair versus lateral decubitus position: differences in suture anchor position and number

- during arthroscopic anterior shoulder stabilization. *Am J Sports Med.* 2021;49(8):2020-2026. doi:10.1177/03635465211013709
2. de Sa D, Sheean AJ, Morales-Restrepo A, Dombrowski M, Kay J, Vyas D. Patient positioning in arthroscopic management of posterior-inferior shoulder instability: a systematic review comparing beach chair and lateral decubitus approaches. *Arthroscopy.* 2019; 35(1):214-224.e3. doi:10.1016/j.arthro.2018.06.057
3. Frank RM, Saccomanno MF, McDonald LS, Moric M, Romeo AA, Provencher MT. Outcomes of arthroscopic anterior shoulder instability in the beach chair versus lateral decubitus position: a systematic review and meta-regression analysis. *Arthroscopy.* 2014;30(10): 1349-1365. doi:10.1016/j.arthro.2014.05.008
4. Harris JD, Gupta AK, Mall NA, et al. Long-term outcomes after Bankart shoulder stabilization. *Arthroscopy.* 2013;29(5):920-933. doi:10.1016/j.arthro.2012.11.010
5. Hurley ET, Manjunath AK, Bloom DA, et al. Arthroscopic Bankart repair versus conservative management for first-time traumatic anterior shoulder instability: a systematic review and meta-analysis. *Arthroscopy.* 2020;36(9):2526-2532. doi:10.1016/j.arthro.2020.04.046
6. Jinnah AH, Mannava S, Plate JF, Stone AV, Freehill MT. Basic shoulder arthroscopy: lateral decubitus patient positioning. *Arthrosc Tech.* 2016;5(5):e1069-e1075. doi:10.1016/j.eats.2016.05.010
7. Kraeutler MJ, McCarty EC, Belk JW, et al. Descriptive epidemiology of the MOON shoulder instability cohort. *Am J Sports Med.* 2018;46(5): 1064-1069. doi:10.1177/0363546518755752
8. Leroux T, Wasserstein D, Veillette C, et al. Epidemiology of primary anterior shoulder dislocation requiring closed reduction in Ontario, Canada. *Am J Sports Med.* 2014;42(2):442-450. doi:10.1177/0363546513510391
9. Li X, Eichinger JK, Hartshorn T, Zhou H, Matzkin EG, Warner JP. A comparison of the lateral decubitus and beach-chair positions for shoulder surgery: advantages and complications. *J Am Acad Orthop Surg.* 2015;23(1):18-28. doi:10.5435/JAAOS-23-01-18
10. Mannava S, Jinnah AH, Plate JF, Stone AV, Tuohy CJ, Freehill MT. Basic shoulder arthroscopy: beach chair patient positioning. *Arthrosc Tech.* 2016;5(4):e731-e735. doi:10.1016/j.eats.2016.02.038
11. Matjasko MJ, Hellman J, Mackenzie CF. Venous air embolism, hypotension, and end-tidal nitrogen. *Neurosurgery.* 1987;21(3):378-382. doi:10.1227/00006123-198709000-00017
12. Moeller EA, Houck DA, McCarty EC, et al. Outcomes of arthroscopic posterior shoulder stabilization in the beach-chair versus lateral decubitus position: a systematic review. *Orthop J Sports Med.* 2019;7(1): 2325967118822452. doi:10.1177/2325967118822452
13. Olds M, Ellis R, Donaldson K, Parmar P, Kersten P. Risk factors which predispose first-time traumatic anterior shoulder dislocations to recurrent instability in adults: a systematic review and meta-analysis. *Br J Sports Med.* 2015;49(14):913-922. doi:10.1136/bjsports-2014-094342
14. Peruto CM, Ciccotti MG, Cohen SB. Shoulder arthroscopy positioning: lateral decubitus versus beach chair. *Arthroscopy.* 2009;25(8): 891-896. doi:10.1016/j.arthro.2008.10.003
15. Rojas J, Familiari F, Bitzer A, Srikumaran U, Papalia R, McFarland EG. Patient positioning in shoulder arthroscopy: which is best? *Joints.* 2019;7(2):46-55. doi:10.1055/s-0039-1697606
16. Verweij LPE, van Spanning SH, Grillo A, et al. Age, participation in competitive sports, bony lesions, ALPSA lesions, > 1 preoperative dislocations, surgical delay and ISIS score > 3 are risk factors for recurrence following arthroscopic Bankart repair: a systematic review and meta-analysis of 4584 shoulders. *Knee Surg Sports Traumatol Arthrosc.* 2021;29(12):4004-4014. doi:10.1007/s00167-021-06704-7
17. Zacchilli MA, Owens BD. Epidemiology of shoulder dislocations presenting to emergency departments in the United States. *J Bone Joint Surg Am.* 2010;92(3):542-549. doi:10.2106/JBJS.I.00450

APPENDIX

APPENDIX TABLE A1
 Descriptions of the Patients in the Lateral Decubitus and Beach-Chair Cohorts
 Who Required Revision Shoulder Stabilization^a

Patient	Injury History	Revision Procedure	Reason for Revision
Lateral Decubitus, n = 3			
29-year-old man	1 acute dislocation while lifting heavy can, with arm in abduction and external rotation	Latarjet procedure 18 mo postop	Shoulder reinjury 7 mo postop, was on a bus ride holding onto a post as the bus did a sudden stop; experienced 2 further dislocations at work
34-year-old man	1 acute dislocation while putting up drywall in home, arm went too far back with weight overhead	Bankart repair with subacromial decompression 5 mo postop	Slight retear of the repaired labrum; prolonged pain and discomfort throughout recovery, with feeling of instability while arm at his side
32-year-old woman, recreational softball athlete	Subjective instability and shoulder pain, no acute dislocation; symptoms began after taking heavy bag of trash outside	Bankart repair 2 y postop	Developed shoulder pain, stiffness, and apprehension several months after initial surgery after carrying a heavy bag of ice, which required lysis of adhesions 6 mo postop; despite this, was able to return to recreational softball 10 mo after initial Bankart repair; however, 25 mo after the initial surgery, slipped while descending stairs and fell on shoulder, resulting in a labral retear
Beach-Chair, n = 8			
45-year-old man, recreational golf athlete	Chronic shoulder instability for 20 y. He was being successfully treated nonoperatively until experiencing 2 recent dislocations, most recently while playing paddle ball.	Latarjet procedure 3.3 y postop	Returned to golf 4 mo after initial Bankart repair; slipped on rocks 21 mo postop, experienced an anterior shoulder dislocation while trying to catch himself; attempted nonoperative treatment but experienced several subsequent dislocations
24-year-old man	Chronic shoulder instability; first visited clinic with shoulder dislocation after reaching for car door handle and most recently visited clinic after rolling onto side while sleeping and experiencing another shoulder dislocation; has had about 20 shoulder dislocations in the past year; no history of connective tissue disorder	Latarjet procedure 9 mo postop	Fell on shoulder while playing soccer 7 mo postop, dislocated shoulder and had it reduced in the emergency department; attempted nonoperative treatment, but pain and instability persisted and significant bone loss was found on CT imaging
18-year-old man, hockey athlete	1 acute dislocation after being hit from behind in hockey game, fell forward and dislocated shoulder while trying to catch himself	Latarjet procedure 2.5 y postop	14 mo postop, fell onto outstretched arm while ice skating and experienced shoulder dislocation; then 18 mo postop, experienced another dislocation after falling off scooter onto the same shoulder; surgery was planned for 19 mo postop, but COVID-19 pandemic resulted in a cancellation; several continued dislocations resulted in a revision in March 2021
18-year-old man, rock climber and ultimate frisbee athlete	2 acute dislocations, 1 while indoor skydiving and most recently with arm outstretched overhead while rock climbing	Bankart repair 16 mo postop	Returned to ultimate frisbee 4 mo postop but was not interested in returning to rock climbing; 14 mo postop, experienced a painful shoulder subluxation while overhead throwing a basketball and put shoulder back in place himself; had a labral retear and increased size of Hill-Sachs lesion

(continued)

Appendix Table A1 (continued)

Patient	Injury History	Revision Procedure	Reason for Revision
23-year-old man	1 acute dislocation while falling onto outstretched arm during a football game	Bankart repair 17 mo postop	After lifting heavy at the gym 3 mo postop, developed intermittent shoulder pain; he felt no pain while at rest but felt pain during physical activity as light as walking
48-year-old man	1 acute dislocation after falling on arm while descending stairs	Bankart repair 3.5 y postop	Developed recurrent instability while playing football 39 mo postop; nonoperative treatment was unsuccessful, and he continued to have difficulty lifting arm with significant nighttime pain
28-year-old woman	6 acute dislocations, the first occurring after someone jumped on the patient's back while dancing; she experienced 5 subsequent dislocations, all requiring reduction in the emergency department	Latarjet procedure 2.5 y postop	Fell onto her outstretched arm 2 y postop and experienced several subsequent dislocations that she was able to self-reduce; shoulder pain was not significant, but significant bone loss was noted on CT imaging, suggesting that recurrence would be expected if not surgically treated
49-year-old woman	1 acute dislocation, slipped and fell at work while pushing a wheelchair; also injured her back during this injury, which required surgery before shoulder surgery	Bankart repair 6 mo postop	Patient still had significant shoulder pain after 2-3 mo postop, with significant range of motion limitations; physical therapy was not helping to improve symptoms

^aNone of the 11 patients who required a revision shoulder stabilization procedure underwent concomitant superior labrum anterior-posterior repair. CT, computed tomography; postop, postoperatively.

APPENDIX TABLE A2
Comparison of Perioperative Variables Between Male and Female Patients^a

Perioperative Variable	Male, n = 227	Female, n = 67	P
Concomitant SLAP repair	22 (9.7)	3 (4.5)	.179
Concomitant surgery	44 (19.4)	13 (19.4)	.997
No. of anchors placed	3.8 ± 1.5	3.3 ± 1.4	.039
Surgery duration, min	68.7 ± 21.8	64.1 ± 19.7	.409

^aCategorical data are presented as No. (%), and continuous data are presented as mean ± SD. SLAP, superior labral anterior-posterior.

APPENDIX TABLE A3
Comparison of Postoperative Outcomes Between Male and Female Patients^a

Postoperative Outcome	Male, n = 227	Female, n = 67	P
Redislocation	17 (7.5)	6 (9.0)	.894
Subjective instability	36 (15.9)	12 (17.9)	.833
Reoperation	10 (4.4)	4 (6.0)	.531
Revision	8 (3.5)	3 (4.5)	.718
Postoperative neuropathy	8 (3.5)	4 (6.0)	.374
All complications	48 (21.1)	17 (25.4)	.572
Patient-Reported Outcome	Male, n = 130	Female, n = 45	P
Ability to RTS	59 (63.4); n = 93	22 (75.9); n = 29	.312
ASES score	87.9 ± 16.2	87.8 ± 14.3	.991
SANE score	77.4 ± 22.9	76.5 ± 19.3	.802
OSI score	41.5 ± 8.7	41.5 ± 6.9	.991

^aCategorical data are presented as No. (%), and continuous data are presented as mean ± SD. ASES, American Shoulder and Elbow Surgeons; OSI, Oxford Shoulder Instability; RTS, return to any level of sport; SANE, Single Assessment Numeric Evaluation.

APPENDIX TABLE A4
Comparison of Perioperative Variables Between Patients <25 and ≥25 Years of Age^a

Perioperative Variable	Age <25 y, n = 141	Age ≥25 y, n = 153	P
Concomitant SLAP repair	11 (7.8)	14 (9.2)	.679
Concomitant surgery	25 (17.7)	32 (20.9)	.490
No. of anchors placed	3.8 ± 1.6	3.5 ± 1.4	.081
Surgery duration, min	62.5 ± 15.1	72.4 ± 25.1	.048

^aCategorical data are presented as No. (%), and continuous data are presented as mean ± SD. Boldface P value indicates a statistically significant difference between groups ($P < .05$). SLAP, superior labral anterior-posterior.

APPENDIX TABLE A5
Comparison of Postoperative Outcomes Between Patients <25 and ≥25 Years of Age^a

Postoperative Outcome	Age <25 y, n = 141	Age ≥25 y, n = 153	P
Redislocation	18 (12.8)	5 (3.3)	.005
Subjective instability	25 (17.7)	23 (15.0)	.640
Reoperation	5 (3.5)	9 (5.9)	.506
Revision	4 (2.8)	7 (4.6)	.633
Postoperative neuropathy	6 (4.3)	6 (3.9)	.885
All complications	33 (23.4)	32 (20.9)	.709
Patient-Reported Outcome	Age <25 y, n = 79	Age ≥25 y, n = 96	P
Ability to RTS	38 (58.5); n = 65	43 (75.4); n = 57	.074
ASES score	90.2 ± 12.2	86.1 ± 17.8	.085
SANE score	78.7 ± 21.0	76.0 ± 22.8	.422
OSI score	42.7 ± 6.3	40.6 ± 9.4	.092

^aCategorical data are presented as No. (%), and continuous data are presented as mean ± SD. Boldface P value indicates a statistically significant difference between groups ($P < .05$). ASES, American Shoulder and Elbow Surgeons; OSI, Oxford Shoulder Instability; RTS, return to any level of sport; SANE, Single Assessment Numeric Evaluation.