11-10-2013

Two-vessel off-pump coronary artery bypass grafting by left thoracotomy in a complex redo case

Shinya Unai, MD
Thomas Jefferson University

Hitoshi Hirose, MD, PhD
Thomas Jefferson University

Gary A. Cook, PA-C
Thomas Jefferson University

Nicholas C. Cavarocchi, MD
Thomas Jefferson University

John W..C. Entwistle III, MD, PhD
Thomas Jefferson University

Follow this and additional works at: <https://jdc.jefferson.edu/surgeryfp>

Part of the [Surgery Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Unai S, Hirose H, Cavarocchi N, Entwistle J. Two-vessel off-pump coronary artery bypass grafting by left thoracotomy in a complex redo case. Presented at Chest 2013. Chicago IL. October 26-31, 2013.

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Surgery Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Two-vessel off-pump coronary artery bypass grafting by left thoracotomy in a complex reoperative case

Shinya Unai, MD, Hitoshi Hirose, MD, PhD, Gary A. Cook MS, PA-C, Nicholas C. Cavarocchi, MD, John W. C. Entwistle III, MD, PhD.
From Thomas Jefferson University Hospital, Philadelphia, PA, USA.

Introduction

The left thoracotomy approach is an alternative technique for coronary revascularization to avoid complications associated with re-sternotomy, such as injury to patent grafts, right ventricle, aorta, during re-sternotomy.

Situations such as calcification of the ascending aorta and previous mediastinitis favor the use of the left thoracotomy approach.

Revascularization of the circumflex territory via a lateral thoracotomy has been reported previously. However, reports of revascularization of the LAD combined with circumflex artery territory via left thoracotomy approach are rare.

We successfully performed an off-pump CABG by left thoracotomy in a complex redo case to revascularize the LAD and obtuse marginal branch (OM) in an 83 year-old-man who had CABG and AVR in the past with a heavily calcified ascending aorta.

Case Report

- 83-year-old caucasian male
- History of CABG 20 years ago, redo CABG in the following year due to graft failure, and an AVR with a mechanical valve 12 years ago, and multiple coronary interventions.
- Presented with unstable angina
- Echo showed normal LV function
- Due to his history of early stent re-stenosis and location of the disease, he was considered not suitable for repeat PCI.
- CT scan showed severe calcification of the ascending aorta (left), and mild disease on the descending aorta.

Pre-operative catheterization

- 70% stenosis of the distal left main
- 80 % stenosis of the mid LAD stent
- 75% stenosis of the ostium of the LCx
- SVG to RCA was diseased but patent
- SVG to the LAD was totally occluded

CT scan

Intra-operative technique

- Double lumen endotracheal intubation
Supine position → right lateral decubitus position with the pelvis corkscrewed
- Left postero-lateral thoracotomy through 5th intercostal space
- The pericardium was opened longitudinally anterior to the phrenic nerve
- Heartstring III and Acrobat-i (Maquet Inc., Wayne, NJ, USA) were used for the anastomosis

proximal anastomosis to descending aorta

SVG to OM

Post-operative catheterization

Postoperative catheterization demonstrates patent grafts.
(Arrowheads: SVG to OM, arrow: radial artery to LAD.)

Hospital course

The overall cardiac function was stable and he was transferred to a rehabilitation facility on postoperative day 17.

Conclusion

Redo CABG is a technical challenge, but we were able to successfully undergo a 4th time redo CABG via the left thoracotomy approach and proximal anastomosis of the conduits in the descending aorta.

This technique in selected patients could be a useful that may reduce the complications related to redo sternotomy.

Contact information

John W. C. Entwistle III, MD, PhD.
john.entwistle@jefferson.edu

Associate Professor of Surgery
Division of Cardiothoracic Surgery
Thomas Jefferson University