

12-12-2022

Total Hip Arthroplasty in Patients With Neurological Conditions: A Systematic Review

Conor S O'Driscoll

Andrew J Hughes

Martin S Davey

Joseph M Queally

Brendan J O'Daly


Follow this and additional works at: https://jdc.jefferson.edu/rothman_institute



Part of the [Orthopedics Commons](#)

[Let us know how access to this document benefits you](#)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Rothman Institute Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.



Systematic Review

Total Hip Arthroplasty in Patients With Neurological Conditions: A Systematic Review

Conor S. O'Driscoll, MB, BAO, BCh, MRCS^{1,2,3,*},
 Andrew J. Hughes, MB, BAO, BCh, MCh, FRCS^{1,3,4},
 Martin S. Davey, MB, BAO, BCh, MCh, MRCS^{2,3},
 Joseph M. Queally, MB, BAO, BCh, MD, MEd, FRCS^{1,5,6},
 Brendan J. O'Daly, MB, BAO, BCh, MSc, FRCS^{1,3,6}

¹ Department of Trauma & Orthopaedics, Tallaght University Hospital, Dublin, Ireland

² Department of Trauma & Orthopaedics, Galway University Hospital, Galway, Ireland

³ Department of Trauma & Orthopaedics, Royal College of Surgeons Ireland, Dublin, Ireland

⁴ Rothman Orthopedic Institute, Thomas Jefferson University Hospital, PA, USA

⁵ Department of Trauma & Orthopaedics, St James Hospital, Dublin, Ireland

⁶ School of Medicine, University College Dublin, Dublin, Ireland

ARTICLE INFO

Article history:

Received 20 June 2022

Received in revised form

20 October 2022

Accepted 1 November 2022

Available online xxx

Keywords:

Total hip arthroplasty

Total hip replacement

Parkinson's disease

Cerebral palsy

Multiple sclerosis

Poliomyelitis

ABSTRACT

Background: As operative techniques and implant design have evolved over time, total hip arthroplasty (THA) is increasingly being carried out for patients with neurological impairment. This patient group places unique surgical challenges to the arthroplasty surgeon, which may include contractures, instability, and altered muscular tone. The purpose of this systematic review is to report the patient outcomes, complications, and implant survival following THA for patients with neurological conditions affecting the hip. Thus, we aim to support orthopaedic surgeon decision-making when considering and planning THA for these patients.

Methods: A systematic review was performed as per Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines using the PubMed/Medline OVID, Cochrane, and Embase databases. All studies reporting the outcomes of THA in the neurological population which met defined inclusion criteria were included.

Results: From an initial screen of 1820 studies, 45 studies with a total of 36,251 THAs were included in the final selection. All 45 studies reported complication rates, with controls included in 16 for comparison. High complication rates were observed following THA in the neurologically impaired population, most notably dislocation with observed rates up to 10.6%. An improvement was noted in all 36 studies (1811 THAs) which reported upon patient-reported outcomes.

Conclusions: THA may be beneficial in the selected patients with neurological conditions, to reduce pain and improve function. There is an increased risk of complications which require careful consideration when planning the operation and open discussion with prospective patients and caregivers before proceeding with surgery.

© 2022 The Authors. Published by Elsevier Inc. on behalf of The American Association of Hip and Knee Surgeons. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Search Strategies and Materials Available Online. Prospero Registration: CRD42022332353; <https://www.crd.york.ac.uk/PROSPERO/>.

All authors contributed to the study conception and design. Material preparation, data collection, and analysis were performed by Conor O'Driscoll, Andrew Hughes, and Martin Davey. The first draft of the manuscript was written by Conor O'Driscoll, and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

* Corresponding author. Tallaght University Hospital, Dublin, Ireland. Tel.: +353857884714.

E-mail address: coodris@tcd.ie

<https://doi.org/10.1016/j.artd.2022.11.001>

2352-3441/© 2022 The Authors. Published by Elsevier Inc. on behalf of The American Association of Hip and Knee Surgeons. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Total hip arthroplasty (THA) reliably alleviates pain and improves quality of life in patients with osteoarthritis [1]. Patient factors associated with a successful recovery include a high level of preoperative physical function and balanced muscular strength [2]. Patients with neurological conditions pose unique surgical

challenges, including contractures, paresis, and muscular imbalance, [3] ranging from flaccidity in conditions affecting the lower motor neurons to spasticity in those affecting the upper motor neurons [4]. Dislocation and aseptic loosening concerns led to a historical reluctance from surgeons to perform THA for patients with neurological conditions, with many proceeding towards salvage procedures in the primary instance such as arthrodesis or resection arthroplasty [5–8]. Operative techniques and implant designs have evolved to broaden the indications for THA, [9–13] with a greater understanding of the biomechanical environment surrounding prosthetic hips, which has naturally extended to the neurological population [3,14–16].

In 2009, Queally et al. identified that the clinical data pertaining to the outcomes of THA in the neurological population were lacking [3]. Such interventions have become increasingly more common over the past 10 years, and as such, the available literature has greatly expanded. The aim of this systematic review is to build upon the previous work of Queally et al and report on the complete literature relating to THA for patients affected by neurological conditions [3]. This review will serve as a guide for orthopaedic surgeons planning THAs in the neurologically impaired population and to enable an informed discussion with patients and their caregivers regarding potential complications and anticipated outcomes.

Material and methods

Search strategy and eligibility

In February 2022, a systematic review of the literature was performed by 2 independent reviewers (C.S.O.D. and A.J.H.) with respect to Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines [17]. The PubMed/Medline OVID, Cochrane, and Embase databases were screened from their inception to February 2, 2022, inclusively. The search strategy was adapted from the study by Queally et al. [3], with predetermined search terms utilized for each of the aforementioned databases, including THA Population, Neurological Impairment, and Outcome (see attached in [Appendix](#)).

Following the removal of duplicate studies, both independent reviewers manually screened the titles and abstracts of the returned studies while applying our predetermined exclusion criteria, with the senior authors (B.J.O.D. and J.M.Q.) acting as arbitrators in cases of discrepancy of opinion. Following the removal of excluded studies, both independent reviewers applied the predetermined inclusion criteria to the remaining studies to evaluate all potential studies for definitive inclusion. Thereafter, the reference lists of all included studies were screened for further studies that potentially may meet the inclusion criteria.

Inclusion criteria encompassed (1) studies reporting the outcomes of THA in patients with neurological conditions, (2) studies published in English language, and (3) published in a peer review journal with full text available. Exclusion criteria included (1) case reports, (2) review articles, (3) abstract-only studies, and (4) cadaveric or biomechanical studies.

Outcomes of interest

The results from each study were tabulated following a quality assessment using the GRADE tool (Grading of Recommendation, Assessment, Development and Evaluation) [18] and Oxford Centre for Evidence-Based Medicine criteria [19].

A predesigned data-collection template was then collated including (1) study population, including neurological condition, study type, follow-up period, patient demographics; (2) implants

utilized and surgical technique; (3) patient-reported outcomes; (4) complications; and (5) arthroplasty revision rate.

Descriptive statistics were performed using Stata software version 16.1 (StataCorp, College Station, TX). A Meta-analysis of the included studies was not performed due to the significant heterogeneity in study location, patient age, disease severity, implant type, surgical technique, and reporting of outcomes.

Results

There were 1820 studies collated in the initial database search, which was subsequently reduced to 1514 following duplicate removal. Following abstract screening, 108 full-text articles were assessed, leaving 45 studies, with 36,251 patients, included in the final review. Of the 45 studies, 36 reported functional outcomes of 1811 patients, using various rating scales, and all 45 reported complications. The PRISMA flow chart with reasons for exclusion is illustrated in [Figure 1](#).

Parkinson's disease

Parkinson's disease (PD) is a progressive neurodegenerative disorder characterized by elevated tone and classical motor symptoms such as bradykinesia, rigidity, rest tremor, as well as both postural and gait impairment, in addition to cognitive effects such as memory impairment which may complicate rehabilitation [20]. PD affects greater than 1% of the population older than 60 years, [21] and advances in medical care have led to increased life expectancy [22]. Between 2000 and 2014, the incidence of THAs performed for patients with PD increased from 946 to 1655 within the United States Nationwide Inpatient Sample (NIS) [14].

Focusing on the primary admission and immediate perioperative period, 2 studies using the United States NIS and carried out by Kleiner et al. and Newman et al. [14,23] observed an increased length of stay for PD patients compared to control. Medical complications were common in the PD cohort and included delirium and respiratory and urinary tract infections, replicating the experiences of PD patients undergoing nonorthopedic surgeries, thus highlighting the need for medical optimization and multidisciplinary care [24].

Eight studies with a total of 1296 THAs reported medium-term follow-up of over 2 years. These were primarily elective primary THAs (949 THAs), with several revision (58 THAs) and trauma (26 THAs) cases also classified. The mean age was 72.6 years, with a range of disability levels included from Hoehn and Yahr Classification stage I and II to severely affected grade IV and V patients [25].

Higher dislocation rates were reported in all studies comparing PD patients to control, ranging from 1.6% to 8.3% [26,27]. This followed through to higher revision rates due to recurrent dislocation in Joint Registry-based studies by Wojtowicz et al. in Sweden and Jansen et al. in Finland, for their cohorts of 495 and 297 PD THAs compared to matched control [28,29]. With respect to surgical indication, the dislocation rates in the trauma and revision THA group of the study by Weber et al., 12.2%, were higher than those in the elective group which experienced no dislocations [30]. Peri-prosthetic infection rates were over 2 times greater within the PD group in each of the 3 studies by Wojtowicz et al., Shah et al., and Rondon et al., which included matched control patients without neurological conditions [26–28].

Patient-reported outcome measures were documented in 6 studies [26,28,30–33]. Notwithstanding the higher complication and implant revision rates experienced by the PD group, an improvement in functional activity and pain was reported in each of the studies postoperatively. A point of note made by Weber et al. was that beyond an initial improvement in function at 1 year




Figure 1. PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-analyses) flowchart.

postoperatively, longer term outcomes were often limited by PD disease progression, as shown by a parallel deterioration in Hoehn and Yahr disease scores [30].

The results of THA in the PD population are displayed in Figure 2.

Cerebral palsy

Cerebral palsy describes a group of permanent disorders of movement and posture attributable to nonprogressive disturbances that occurred in the developing brain [34]. Spasticity is the most common movement disorder, occurring in 80%, [35] which may lead to periarticular contractures about the hip joint and migration of the femoral head leading to subluxation and dislocation [36]. Abnormal loading leads to dysplastic changes of the femoral head and acetabulum with associated pain and disability [3,37].

Within the primary admission period, Moon et al. using the US NIS database reported a longer length of stay and increased risk of perioperative medical complications despite a younger mean age at the time of surgery in their cohort of 2062 Cerebral Palsy (CP) patients matched 10:1 to a non-CP control group [38]. Over the first 90 days, Moore et al. in their study of 864 CP patients matched 4:1,

from the US Mariner patients records database, found a statistically significant increased odds ratio of medical complications such as urinary (odds ratio [OR] 2.42, 95% confidence interval 1.25–4.58) or respiratory tract infection (OR 3.77, 95% confidence interval 1.64–8.56) and periprosthetic fracture (OR 2.55, 95% confidence interval 1.42–4.46) [39].

Reviewing the England and Wales Joint Registry data from 389 CP patients undergoing THA, King et al. reported cumulative revision rates of 2.6% and 6.4% at 1 year and 5 years, respectively [15]. Hybrid implants had the lowest 5-year revision rate of 1.2%, with uncemented (7.1%) and resurfacing (11.5%) implants faring less favorably. For comparison, the unmatched control cumulative revision rate was 0.79% at 1 year and 2.9% at 5 years. Among seven other retrospective studies, at varying lengths of follow-up, revision rates ranged from 0% to 27%. These studies with a combined 189 patients at mean 94 months of follow-up, reported 20 dislocations (10.6%), 11 cases of aseptic loosening (5.8%), 6 infections (3.2%), and 6 periprosthetic fractures (3.2%) [40–46]. While there was considerable heterogeneity across the studies with regard to severity of CP symptoms, functional improvements were noted in multiple studies, including cases of both mild and severe impairment [41]. Results for the cerebral palsy THA population are displayed in Figure 3.

Author + Year	LO E	GRAD E	Population	Number	Implant/Technique	Mean Age (Years)	Follow Up (Years)	Functional Outcome	Survivorship	Complications
Shah N. V. et al. (2020) [21]	4	Low	Multicentre, New York State DOH Database	235, 1:1 Control	Not specified	74.3	Minimum 2	NA	Revision Rate, PD 6.4%; C 4.7%	Dislocation: PD 7.3%, C 6.2%; Wound Infection: PD 19.8%, C 6.2%; Prosthetic Joint Infection: PD 9.1%, C 3.1%
Wojtowicz, A. L., et al. (2019) [28]	3	Low/Moderate	Swedish Hip Arthroplasty Register, 1999-2012	490, 1:1 Control Matched	Implants Cemented, 442 (90%); Uncemented, 23 (5%); Hybrid, 6 (1%); Reverse Hybrid, 19 (4%)	73	4.7	EQ-5D Pre-op PD 0.32, C 0.40; 1 Year PD 0.62, C 0.82	Mortality: 90 Day, PD 0.62%, C 0.41%; 1 Year, PD 2.56%, C 2.11%; 9 Year, PD 54.33%, 28.05% Implant Revision Rate: 90 Day PD 1.03%, C 0.41%; 1 Year, PD 2.10%, C 0.41%; 9 Year PD 5.44%, C 1.75%	Complications leading to Revision, Dislocation, PD 11 (2.2%), C (0.2%); Aseptic Loosening, PD 5 (1%), C (0.6%); Prosthetic Joint Infection, PD 4 (0.8%), C (0.2%); Fracture, PD 2 (0.4%), C (0.2%)
Rong, X., et al. (2019) [31]	4	Low	West China Hospital, China, 2009-2016	Total 28, Subgroup 1, H&Y Stage I and II, 12 (43%); Subgroup 2, H&Y Stage IV and V, 12 (43%)	Not specified	65 (47-80)	4.3	HHS Pre-op: 30.00, Post-op, 71.39; SF-12(QOL) Pre-op PCS 13.96, MCS 17.96 Post-op PCS 17.54, MCS 21.41; WOMAC Pain (0-20) Pre-op 10.63, Post-op 2.96, Function (0-68) Pre-op 44.08, Post-op 34.63	Implant Revision Rate at 5 years: 5.9%	Periprosthetic Osteolysis and Polyethylene Wear, 1 (3.8%); Periprosthetic Fracture, 1 (3.8%); Infection/Dislocation/Aseptic Loosening, nil
Kleinert, J. E., et al. (2019) [41]	3	Low	Healthcare Cost and Utilization NIS Database, USA, 2006-2014	4001, 1:3 Control Matched	Not specified	74.5	Primary admission	NA	Mortality (in hospital) PD 0.50%, C 0.47%	All Complications (including dislocation, prosthetic loosening and PJI) PD 1.5%
Rondan, A. J., et al. (2018) [26]	3	Low/Moderate	Thomas Jefferson University, Philadelphia, 2008-2016	52, 93 Control Matched	Not specified	68.7	5.3	SF-12 (26 PD, 47 Control) Pre-op: PD 28.4, C, 34.4. Post-op: PD, 37.4; C, 44.7; Change: PD, 62.2; C, 11.8	Implant Survivorship 2 Year 94.3%; 5 Year 85.3%; 10 Year 78.7%	Periprosthetic Fracture 4 (7.7%); Dislocation 4 (7.7%); Aseptic Loosening 4 (7.7%); Periprosthetic Joint Infection 1 (1.9%)
Newman, J. M., et al. (2018) [23]	3	Low	NIS Database, Sample from over 1000 US Hospitals, 2002-2013	10528, 1:3 Control Matched	Not specified	73	Primary admission	NA	NA	Surgical Complication PDC Odds Ratio 1.31, Haematomata/Seroma OR 1.3, Peripheral Nerve Injury, OR 3.0, Irrigation/Debridement, OR 0.75; No infection or dislocation reported
Larocque, J. Y., et al. (2018) [32]	4	Low	Pine-Subpierre Hospital, France, 2002-2012	63 (59 patients), 42 Primary, 21 Revision	Approach: Direct lateral Implants Acetabular Cementless with DM, Femoral Primary, Uncemented 38 (90%) Modular Stem 4 (10%); Revision, Uncemented 17 (81%) Cemented 4 (9%)	72.5 (55-79)	8.3	Pain Relief at 2 year follow up: 53/57 rated good to excellent	Implant survivorship 2 Year 91.5%, 5 Year 79.7%	Periprosthetic Fracture, 4 (6.4%); Surgical Site Infection, 2 (3.2%); Dislocation, 1 (1.6%)
Sponer, P., et al. (2017) [33]	4	Low/Very Low	University Hospital Hradec Králové, Czech Republic, 2005-2012	Total 24 (10 Elective, 14 Trauma)	Elective Acetabular Cemented 8 (80%) Uncemented 2 (20%); Femoral Stem Cemented 9 (90%) Cementless 1 (10%) Trauma Acetabular Cemented 13 (93%), Cementless 1 (7%); Femoral Stem Cemented 12 (86%), Cementless 2 (14%)	Elective 74 (65-82), Trauma 76 (67-83)	Elective 6.8, Trauma 4.5	Merkel d'Anjou Score (0-18) Elective Pre-op: 8, 6 Months, 14, 36 Months, 14, Trauma Pre-op: 3, 6 Months, 12, 36 Months, 12	Nil Implant Revision	Dislocation E 1 (10%), T 1 (7.1%), Periprosthetic Fracture E 1 (10%), T 1 (7.1%), Prosthetic Joint Infection E 0, T 1 (7.1%)
Jansen, E., et al. (2014) [29]	4	Low	Multicentre, PERFECT Database, Finland, 2008-2009	297, 1:3 Control Matched. Note with 500 TKA	Implants Cemented 165 (55.6%)	72	1-13	NA	Implant Survival PD 1 Year, 98% (CI 96.4-99.6), 3 Year, 96.8% (CI 94.6-99.0), Implant Revision THA First 2 years post op PDC, 1.13; 1 (CI 0.50-2.56)	Complications leading to Revision, Dislocation 1 Year 3.36%, Total (6.06%, Relative Risk 1 year, PDC 2.23 Total 1.29)
Webster, M., et al. (2002) [30]	4	Low	Mayo Clinic, Rochester, 1970-1994	107 (98 Patients), 58 Primary, 49 Revision; Trauma, H&Y Stage Primary 1-19%, II-69%, III-10%, Unknown 2% Revision Trauma 1-6%, II-24%, III-65%, IV-4%, Unknown-2%	Approach: Anteriorlateral, 56 (52%), transtrochanteric, 36 (34%), posterior, 12 (11%); direct lateral, 3 (3%). Procedures: Trauma Acetabular Cemented 13 (93%), Cementless 1 (7%); Femoral Stem Cemented 12 (86%), Cementless 2 (14%)	72 (57-87)	Followed up over 2 years 9, 44 (65%); R/T 31 (65%)	Pain Good to Excellent relief 93%. Functional Primary Group Improvement noted during first year with subsequent deterioration in line with primary Parkinson's Disease related disability. 78% Neurological Progression 57% Hoehn and Yahr Stage 4 or 5.	Periprosthetic mortality, 4 (3.7%) LRTI, PE 1, CVA 1 THA Reoperated, 9 (8.4%) PJI 1, PPI 1, Trochanteric Nonunion 1, Instability 1, trochanteric wire removal 1, aseptic loosening 3. THA Survival 93% at 5 years for Primary and Total Groups	Dislocation Primary 0, Revision Trauma 0 (1.25%), Prosthetic Joint Infection P 0; R T 1 (2%), Trochanteric Nonunion P 2 (3.4%); R/T 2 (4%), DVT P 2 (3.4%); R/T 1 (2%), PE P 2 (3.4%); R/T 0, Transient Peroneal Palsy P 1 (1.7%); R/T 1 (2%)

Figure 2. Total hip arthroplasty in Parkinson's disease results.

Multiple sclerosis

Multiple sclerosis (MS) is a chronic inflammatory autoimmune condition primarily affecting the central nervous system via demyelination of the axonal sheaths, which disrupts transmission [47]. MS is the leading cause of nontraumatic disability in young adults, affecting approximately 400,000 in the United States alone [48]. Neuromuscular manifestations are common and vary according to the location affected [4]. These may include rigidity, spasticity, weakness, contractures, and functional limitation, with increased risk of falls and poor postural control [49].

Newman et al. found that during the primary admission, the rate of perioperative and surgical complications was higher for MS

patients than for the control, based on an US NIS Inpatient Sample of 5899 MS patients who underwent THA [16]. There was also an increased average length of hospital stay and likelihood of requiring admittance to step-down care facilities before returning home from hospital in the MS cohort.

Among those studies that reported THA complications individually, the 2018 retrospective review of Newman et al. with 41 THAs in the Cleveland Clinic observed dislocation, infection, and aseptic loosening at higher rates than those in control [50]. With regard to implant revision rates, Quinlan et al. estimated the rate at 2 years to be 4.23% from their analysis of US Medicare analytical files [51], while Rondon et al. reported a corresponding 2-year rate of 2.1% among their 62 THAs [52]. At longer term follow-up of 7 and 8

Author + Year	LO E	GRAD E	Population	Number	Implant/Technique	Mean Age (Years)	Follow Up (Years)	Functional Outcome	Survivorship	Complications
Moore, H.G. et al. (2021) [39]	4	Low	Marmor administrative database, USA, 2010-2018	864, 1:4 Control matched.	Not specified	56.3	90 day / 7 year	NA	5 year survival 94.2%, Control 95.2%	90 Day Dislocation 23 (2.7%) OR versus control 1.52, Surgical Site Infection 14 (1.6%) OR 0.72, Periprosthetic Fracture 23 (2.7%) OR 2.55 Kosmosium 131 (15.2%) OR 1.13
Moore, A. S., et al. (2020) [38]	4	Low	Healthcare Cost and Utilization NIS Database, USA, 2005-2014	2062, 1:10 Control Unmatched	Not Specified	CP 69.2, Control 64.8	Primary admission	NA	Not reported	Length of Stay, CP 3.8 days, Control 3.2 days. Surgical site infection CP 0.3%, C 0.1%, Perioperative Haemorrhage, CP 0.7%, C 0.8%, Postoperative Anemia CP 30.2%, C 24.4%, Mechanical Failure CP 0.74% C 0.45%, Overall CP 34.3%, C 27.9%
Hosada, M.T. et al. (2017) [40]	3	Low/Moderate	Mayo Clinic, Rochester, USA, 1990-2013	41, 139 With 2 Year Follow Up; 1:2 Control Matched. CP GMFCS I, 3 (7%), II, 18 (44%), III, 12 (29%), IV, 6 (15%)	Approach: Posterior 20 (49%), Anteriorlateral 19 (46%). Implants Acetabulum Uncemented all. Dual Mobility implant 5 (12%), Lipped Liner 2 (5%), Femoral Cemented, Femoral head augmentation 4 (10%), Procedures Adductor tenotomy 7 (17%), Psoas tenotomy 2 (5%)	49 (21-74)	7 (2-20)	Pain: Pre-op moderate to severe hip pain all Post-op moderate to severe pain nil. Ambulation Pre-op Independent 10%, Use of Aid 70%, Non-Ambulatory 19.5%, Post-op Independent 54%, Use of Aid 46%, Non-Ambulatory 0%. Flexion contracture greater than 15 degrees, Pre-op 9 (22%), Post-op 0. Harris Hip Score (0-100) Pre-op 36, Post-op 78	THA Revision 5/39 (12.8%) CP mean 3yrs. Aseptic loosening: 2 (5.1%); Recurrent instability: 1 (2.6%); Deep Infection: 1 (2.6%); Mean THA Survival, 2yr 92%, 5yr 88%, 10yr 81%, 15yr 81%	Dislocation: 3 (7.3%) OR 3.0; Aseptic loosening, 2 (4.9%) OR 1.0; Wound Dehiscence, 1 (2.4%); Deep Infection, 1 (2.4%) (OR 1.0); DVT, 1 (2.4%)
Morrin, C., et al. (2016) [41]	4	Low	Instinct Care, France, 2001-2014	40 THA on 31 pts. Non-ambulatory CP GMFCS V, 21 (52.5%) previous surgery including 9 (22.5%) adductor tenotomy, 9 (22.5%) femoral and pelvic osteotomies, 1 (2.5%) Proximal Femoral Replacement, 2 (5%) Ostoma Resection, 23 (57.5%) Scissors with Vertebral fusion	Approach Lateral with trochanteric implant, all Implants Acetabulum Dual Mobility implant, all 15 (37.5%) supported by femoral bone block, 5 (12.5%) with 4-hole plate. Femoral Small "dysplastic" implant, Cemented, 37 (92.5%) Uncemented, 3 (7.5%)	19.2 (13.5-31.7)	5.3 (0-7.25)	Function Independent Sitting Pre-op 5 (12.5%), Post-op 6 (15%), Pain Permanent Pre-op 16 (40%), Post-op 0. Pain Sitting Pre-op 20 (50%), Post-op 1 (2.5%). Pain on Transfer Pre-op 28 (70%) Post-op 0 Motion Flexion >80 Pre-op 19 (47.5%) Post-op 34 (85%)	Implant Revision: 6 (15%), Deep Infection with femoral loosening 1; Delayed Femoral Osteotomy 2; Greater Trochanter Attachment 1; Osteolysis 1; Dislocation (Intraoperative) 1.	Deep Infection, 2 (5%) (1 early, 1 late); Dislocation, 3 (7.5%); Osteolysis/Loosening, 2 (5%); Dislocation (Intraoperative) 1 (2.5%)
King, G., et al. (2016) [15]	3	Low/Moderate	Multicentre-England and Wales National Joint Registry, 2003-2012	389 (265 Surgeons) 425813 Control (453 Surgeons)	Implants Cemented CP 72 (18%), C 166654 (39%), Uncemented CP 163 (42%), C 161539 (38%), Hybrid CP 85 (22%), C 64701 (15%). Reverse Hybrid CP 14 (3.6%), C 10359 (2.5%). Resurfacing (all MoM) CP 55 (14%), C 22531 (5.3%)	CP 53 Control 69	Unspecified	Oxford Hip Score (0-48) CP (47 Complete Pairs), Pre-Op 12, 6 months 34, Control (92073 Complete Pairs), Pre-Op 18, 6 months 41, EQ-5D Health Scale (0-100) CP (43 Complete Pairs), Pre-Op 66, 6 months 70, Control (80341 Complete Pairs) Pre-Op 70, 6 months 80.	THA Implant Revision Total CP 22 (5.7%), C 9776 (23.5%), 1 Year CP 10 (2.6%), C 3212 (7.9%), At 5 years by implant type CP Population Cemented 1.5%, Hybrid 1.2%, Uncemented 1.1%, Resurfacing 1.1%, Mortality 90 Days CP 0.26%, C 0.54%, 1 Year CP 1.4%, C 1.6%, 3 Year CP 3.7%, C 5.2%, 5 Year CP 6.9%, C 10%	Complications Leading to Revision CP with prosthetic time incident rate (PIR) per 1000 patient years Periprosthetic Fracture 7 cases PIR 5.0 (CI 2.4-10.5) Control PIR 0.7 (CI 0.66-0.75) Aseptic Loosening 6 cases PIR 4.3 (CI 1.9-9.6) Control PIR 1.6 (CI 1.32-1.43) Pain 5 cases PIR 3.6 (CI 1.5-8.6) control PIR 1.37 (CI 1.32-1.43) Dislocation 4 cases PIR 2.9 (CI 1.1-7.0) control PIR 1.22 (CI 1.16-1.42)
Yoon, B.H., et al. (2015) [42]	4	Very Low	KEPCO Medical Centre, South Korea, 2005-2007	5 THA. Prior surgery, 2 (40%); Resection arthroplasty, 1; Open reduction, 1	Approach Posterolateral. Implants Uncemented Ceramic on Ceramic	35.9 (20.2-55.6)	6.8 (5.8-8.3)	Pain Complete Pain Relief 3 (60%). Reduction in Preoperative Pain 2 (40%). GMFCS Function Score Improvement 3 (60%), Unchanged 2 (40%)	Nil Revision	THA Revision Total 4 (27%), Aseptic loosening 2 (15.4%), Infection 1 (7.7%), Recurrent dislocation 1 (7.7%)
Sanders, R. J., et al. (2013) [43]	4	Low/Very Low	Sant Maartenskliniek, Netherlands, 2008-2010	10 (9 Elective, 1 Trauma) GMFCS II, 3 (30%), III, 3 (30%), IV, 4 (40%)	Approach Posterolateral. Implants Dual Mobility Cemented CP Adductor tenotomy, 1 (10%)	54 (43-61)	3.2	SF-36 (0-100) Post-op Physical Function 35, Pain 75.6, General Health 59.5.	Reoperation Periprosthetic Fracture 1 (10%)	Nil Dislocation
Schroeder, K., et al. (2010) [44]	4	Low	Heidelberg University Hospital, Germany, 1988-2004	15 (13 Patients)	Approach Transgluteal Lateral Baur Implants Uncemented 11 (73%) incl 1 cemented acetabular cup, Cemented 4 (27%), Procedures 6 (40%) incl 5 adductor tenotomy, 1 lengthening of adductor tendon, 4 psoas and rectus tendon release, 1 transposition of outer rotators	42 (32-58)	10.5 (2-18)	Pain (0-10) Pre-op 8.4, Post-op 1.1.	THA Revision Total 4 (27%), Aseptic loosening 2 (15.4%), Infection 1 (7.7%), Recurrent dislocation 1 (7.7%)	
Rapraet, B. S., et al. (2010) [45]	4	Low	Hospital for Special Surgery, USA, 1972-2006	59 (56 Patients) Prior surgery 37 (65%), 12 soft tissue release, 17 osteotomy, 1 resection arthroplasty, 2 arthrodesis	Approach Anterolateral, 5 (9%) Transstrochanteric, 14 (24%), Abductor tendon release if abduction preoperative limited to 30 degrees or less, 28 (47%), Implants Cemented 20 (34%) (early cases), Hybrid (cemented stem, uncemented acetabulum), 35 (59%), with screws placed in 17 (29%), Uncemented 2 (3%) S ROM Components 2 (3%), Supratorial augmentation with autologous femoral head bone graft 4 (7%)	30.7 (14-61)	9.7 (2-28)	Pain (0-10) Pre-op 8, Post-op 0.7 Function Same or improved in all cases pre-operative to post-operative, 52 (88%) return to GMFCS Level of function before onset of hip pain.	THA Revision Total 9 (15.3%), Dislocation/Subluxation 5 (8.5%), Loosening and Pain 2 (3.3%), Infection 1 (1.7%), Periprosthetic Fracture 1 (1.7%), Reported Proportion of subjects who had Prostheses removed, 5 (8.5%), 1 year 5.08%, 3yr 6.95%, 6yr 12.42%, 8yr 15.66%, 17yr 29.16%.	Dislocations 1 (1.7%), 2 early within 6 weeks, 6 late after 6 months. Infection 3 (5.0%), Periprosthetic Fracture 1 (1.7%), Aseptic Loosening, 4 (6.8%) S acetabular, 1 femoral. Trochanteric Nonunion 1 from 14 osteotomies (7%). Trochanteric Bursitis 5 (8.5%), PE 1 (1.7%).
Webster, M. and M. E. Cabanela (1999) [46]	4	Low/Very Low	Mayo Clinic, Rochester, USA	Total 16 (11 Elective, 5 Trauma)	Approach Anterolateral, 5 (30%) Transtrochanteric, 7 (44%), Posterolateral, 1 (6%), Implants Cemented, 12 (75%) Uncemented, 2 (12.5%), Hybrid, 2 (12.5%)	48.5 (22-79)	9.7 (2.5-21)	Pain 87% good to excellent relief. Function Improvement 79%	Revision, 1 (6.3%) aseptic loosening at 13 years, 2 (13%) repeat surgeries, 1 trochanter avulsion ORIF, 1 adductor tenotomy.	Periprosthetic Fracture, 2 (12.5%); Aseptic loosening 1 (6.3%); Nil Dislocation

Figure 3. Total hip arthroplasty in cerebral palsy results.

Author + Year	LO E	GRAD E	Population	Number	Implant/Technique	Mean Age (years)	Follow Up (years)	Functional Outcome	Survivorship	Complications
Quillen, N. D., et al. (2019) [51]	3	Low/V ery Low	Multicentre, Paardiver Patient Records Database from Medicare Standard Analytical, 2005-2014.	3360, 1:10 Control Matched	NA	65.6 (47.0% < 65, 69, 20.7%, 70-74, 15.2%, 75-80, 8.3%, 80-84, 5.0%, > 85, 2.0%.	NA	NA	In Hospital Mortality (1 Year) MS 230(6.6%), C 231 (0.69%), Revision THA (2 Year) MS 142 (4.23%), C 12.3(3%)	Hospital admission (30 day) MS 250(7.44%), C 127(3.79%), ED visit (30 day) MS 232(6.9%), C 100(4.79%), Prosthetic Joint Infection (2 Year) MS 162(4.82%), C 119(3.57%), Hip Dislocation (1 Year) MS 127(3.78%), C 78(2.34%), Length of Stay MS 3.73, C 3.46.
Rendon, A. J., et al. (2018) [52]	3	Low	Robman Institute Thomas Jefferson University, Philadelphia, USA, 2006-2016.	62 THA, 1:2 Control Matched. Note study also included TKA.	NA	57.2	6.2	SF-12 Note including TKA. Mean SF12 MS Group Pre-Op 26.6, Post-Op 37.7, Control Group Pre-Op 29.0, Post-Op 44.8	Implant Survival THA 2 year 97.9%, 5 year, 91.9%, 7 year 77.8%	Infection 2.1, 2%
Newman, J. M., et al. (2018) [50]	3	Low	Cleveland Clinic, Ohio, USA, 2008-2016.	41 (34 Patients), 1:2 Control Matched	Implants Un cemented Approach Anterior/Anterolateral MS 8 (20%), C 16 (20%), Direct Lateral MS 6 (15%), C 20 (24%). Posteroslateral MS 27 (66%), C 46 (56%).	57 (38-79)	4 (2-8)	Functional AJR In last follow up mHHS (0-100) MS 66, C 80. HOS JR (0-100) MS 79, C 88. Mean Physiotherapy Duration MS 5wk, C 3wk. Mean Return to Baseline MS 7wk, C 5wk.	Implant Revision 8 Year MS 3 (7.3%), PJI and multiple dislocations 1 (2.4%), Periprosthetic Fracture 1 (2.4%), Aseptic Loosening 1 (2.4%), Control 1 (1.2%), Periprosthetic Fracture MS 1 (2.4%), C 0. Periprosthetic Fracture MS 1 (2.4%), C 1 (1.2%)	Length of Stay MS 4 days (2 - 8), C 4 days (1 - 19). Dislocation MS 4 (9.8%), C 0. Aseptic Loosening MS 3 (7.3%), C 0. Prosthetic Joint Infection MS 2 (4.9%), C 0. Wound Infection MS 1 (2.4%), C 0. Stress Fracture MS 1 (2.4%), C 0. Periprosthetic Fracture MS 1 (2.4%), C 1 (1.2%)
Gutman, J. M., et al. (2018) [53]	4	Low/V ery Low	NYU MS Care Centre, New York, USA, 2012 - 2016.	13 THA (Elective 10 Elective, 3 Trauma). Note study also included TKA.	Approach/Components not specified	53	3;75 (0.25-12.7)	Ambulatory Aid Requirements Reduction, 5 (38%), No Change, 3 (23%), Increase, 5 (38%) (following fracture in 3)	Revision 1 (7.7%), Recurrent Instability	Not Reported Specifically for THA.
Newman, J. M., et al. (2017) [46]	3	Low/V ery Low	Multicentre Healthcare Cost and Utilization NIS Database, USA, 2002-2013.	8899, 1:3 Control Matched	NA	57 (SD 10.9)	Primary Admission	NA	NA	Length of Stay (Days) MS 3.55, C 3.41. All peri-operative complications OR 1.37. Any Surgical Complication OR 1.18. Any Medical Complication OR 1.55

Figure 4. Total hip arthroplasty in multiple sclerosis results.

years, revision rates of 22.2% and 7.3%, respectively, were reported [50,52].

Newman et al. also reported variable functional outcome scores depending upon MS phenotype, with a higher mean Hip Disability and Osteoarthritis Outcome Score Joint Replacement score of 95 in the primary progressive subtype, as opposed to 70 in the secondary progressive group [50]. MS patients demonstrated lower modified Harris Hip Score and Hip Disability and Osteoarthritis Outcome Score Joint Replacement hip function scores than control despite receiving increased physiotherapy care although this may be partly reflective of progression of their disease during the period of this study as demonstrated by worsening of mMSIS scores. Results from the included MS THA studies are displayed in [53] Figure 4.

Poliomyelitis

Poliomyelitis is an infectious viral disease caused by the polio enterovirus, in which loss and degradation of anterior horn cells in the lower motor neuron system leads to varying degrees of muscle wasting, hyporeflexia, and flaccid paralysis [54]. Hip joint instability and muscular imbalance, [55] particularly gluteus medius weakness, [56] may lead to subluxation and abnormal loading throughout hip development during childhood, with resultant bony dysplasia and painful degenerative hip arthritis [3]. Commonly associated leg length discrepancy [56–59] and soft-tissue envelope laxity pose additional challenges when seeking optimum tension intraoperatively [60].

The largest study was by DeDeugd et al. in the Mayo Clinic including 59 patients spread over 42 years [60]. They found an improvement in functional outcomes both in a polio-affected and

unaffected limb. Their component revision rate at 6-year follow-up was 5.1%, with osteolysis (10.2%) and dislocation (5.1%) being the most commonly encountered complications. A further 8 retrospective case series with a total of 78 THAs also reported improved functional outcomes through a variety of scoring systems, [56–59,61–64] which are detailed in Figure 5.

Charcot hip

Charcot or neuropathic arthropathy occurs in patients with reduced sensory and nociceptive feedback in a joint susceptible to repetitive microtrauma, leading to progressive joint destruction and deformity [65]. Five studies met our inclusion criteria which reported upon THA outcomes in the neuropathic or “Charcot” hip population [65–69].

The largest of these was from Chalmers et al. in 2018 who described a case series of 12 THAs in 11 patients, with a range of underlying medical conditions, including Charcot-Marie-Tooth disease and diabetes mellitus [65]. They reported considerable improvements in pain and function, which correlated with a mean rise in HHS at 5-year follow-up, notwithstanding a high rate of complications, as seen in Figure 6.

Neurological assorted

Further studies reported experiences with THA in the setting of an assortment of other neurological conditions, such as stroke, traumatic brain injury, spinal cord injury, brain tumors, and spondylotic neuropathy. A number of these studies grouped multiple underlying conditions together. Although these studies did not

Author + Year	LO E	GRAD E	Population	Number	Implant/Technique	Mean Age (years)	Follow Up (Years)	Functional Outcome	Survivorship	Complications
Zhang, T.F., et al. (2021) [61]	4	Low	The First Affiliated Hospital, Guangzhou, China, 2006-2016	17, (100% on affected limb, Trauma	Approach Posteroslateral Implants Un cemented	66.95	6.4	Oxford (0-48) Pre-op 32.9, Post-op 36.2. UCLA (0-16) Pre-op 4.9%, Post-op 3.9%	Nil revision, Repeat Operation, 1 (5.9%) (Dislocation, Closed reduction)	Dislocation , 1 (5.9%), Periprosthetic Fracture , 1 (5.9%), Nil aseptic loosening/infection.
Sankatsis, M., et al. (2018) [50]	4	Very Low	Saga University Hospital, Japan, 2001-2011.	6 (5 Patients), 2 (40%) on paralytic side, 4 (60%) on non-paralytic side. Previous surgery 6 (100%), 2 foot arthrodesis, 1 achilles tendon lengthening, 1 hip osteotomy and unspecified (unkn surgery)	Approach Posteroslateral Implants Un cemented	54.7 (49-63)	8.4 (4.5-15)	Oxford Orthopaedic Association Hip Score (0-100) Pre-op 42.8, Post-op 78.8	Nil revision	Nil Dislocation/loosening/infection/nerve palsy observed
DeDeugd, C. M., et al. (2018) [60]	4	Low	Mayo Clinic, Minnesota, USA, 1970-2012.	59 (21 Patients) 32 (54%) on polio affected limb, 27 (46%) on polio unaffected limb	Approach Anterolateral, 36 (61%); Posteroslateral, 23 (39%). Implants Femoral Uncemented, 38 (64%) used in modern cases. Cemented, 21 (36%) used in older cases. Acetabular Uncemented, 58 (98%), Cemented, 1 (2%). Elevated or face changing acetabular lining used in 3 (5%)	66 (38-88)	6 (2-20)	HHS (0-100) Affected Side Pre-op 23, Post-op 81. Unaffected Side Pre-op 54, Post-op 80. Hip Flexion Affected Side Pre-op 73, Post-op 99, Unaffected Side Pre-op 89, Post-op 96. Hip Abduction Affected Side Pre-op 15, Post-op 21. Unaffected Side Pre-op 22, Post-op 28.	Component Revision 3 (9%), Osteolysis and Loosening 2 (3.4%), Trochanteric osteolysis, 1 (1.7%), Repeat Operations, 2 (3.4%) Periprosthetic Fracture, Open Reduction Internal Fixation.	Osteolysis 0 (10.2%), 3 Revised, 3 Unrevised. Dislocation , 3 (5.1%), Periprosthetic Fractures , 2 (3.4%), Common peroneal nerve palsy 1 (1.7%), Superficial Wound Dehiscence, 1 (1.7%), Superficial Surgical Site Infection 1 (1.7%)
Sobrin, F. B., et al. (2017) [57]	4	Very Low	Hospital General Universitario Gregorio Marañon, Madrid, Spain, 2008-2012.	5, (100% on polio affected limb	Approach Posteroslateral 4 (80%) Trochanteric 1 (20%) Implants Uncemented 5 (100%). Dual Mobility Cup 2 (40%), Autologous Bone graft 2 (40%), Modular Femoral Stem 2 (40%).	47 (38-64)	4.6	HHS (0-100) Pre-op 30, 6 Months 75, 2 Year 81. Limb Length Discrepancy (Median) , Pre-op 30mm, 3 Months 17mm.	revision 1 (20%), Recurrent dislocation	Periprosthetic Fracture , 1 (20%) Dislocation , 1 (20%) Pseudotumor, 1 (20%) - at site of greater trochanter osteotomy in trans-trochanteric approach.
Faldini, C., et al. (2017) [62]	4	Low/V ery Low	Rizzoli Orthopaedic Institute Bologna, Palermo, Italy, 1999-2011.	14, Elective THA	Approach Direct Lateral. Implants Uncemented 13 (93%), Hybrid (Cemented Femoral and Uncemented Acetabular) 1 (7%).	51 (26-66)	7.7 (4.3-13)	HHS (0-100) Pre-op 52 (32-78), Post-op 83.3 (72-91).	Implant Revision , 2 (14.3%) Periprosthetic Fracture, 1 (7.1%), Aseptic loosening 1 (7.1%).	Transient Sensory Sciatic Nerve Palsy, 1 (7.1%) following limb lengthening of 34mm, Aseptic Loosening, 1 (7.1%), Periprosthetic Fracture , 1 (7.1%).
Choi, Y. J., et al. (2016) [58]	4	Very Low	Kyung Hee University, South Korea, 2004-2012.	11, 4 (36%) Polio affected side, 7 (64%) Polio unaffected side	Approach Posterior Implants Un cemented	57 (41-64)	6.7 (2.7-10.9)	HHS (0-100) Pre-op 52.5 (21.78), Post-op 85.8 (70-100) UCLA (0-16) Pre-op 3.9 (2.7) Post-op 6.2 (4.10). WOMAC Score (0-68) Pre-op 52.4 (41-69), Post-op 12.5 (6.55).	Repeat operations, 1 (9%), Rotational osteotomy of ipsilateral femur due to in-toeing	Dislocation , 1 (9.1%), Instability , 1 (9.1%), Nil loosening/osteolysis/infection/periprosthetic fracture.
Bertoni, M. et al. (2016) [63]	4	Low	Italian Hospital of Buenos Aires, Argentina.	6, (100% on polio affected limb, Elective THA, 2 (1) prior achilles tendon lengthening, 1 (1) prior pelvic and femoral osteotomy	Approach Posteroslateral, 5 (83%); Transtrochanteric, 1 (17%) Implants Cemented 4 (67%), Hybrid 1 (17%), Uncemented 1 (17%).	51.3	10 (7-12)	HHS (0-100) Pre-op 67.6, Post-op 87.3. VAS (0-10) Pre-op 7.7, Post-op 2.	Dislocation/Instability	Dislocation , 1 (33%), Nil osteolysis/aseptic loosening/fracture/infection
Yoon, B. H., et al. (2014) [59]	4	Very Low	Seoul National University Bundang Hospital, South Korea, 2000-2009.	10, 4 (40%) on polio affected limb, 6 (60%) on polio unaffected limb, 5 (50%) prior achilles tendon lengthening, 3 (30%) prior foot arthrodesis.	Approach Posteroslateral 9 (90%) Transgluteal Lateral 1 (10%) Implants Un cemented	48 (32-59)	7 (3.4-13)	HHS (0-100) Pre-op 70, Post-op 92. Pain (0-10) Pre-op 7, Post-op 0.9. Limb Length Discrepancy in cm Pre-op 2.1 (0-4), Post 1.7 (0-4).	Nil reported	Dislocation , 1 (10%), Nil osteolysis/aseptic loosening/fracture/infection reported
Hosalkar, H. S., et al. (2010) [64]	4	Very Low	Rady Childrens Hospital, San Diego, USA, 1991-2005.	9 THA 8 Patients. Note study also included TKA.	Approach Posterior Implants Un cemented	66.5 (52-77)	Over 2 Years	HHS (0-100) Pre-op 94, Post-op 173. Range of Motion Flexion/Extension Pre-op 20-83, Post-op 0-110.	Nil Reported	Nil Dislocation observed.

Figure 5. Total hip arthroplasty in poliomyelitis results.

Author + Year	LO E	GRAD E	Population	Number	Underlying Neurological	Implant/Technique	Mean Age (Years)	Follow Up (Years)	Functional Outcome	Survivorship	Complications
Inoue, D. et al. (2018) [65]	4	Very Low	Orthopedic Department, Kitazawa University, Japan.	2	Congenital insensitivity to Pain	Approach Posterolateral Implants Uncemented	38	0.5	Functional improvement ambulation 1 (50%)	NA	Dislocation, 1 (50%)
Chalmers, B. F. et al. (2018) [65]	4	Low	Mayo Clinic, Minnesota, USA, 2007-2014.	12 (11 Patients)	Charcot-Marie-Tooth 3 (25%), TDM 3 (25%), Spinal Cord Injury 2 (17%), Lysosomal storage disorder 1 (8%), Guillain-Barre 1 (8%), Head Injury 1 (8%), CVA 1 (8%).	Approach Anterolateral 9 (75%), Posterolateral 2 (25%), Implants Acetabular uncemented with adjunct screw fixation, 2 (17%) Dual Mobility, Femoral Uncemented 11 (92%), Cemented 1 (8%).	54 (31-79)	5 (2-9)	HHS (0-100) Pre-op 43 (34-56), Post-op 81 (67-90). Pain Pre-op No pain nil, Mild pain 4 (33%), Moderate Pain 7 (58%), Severe pain 1 (8.3%). Post-op No Pain 9 (75%), Mild Pain 3 (25%), Ambulation Pre-op Wheelchair 5 (45%), Full time aid 6 (50%), Post-op Wheelchair 0, Full time aid 7 (64%), Aid for long distance 2 (18%), Unaided 2 (18%), Limb Length Discrepancy Pre-op 36.7mm, Post-op 6.7mm	Revision 4 (25%), Recurrent dislocation 2 (17%), Aseptic femoral loosening 1 (8%), Periprosthetic Fracture (8%).	Dislocation, 2 (17%), Deep Prosthetic Joint Infection, 2 (17%), Superficial Surgical Site Infection, 1 (8%).
Rapala, K. and M. Oksela (2007) [67]	4	Very Low	Konarskiego Department of Orthopedics, Poland, 1994-1995	3 (2 patients)	Syphilis	Implants Cemented	NA	9.75 (9.5-10)	Improvement all	Revision 1 (33%), recurrent dislocation	Dislocation 2 (67%)

Figure 6. Total hip arthroplasty in Charcot joint results.

differentiate results by the underlying condition, the functional results were positive in each case.

The outcomes of THA on the affected hemiplegic limb following stroke were investigated by Henawy and Badie through a retrospective review of 24 patients in both the trauma and elective settings between 2013 and 2015 [68]. Following stroke, patients may experience upper motor neuron signs such as spasticity in the affected limb, posing similar challenges to that of CP, both intra-operatively and in the rehabilitation period. Excellent functional improvements were reported following arthroplasty, and they quantified this using the Harris and Merle d'Aubigne Hip Scores [68]. Studies by Wang et al., Ryu et al., Abdelazim and Michael, Alesh et al., and Park et al. also included patients with a history of stroke among their neurological THA groups, [69–75] which are detailed in Figure 7.

Discussion

Surgery setting

Often, arthroplasty surgeons operate on a limited number of patients with neurological conditions. A British joint registry-based study of 389 THAs for CP patients by King et al. found that only 23% of surgeons performed more than 1 THA procedure on a CP patient during the study period of 2003-2012 [15]. Across the arthroplasty field, it has been shown that a higher surgeon procedure volume is associated with lower dislocation and revision rates [76,77]. A study of nearly 38,000 Canadian patients found that surgeons who performed <35 THAs a year had a

dislocation rate of 1.9% vs 1.3% for surgeons with greater volumes [77]. While patient and caregiver preference may favor a local hospital with accessible follow-up, it is worth considering that arthroplasty procedures in patients with neurological conditions should ideally be performed by specialists with sufficient case volume. In addition, given the increased perioperative care needs as detailed in the following section, centers with well-integrated multidisciplinary teams are preferable. This can create a virtuous circle of specialty knowledge, surgeon's and theatre staff's familiarity with novel components, and multidisciplinary expertise.

Perioperative care

Advance THA planning allows for medical optimization, surgical planning with possible lead in time for implant delivery, as well as patient and caregiver education. Multidisciplinary input from health-care professionals is crucial. Many patients with neurological conditions, such as PD, MS, and CP, receive regular disease-modifying medication. A perioperative neurological assessment can help optimize their regimen to minimize spasticity and tremor. Patients with spasticity may also benefit from botulinum toxin injection prior to THA [40,78]. Postoperatively, patients are at increased risk of medical complications such as delirium, respiratory tract infections, falls, and periprosthetic fractures, alongside an increased length of stay, [14,16,23,38] and dedicated medical care helps mitigate these risks [79].

Higher order 3-dimensional imaging, such as CT scans of the hip and pelvis, can be used for accurately framing bony dysplasia, [80] commonly encountered in neurological conditions associated with

Author + Year	LO E	GRAD E	Population	Number	Underlying Neurological	Implant/Technique	Mean Age (Years)	Follow Up (Years)	Functional Outcome	Survivorship	Complications
Wang, Y. et al. (2021) [69]	4	Low	The 3 rd Hospital of Hebei, China, 2014-2019	52, Trauma	CVA 37 (71%), Parkinson Disease 6 (12%), Polioomyelitis 5 (10%), Epilepsy 2 (4%), Dementia 2 (4%)	Approach Posterolateral	71.1 (60-85)	3.5 Minimum	HHS (0-100) Post-op 82.3 VAS (0-10) Post-op 1.6	Revision 0, Reoperation 5 (10%), Periprosthetic Fracture 2 (4%) ORIF, Dislocation 3 (6%) Closed Reduction.	Dislocation 3 (6%), Periprosthetic Fracture 2 (4%), Heterotopic Ossification 2 (4%), Nil Loosening/Infection
Ryu, H.G. et al. (2021) [70]	4	Low	Guro Hospital, Seoul, South Korea, 2013-2015	35, Trauma, Control 127	Unspecified, including Parkinson's Disease, Cerebral Palsy, Polioomyelitis, Hemiplegia, Paraplegia.	Approach Posterolateral 20 (57%), Anterolateral 15 (43%), Implants Uncemented Dual Mobility	77.6 Control 76.2	6.2 (5-7.3)	HHS (0-100) Post-op 81.3, Control 82.2. UCLA (0-10) Pre-op 4.6, Post-op 3.7 Control Pre-op 4.7, Post-op 4.1.	Revision unspecified Reoperation 3 (8.6%), 2 dislocation, 1 superficial wound infection	Dislocation 2 (5.7%), Control 5 (3.9%) Infection 1 (2.9%) Loosening/Periprosthetic Fracture unspecified
Henawy, A. T., & Abdel Badie, A. (2017) [68]	4	Low	Suez Canal University Hospital, Egypt, 2013-2015	24 (16 Trauma, 8 Elective).	CVA, Mean Time post-stroke 10 months (6-67)	Approach Lateral transgluteal. Implants Acetabulum Uncemented Dual Mobility, all Femoral Cemented, 18 (75%), Uncemented, 6 (25%)	68 (53-79)	Minimum 1 year Range 1-2	Merle D'Aubigne (0-18) Pre-op 8 (0-10), Post-op 17 (14-18). HHS Hip Score (0-100) Pre-op 36 (0-73), Post-op 94 (88-100)	Revision, 1 (4%) Prosthetic Joint Infection	DVT, 3 (12.5%), Prosthetic Joint Infection, 1 (4%), Dislocation, 0, Osteomyelitis, 4 (17%), Loosening, 0.
Blitzard, D. J., et al. (2016) [71]	4	Low	Multicentre Database, Medicare Standard Analytical Files, 2005–2011.	CSM 6021, CSM + D 1173, Control 707460.	Cervical spondylotic myelopathy with subgroup of CSM previously decompressed	NA	<65 CSM (19%), CSM+D (20%), Control (9.5%), 65-69 CSM (30%), CSM+D (34%), Control (22%), 70-74 CSM (38%), CSM+D (41%), Control (22%), 75-79 CSM (28%), CSM+D (28%), Control (20%), 80-84 CSM (19%), CSM+D (14%), Control (15%), >85 CSM (9%), CSM+D (3%), Control (1%)	Minimum 1 year	NA	90 Day: Periprosthetic Infection CSM 1.8%, CSM + D 2%, Control 1.1%, Arthroscopy I&D CSM 1.5%, CSM + D 2.2%, Control 0.8%, Dislocation CSM 3.5%, CSM + D 3.1%, Control 1.5%, Periprosthetic Fracture CSM 1.4%, CSM + D 1.5%, Control 0.7%, Wound Complication CSM 2%, CSM + D 3.2%, Control 0.7%, Overall Follow Up: Periprosthetic Infection CSM 4.6%, CSM + D 6.6%, Control 2.5%, Arthroscopy I&D CSM 3.1%, CSM + D 3.7%, Control 1.5%, Dislocation CSM 5.7%, CSM + D 6.1%, Control 2.8%, Periprosthetic Fracture CSM 2.8%, CSM + D 3.1%, Control 1.6%, Wound Complication CSM 4.9%, CSM + D 7.5%, Control 2.2%.	90 Day: Periprosthetic Infection CSM 1.8%, CSM + D 2%, Control 1.1%, Arthroscopy I&D CSM 1.5%, CSM + D 2.2%, Control 0.8%, Dislocation CSM 3.5%, CSM + D 3.1%, Control 1.5%, Periprosthetic Fracture CSM 1.4%, CSM + D 1.5%, Control 0.7%, Wound Complication CSM 2%, CSM + D 3.2%, Control 0.7%, Overall Follow Up: Periprosthetic Infection CSM 4.6%, CSM + D 6.6%, Control 2.5%, Arthroscopy I&D CSM 3.1%, CSM + D 3.7%, Control 1.5%, Dislocation CSM 5.7%, CSM + D 6.1%, Control 2.8%, Periprosthetic Fracture CSM 2.8%, CSM + D 3.1%, Control 1.6%, Wound Complication CSM 4.9%, CSM + D 7.5%, Control 2.2%.
Abdelrazek, H. and F. Michael (2015) [72]	4	Low	Ain Shams University, Egypt, 2012-2014.	36 (12 Trauma, 18 Revision)	CVA 12 (47%), Parkinson Disease 6 (20%), Polioomyelitis 6 (20%), Brain Tumor 2 (7%), MS 2 (7%)	Approach Modified lateral. Implants Acetabulum Cemented Dual Mobility, Femoral Unspecified	64.6 (48–79)	1.1	Functional Evaluation Follow up results Excellent 7 (23%), Good 12 (40%), Fair, 6 (20%); Poor, 4 (13%)	Mortality, 1 - CVA 6 months post op, Nil implant revision reported	Nil Dislocation/Prosthetic Joint Infection
Li, J., et al. (2014) [73]	4	Low	Changshu Hospital, China, 2007–2009.	12, Trauma	Parkinson Disease 8 (67%), Polioomyelitis 4 (33%)	Approach Posterolateral Implants Uncemented Large Diameter Metal on Metal.	65.7 (56-76)	5 (3-6.6)	HHS (0-100) Pre-op 10.1, Post-op 76.4. UCLA (0-10) Pre-op 2.3, Post-op 6.7.	Nil Revision	Nil Dislocation/Infection/Fracture/Nerve Injury/DVT/loosening observed Heterotopic Ossification, 1 (8.3%)
Alesh, H., et al. (2014) [74]	4	Low	University of Pennsylvania USA, 1993-2011.	30 THA, 27 Patients	Cerebral Palsy 12 (40%), Traumatic Brain Injury 9 (30%), CVA 3 (10%), MS 2 (7%), Spinal Cord Injury 1 (3%).	Approach Posterolateral. Implants Acetabular Uncemented, Constrained liner 2 (7%), Augmentation 2 (7%), Femoral Uncemented, with modular components 3 (10%), Cemented procedure, Adductor Tenotomy 11 (37%) (7 CP, 4 Other), Hipoplasty lengthening 6 (20%), Achilles tendon lengthening 6 (20%) (5 CP, 1 Other), Heterotopic ossification excision 7 (23%) (1 CP, 6 Other)	48.6 (SD 12.2)	2.5 (2.1-12.1)	Pain Score (Minimal 1-2, Mild 3-7, Severe 8-10) Pre-op Severe 30 (100%), Post-op 29, Minimal 14 (50%), Mild 11 (39%), Severe 3 (11%). HHS (0-100) Pre-op 15 (5-35), Post-op 67 (24-91).	Implant Revision 1, (3.3%) Reoperation Arthroscopy due to Deep Infection	Nil Dislocation reported. Periprosthetic Fracture, 1 (3%), Infection, 4 (13%) - Deep 1, superficial 3
Park, K.S. et al. (2014) [75]	4	Low	Cheonam National University Hoswan Hospital, South Korea, 2008-2010.	19, Trauma	CVA, 15 (79%) (11 Haemorrhagic), Parkinson Disease, 4 (21%)	Approach Modified Minimally Invasive 2 Incision, Implants Uncemented, Large Diameter head >38mm, metal on metal Cemented procedure, Adductor tenotomy 4 (22%), Prophylactic femoral cabling 6 (32%).	72.6	1.4	Latest Follow Up: HHS (0-100) 41.5, WOMAC (0-68) 42.9.	Nil Implant Revision	Dislocation/Infection/Periprosthetic Fracture - Nil

Figure 7. Total hip arthroplasty in assorted neurological conditions results.

hip dysplasia, or in cases where there is concern for acetabular bone loss [81]. CT scanning may also be required if surgeons plan to use image-based navigation or robotic systems, which may be helpful in aiding accurate implant positioning in particularly challenging cases [82].

Physiotherapists can develop an individualized prehabilitation program tailored to the patient's capabilities. Advance occupational therapy assessment can anticipate assistive supports or ambulatory devices and address potential barriers to discharge from hospital, particularly for patients who experience concomitant upper-extremity spasticity and deformity [75]. Preoperative patient education regarding muscle rehabilitation and hip movement restrictions has also been shown in other arthroplasty settings to reduce the risk of dislocations [83]. Specialized postoperative physiotherapy can aid a challenging rehabilitation process, [84] with points of focus including maintenance of hip precautions, achieving a stable gait pattern, and maximizing neuromuscular control [75].

Operative considerations

The most common complication observed in our review was dislocation, and reducing its risk is a major focus of operative planning in patients with neurological conditions. Other complications such as infection were also observed at an increased rate within the neurological population, and standard measures to mitigate them, such as medical optimization, antibiotic prophylaxis, operating environment, and meticulous asepsis, are advised [85].

An uncemented prosthesis were favored in 69% of studies throughout our review. In those patients with particularly high risk of dislocation, such as severe Gross Motor Function Classification System level V CP, advanced PD, and poliomyelitis, implant designs such as dual-mobility (DM) implants and acetabular liners, with additional levels of constraint, were utilized. DM implants have a broad application in high-risk revision surgeries and are increasing in popularity. Neurological conditions in which DM implants have been used successfully over the last 10 years include PD, [32] CP, [40,41,43] polio, [57] Charcot arthropathy of the hip, and stroke [65,69,70,72]. The modular design of DM cups also allows for screw placement for additional cup fixation in cases of significant bone loss, which has been reported as a challenge when performing THA for patients with neurological conditions [45,65]. Lazennec et al. utilized DM components in their study of 63 PD THA cases, of whom 33% were undergoing revision arthroplasty and reported a single dislocation episode (1.6%) at a mean 8.3 years of follow-up [32]. Dislocation events observed in the setting of DM constructs cite the use of smaller outer polyethylene insert diameter (<38 mm), which corresponds to a smaller inner head size and cup malposition [86]. Morin et al. also reported a low dislocation rate of 2.5% in their group of 40 high Gross Motor Function Classification System grade 5 cerebral palsy THAs; however, this event was associated with an intraprosthesis dissociation requiring operative revision [41]. Intraprosthesis dissociation requiring operative reduction was also reported by Ryu et al. in their study of 35 patients affected by neurological conditions [70].

Constrained acetabular liners were used in limited numbers throughout the studies [33,40,56,60,75]. Such liners may be beneficial in reducing dislocation by containing the femoral head beyond its equator, thus preventing the head from dislocating out of socket. However, concerns arise regarding the significant reduction in primary arc range when constrained liners are utilized, which may paradoxically increase the risk of instability, and there is a high failure rate with poor initial placement, through liner dissociation, component loosening, and recurrent dislocations [87].

A simpler measure, widely adopted throughout the reviewed studies, is to use femoral heads with a larger diameter of greater than 36 mm, which increases the head-neck ratio, while also increasing the primary arc range prior to an impingement event, which could result in a dislocation [88]. Favorable results have been shown in large joint registry studies and have been adopted for neurological patients undergoing THA [45,89].

Surgical approach

There is no clear consensus regarding the optimal surgical approach, and thus, further research in this area would be of benefit. Of the included studies that detailed their surgical approach, the posterior approach was favored most in 14 of the 25 studies, highlighting the necessity to achieve a robust soft-tissue repair to minimize dislocation risk [90]. Achieving a robust envelop is difficult due to the degree of soft-tissue laxity in common neurological conditions, and for this reason, surgeons made use of additional combined anteversion and extended post-operative restrictions such as abduction braces and knee immobilizers, while hip spica casts were prevalent in the past [40,45].

Limitations

There are a number of limitations regarding the findings of this review. Bias may have been introduced in the reporting of complications by virtue of the retrospective nature of included studies which are heterogenic by nature. Our initial search found a number of studies from the same institution with overlapping patient groups, whereby the series of the largest cohort was selected for inclusion. The statistical analysis for subgroups was limited due to discrepancies in reporting of results among the included studies.

Conclusion

THA reliably improves symptoms of painful hip arthritis for patients with neurological conditions affecting the hip. It is important for surgeons, patients, and caregivers to be aware of the increased risk of associated complications, most notably dislocation with specific risk data presented herein, before proceeding with surgery. As a technically challenging operation, it should be undertaken by experienced arthroplasty surgeons allowing for familiarity with novel techniques and implants, with sufficient multidisciplinary support to meet their perioperative care requirements.

Conflicts of interest

The authors declare there are no conflicts of interest.

For full disclosure statements refer to <https://doi.org/10.1016/j.artd.2022.11.001>.

References

- [1] Molina CS, Thakore RV, Blumer A, Obrebsky WT, Sethi MK. Use of the national surgical quality improvement program in orthopaedic surgery. *Clin Orthop Relat Res* 2015;473:1574–81. <https://doi.org/10.1007/s11999-014-3597-7>.
- [2] Buirs LD, Van Beers LW, Scholtes VA, Pastoors T, Sprague S, Poolman RW. Predictors of physical functioning after total hip arthroplasty: a systematic review. *BMJ Open* 2016;6:e010725. <https://doi.org/10.1136/bmjopen-2015-010725>.
- [3] Queally JM, Abdulkarim A, Mulhall KJ. Total hip replacement in patients with neurological conditions. *J Bone Joint Surg Br* 2009;91:1267–73. <https://doi.org/10.1302/0301-620X.91B10.22934>.
- [4] Pomeroy E, Fenelon C, Murphy EP, Staunton PF, Rowan FE, Cleary MS. A systematic review of total knee arthroplasty in neurologic conditions:

- survivorship, complications, and surgical considerations. *J Arthroplasty* 2020;35:3383–92. <https://doi.org/10.1016/j.arth.2020.08.008>.
- [5] Castle ME, Schneider C. Proximal femoral resection-interposition arthroplasty. *J Bone Joint Surg Am* 1978;60:1051–4.
- [6] Root L, Goss JR, Mendes J. The treatment of the painful hip in cerebral palsy by total hip replacement or hip arthrodesis. *J Bone Joint Surg Am* 1986;68:590–8.
- [7] Flynn JM, Miller F. Management of hip disorders in patients with cerebral palsy. *J Am Acad Orthop Surg* 2002;10:198–209. <https://doi.org/10.5435/00124635-200205000-00006>.
- [8] Koffman M. Proximal femoral resection or total hip replacement in severely disabled cerebral-spastic patients. *Orthop Clin North Am* 1981;12:91–100. <https://doi.org/10.1002/bjs.18004519315>.
- [9] Wiles P. The surgery of the osteoarthritic hip. *Br J Surg* 1958;45:488–97. <https://doi.org/10.1002/bjs.18004519315>.
- [10] Ferguson RJ, Palmer AJ, Taylor A, Porter ML, Malchou H, Glyn-Jones S. Hip replacement. *Lancet* 2018;392:1662–71. [https://doi.org/10.1016/S0140-6736\(18\)31777-X](https://doi.org/10.1016/S0140-6736(18)31777-X).
- [11] Subramanian P, Wainwright TW, Bahadori S, Middleton RG. A review of the evolution of robotic-assisted total hip arthroplasty. *Hip Int* 2019;29:232–8. <https://doi.org/10.1177/1120700019828286>.
- [12] Singh JA, Lewallen DG. Increasing obesity and comorbidity in patients undergoing primary total hip arthroplasty in the U.S.: a 13-year study of time trends. *BMC Musculoskelet Disord* 2014;15:441. <https://doi.org/10.1186/1471-2474-15-441>.
- [13] Ravi B, Croxford R, Reichmann WM, Losina E, Katz JN, Hawker GA. The changing demographics of total joint arthroplasty recipients in the United States and Ontario from 2001 to 2007. *Best Pract Res Clin Rheumatol* 2012;26:637–47. <https://doi.org/10.1016/j.berh.2012.07.014>.
- [14] Kleiner JE, Eltorai AEM, Rubin LE, Daniels AH. Matched cohort analysis of total hip arthroplasty in patients with and without Parkinson's disease: complications, mortality, length of stay, and hospital charges. *J Arthroplasty* 2019;34:S228–31. <https://doi.org/10.1016/j.arth.2019.03.023>.
- [15] King G, Hunt LP, Wilkinson JM, Blom AW. National joint registry for England, Wales, and northern Ireland. Good outcome of total hip replacement in patients with cerebral palsy: a comparison of 389 patients and 425,813 controls from the national joint registry for England and Wales. *Acta Orthop* 2016;87:93–9. <https://doi.org/10.3109/17453674.2015.1137439>.
- [16] Newman JM, Naziri Q, Chughtai M, Khlopas A, Kryzak TJ, Navale SM, et al. Does multiple sclerosis affect the inpatient perioperative outcomes after total hip arthroplasty? *J Arthroplasty* 2017;32:3669–74. <https://doi.org/10.1016/j.arth.2017.07.006>.
- [17] Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009;339:b2700. <https://doi.org/10.1136/bmj.b2700>.
- [18] Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336:924–6. <https://doi.org/10.1136/bmj.39489.470347.AD>.
- [19] Howick J, Chalmers I, Glasziou P, Greenhalgh T, Heneghan C, Liberati A, et al. The 2011 Oxford CEBM levels of evidence. Oxford, UK: University of Oxford; 2011. p. 12. <https://www.cebm.ox.ac.uk/resources/levels-of-evidence/cebm-levels-of-evidence> [accessed 02.02.22].
- [20] Kalia LV, Lang AE. Parkinson's disease. *Lancet* 2015;386:896–912. [https://doi.org/10.1016/S0140-6736\(14\)61393-3](https://doi.org/10.1016/S0140-6736(14)61393-3).
- [21] Tynes OB, Storstein A. Epidemiology of Parkinson's disease. *J Neural Transm (Vienna)* 2017;124:901–5. <https://doi.org/10.1007/s00702-017-1686-y>.
- [22] Golbe LI, Leyton CE. Life expectancy in Parkinson disease. *Neurology* 2018;91:991–2. <https://doi.org/10.1212/WNL.0000000000006560>.
- [23] Newman JM, Sodhi N, Dalton SE, Khlopas A, Newman RP, Higuera CA, et al. Does Parkinson disease increase the risk of perioperative complications after total hip arthroplasty? A nationwide database study. *J Arthroplasty* 2018;33:S162–6. <https://doi.org/10.1016/j.arth.2018.01.006>.
- [24] Pepper PV, Goldstein MK. Postoperative complications in Parkinson's disease. *J Am Geriatr Soc* 1999;47:967–72. <https://doi.org/10.1111/j.1532-5415.1999.tb01292.x>.
- [25] Hoehn MM, Yahr MD. Parkinsonism: onset, progression and mortality. *Neurology* 1967;17:427–42. <https://doi.org/10.1212/wnl.17.5.427>.
- [26] Rondon AJ, Tan TL, Schlitt PK, Greenky MR, Phillips JL, Purtill JJ. Total joint arthroplasty in patients with Parkinson's disease: survivorship, outcomes, and reasons for failure. *J Arthroplasty* 2018;33:1028–32. <https://doi.org/10.1016/j.arth.2017.11.017>.
- [27] Shah NV, Solow M, Lavian JD, Bloom LR, Grieco PW, Stroud SG, et al. Total hip arthroplasty in Parkinson's disease patients: a propensity score-matched analysis with minimum 2-year surveillance. *Hip Int* 2020;30:684–9. <https://doi.org/10.1177/1120700019862247>.
- [28] Wojtowicz AL, Mohaddes M, Odin D, Bülow E, Nemes S, Cnudde P. Is Parkinson's disease associated with increased mortality, poorer outcomes scores, and revision risk after THA? Findings from the Swedish hip arthroplasty register. *Clin Orthop Relat Res* 2019;477:1347–55. <https://doi.org/10.1097/CORR.0000000000000679>.
- [29] Jämsen E, Puolakka T, Peltola M, Eskelinen A, Lehto MU. Surgical outcomes of primary hip and knee replacements in patients with Parkinson's disease: a nationwide registry-based case-controlled study. *Bone Joint J* 2014;96-B:486–91. <https://doi.org/10.1302/0301-620X.96B4.33422>.
- [30] Weber M, Cabanela ME, Sim FH, Frassica FJ, Harmsen WS. Total hip replacement in patients with Parkinson's disease. *Int Orthop* 2002;26:66–8. <https://doi.org/10.1007/s00264-001-0308-2>.
- [31] Rong X, Dahal S, Luo ZY, Zhou K, Yao SY, Zhou ZK. Functional outcomes after total joint arthroplasty are related to the severity of Parkinson's disease: a mid-term follow-up. *J Orthop Surg Res* 2019;14:396. <https://doi.org/10.1186/s13018-019-1447-8>.
- [32] Lazennec JY, Kim Y, Pour AE. Total hip arthroplasty in patients with Parkinson disease: improved outcomes with dual mobility implants and cementless fixation. *J Arthroplasty* 2018;33:1455–61. <https://doi.org/10.1016/j.arth.2017.11.062>.
- [33] Sponer P, Kučera T, Grinac M, Bezrouk A, Waciakowski D. The outcomes of total hip replacement in patients with Parkinson's disease: comparison of the elective and hip fracture groups. *Parkinsons Dis* 2017;2017:1597463. <https://doi.org/10.1155/2017/1597463>.
- [34] Rosenbaum P, Paneth N, Leviton A, Goldstein M, Bax M, Damiano D, et al. A report: the definition and classification of cerebral palsy April 2006. *Dev Med Child Neurol Suppl* 2007;109:8–14.
- [35] Mathewson MA, Li RL. Pathophysiology of muscle contractures in cerebral palsy. *Phys Med Rehabil Clin N Am* 2015;26:57–67. <https://doi.org/10.1016/j.pmr.2014.09.005>.
- [36] Adams CT, Lakra A. Clinical and functional outcomes of total hip arthroplasty in patients with cerebral palsy: a systematic review. *J Orthop* 2020;21:19–24. <https://doi.org/10.1016/j.jor.2020.01.031>.
- [37] Scrutton D, Baird G, Smeeton N. Hip dysplasia in bilateral cerebral palsy: incidence and natural history in children aged 18 months to 5 years. *Dev Med Child Neurol* 2001;43:586–600. <https://doi.org/10.1017/s0012162201001086>.
- [38] Moon AS, Pinto MC, Cichos KH, McGwin Jr G, Ponce BA, Ghanem ES. Total joint arthroplasty in patients with cerebral palsy. *J Am Acad Orthop Surg* 2020;28:171–7. <https://doi.org/10.5435/JAAOS-D-18-00828>.
- [39] Moore HG, Gardezi M, Burroughs PJ, Rubin LE, Frumberg DB, Grauer JN. Total hip arthroplasty in patients with cerebral palsy: a matched comparison of 90-day adverse events and 5-year implant survival. *J Arthroplasty* 2021;36:3534–7. <https://doi.org/10.1016/j.arth.2021.05.039>.
- [40] Houdek MT, Watts CD, Wyles CC, Trousdale RT, Milbrandt TA, Taunton MJ. Total hip arthroplasty in patients with cerebral palsy: a cohort study matched to patients with osteoarthritis. *J Bone Joint Surg Am* 2017;99:488–93. <https://doi.org/10.2106/JBJS.16.00528>.
- [41] Morin C, Ursu C, Delecourt C. Total hip replacement in young non-ambulatory cerebral palsy patients. *Orthop Traumatol Surg Res* 2016;102:845–9. <https://doi.org/10.1016/j.otsr.2016.07.010>.
- [42] Yoon BH, Lee YK, Ha YC, Koo KH. Contemporary ceramic total hip arthroplasty in patients with cerebral palsy: does it work? *Clin Orthop Surg* 2015;7:39–45. <https://doi.org/10.4055/cios.2015.7.1.39>.
- [43] Sanders RJ, Swierstra BA, Goosen JH. The use of a dual-mobility concept in total hip arthroplasty patients with spastic disorders: no dislocations in a series of ten cases at midterm follow-up. *Arch Orthop Trauma Surg* 2013;133:1011–6. <https://doi.org/10.1007/s00402-013-1759-9>.
- [44] Schroeder K, Hauck C, Wiedenhöfer B, Braatz F, Aldinger PR. Long-term results of hip arthroplasty in ambulatory patients with cerebral palsy. *Int Orthop* 2010;34:335–9. <https://doi.org/10.1007/s00264-009-0771-8>.
- [45] Raphael BS, Dines JS, Akerman M, Root L. Long-term follow up of total hip arthroplasty in patients with cerebral palsy. *Clin Orthop Relat Res* 2010;468:1845–54. <https://doi.org/10.1007/s11999-009-1167-1>.
- [46] Weber M, Cabanela ME. Total hip arthroplasty in patients with cerebral palsy. *Orthopedics* 1999;22:425–7. <https://doi.org/10.3928/0147-7447-19990401-12>.
- [47] Yamout BI, Alroughani R. Multiple sclerosis. *Semin Neurol* 2018;38:212–25. <https://doi.org/10.1055/s-0038-1649502>.
- [48] Tullman MJ. Overview of the epidemiology, diagnosis, and disease progression associated with multiple sclerosis. *Am J Manag Care* 2013;19(2 Suppl):S15–20.
- [49] Pidgeon TS, Borenstein T, Daniels AH, Murali J, Hayda RA. Understanding multiple sclerosis: essentials for the orthopaedic surgeon. *JBJS Rev* 2014;2:e3. <https://doi.org/10.2106/JBJS.RVW.M.00120>.
- [50] Newman JM, Khlopas A, Sodhi N, Curtis GL, Sultan AA, George J, et al. Are adverse outcome rates higher in multiple sclerosis patients after total hip arthroplasty? *Bone Joint J* 2018;100-B:875–81. <https://doi.org/10.1302/0301-620X.100B7.BJJ-2017-1569.R1>.
- [51] Quinlan ND, Chen DQ, Werner BC, Barnes CL, Browne JA. Patients with multiple sclerosis are at increased risk for postoperative complications following total hip and knee arthroplasty. *J Arthroplasty* 2019;34:1606–10. <https://doi.org/10.1016/j.arth.2019.04.022>.
- [52] Rondon AJ, Schlitt PK, Tan TL, Phillips JL, Greenky MR, Purtill JJ. Survivorship and outcomes in patients with multiple sclerosis undergoing total joint arthroplasty. *J Arthroplasty* 2018;33:1024–7. <https://doi.org/10.1016/j.arth.2017.10.056>.
- [53] Gutman JM, Kim K, Schwarzkopf R, Kister I. Total hip and knee arthroplasty in patients with multiple sclerosis. *Int J MS Care* 2018;20:244–50. <https://doi.org/10.7224/1537-2073.2017-093>.
- [54] Bennett JE, Dolin R, Blaser MJ. *Mandell, Douglas, and Bennett's principles and practice of infectious diseases*. 8th ed. Philadelphia, PA: Saunders; 2014.

- [55] Lau JH, Parker JC, Hsu LC, Leong JC. Paralytic hip instability in poliomyelitis. *J Bone Joint Surg Br* 1986;68:528–33. <https://doi.org/10.1302/0301-620X.68B4.3733824>.
- [56] Sonekatsu M, Sonohata M, Kitajima M, Kawano S, Mawatari M. Total hip arthroplasty for patients with residual poliomyelitis at a mean eight years of follow-up. *Acta Med Okayama* 2018;72:17–22. <https://doi.org/10.18926/AMO/55658>.
- [57] Sobrón FB, Martínez-Ayora Á, Cuervas-Mons M, Quevedo T, Laguna R, Vaquero J. Total hip arthroplasty in patients of post polio residual paralysis: a retrospective case series. *Indian J Orthop* 2017;51:434–9. <https://doi.org/10.4103/0019-5413.209951>.
- [58] Cho YJ, Lee CH, Chun YS, Rhyu KH. Outcome after cementless total hip arthroplasty for arthritic hip in patients with residual poliomyelitis: a case series. *Hip Int* 2016;26:458–61. <https://doi.org/10.5301/hipint.5000372>.
- [59] Yoon BH, Lee YK, Yoo JJ, Kim HJ, Koo KH. Total hip arthroplasty performed in patients with residual poliomyelitis: does it work? *Clin Orthop Relat Res* 2014;472:933–40. <https://doi.org/10.1007/s11999-013-3338-3>.
- [60] DeDeugd CM, Perry KI, Trousdale WH, Taunton MJ, Lewallen DG, Abdel MP. Total hip arthroplasty in patients affected by poliomyelitis. *Bone Joint J* 2018;100-B:733–9. <https://doi.org/10.1302/0301-620X.100B6.BJJ-2018-0127.R1>.
- [61] Zhuang TF, Huan SW, Luo SM, She GR, Wu WR, Chen JY, et al. Outcomes of dual mobility articulation total hip arthroplasty in ipsilateral residual poliomyelitis. *Int Orthop* 2022;46:489–96. <https://doi.org/10.1007/s00264-021-05222-y>.
- [62] Faldini C, De Fine M, Di Martino A, Fabbri D, Borghi R, Chehrassan M, et al. Outcomes of total hip replacement in limbs affected by poliomyelitis. *Hip Int* 2017 Mar 31;27:198–204. <https://doi.org/10.5301/hipint.5000451>.
- [63] Buttaro MA, Slullitel PA, García Mansilla AM, Carlucci S, Comba FM, Zanotti G, et al. Long-term outcome of unconstrained primary total hip arthroplasty in ipsilateral residual poliomyelitis. *Orthopedics* 2017;40:e255–61. <https://doi.org/10.3928/01477447-20161108-03>.
- [64] Hosalkar HS, Fuller DA, Rendon N, Esquezenai A, Keenan MAE. Outcomes of total joint arthroplasties in adults with post-polio syndrome: results from a tertiary neuro-orthopaedic center. *Curr Orthop Pract* 2010;21:273–81. <https://doi.org/10.1097/BCO.0b013e3181d2dc6c>.
- [65] Chalmers BP, Tibbo ME, Trousdale RT, Lewallen DG, Berry DJ, Abdel MP. Primary total hip arthroplasty for Charcot arthropathy is associated with high complications but improved clinical outcomes. *J Arthroplasty* 2018;33:2912–8. <https://doi.org/10.1016/j.arth.2018.04.002>.
- [66] Inoue D, Kabata T, Kajino Y, Taga T, Yamamoto T, Takagi T, et al. Clinical results of total hip arthroplasty in two patients with Charcot hip joints due to congenital insensitivity to pain with anhidrosis. *Case Rep Orthop* 2018;2018:1743068. <https://doi.org/10.1155/2018/1743068>.
- [67] Rapała K, Obrebski M. Charcot's arthropathy of the hip joints: a late manifestation of tabes dorsalis successfully treated by total joint arthroplasty. report of 2 cases. *J Arthroplasty* 2007;22:771–4. <https://doi.org/10.1016/j.arth.2006.05.032>.
- [68] Henawy AT, Abdel Badie A. Dual mobility total hip arthroplasty in hemiplegic patients. *SICOT J* 2017;3:40. <https://doi.org/10.1051/sicotj/2017024>.
- [69] Wang Y, Deng X, Wang Z, Zhu Y, Chen W, Zhang Y. Total hip arthroplasty or hemiarthroplasty for femoral neck fractures in elderly patients with neuromuscular imbalance. *Aging Clin Exp Res* 2021;34:2825–33. <https://doi.org/10.1007/s40520-021-01976-y>.
- [70] Ryu HG, Roh YJ, Oh KJ, Hwang JH, Kim Y, Cho HW, et al. Dual mobility articulation total hip arthroplasty for displaced neck fracture in elderly with neuromuscular disorder. *Injury* 2021;52:1480–6. <https://doi.org/10.1016/j.injury.2021.01.005>.
- [71] Blizzard DJ, Klement MR, Penrose CT, Sheets CZ, Bolognesi MP, Seyler TM. Cervical myelopathy doubles the rate of dislocation and fracture after total hip arthroplasty. *J Arthroplasty* 2016;31(9 Suppl):242–7. <https://doi.org/10.1016/j.arth.2016.05.070>.
- [72] Abdelazim H, Michael F. Dual mobility cup for prevention of early total hip arthroplasty dislocation in patients with neurological disorders. *Eur Orthop Traumatol* 2015;6:427–32. <https://doi.org/10.1007/s12570-015-0336-9>.
- [73] Li J, Zheng W, Zhao J, Liu D, Xu W. Large diameter metal on metal total hip replacement for femoral neck fractures with neurological conditions: a retrospective assessment. *Indian J Orthop* 2014;48:605–11. <https://doi.org/10.4103/0019-5413.144236>.
- [74] Alish H, Kamath AF, Baldwin KD, Keenan M, Lee GC. Outcomes of total hip arthroplasty in spastic patients. *J Arthroplasty* 2014;29:1566–70. <https://doi.org/10.1016/j.arth.2014.03.005>.
- [75] Park KS, Seon JK, Lee KB, Yoon TR. Total hip arthroplasty using large-diameter metal-on-metal articulation in patients with neuromuscular weakness. *J Arthroplasty* 2014;29:797–801. <https://doi.org/10.1016/j.arth.2013.08.012>.
- [76] Jolbäck P, Rolfson O, Cnudde P, Odin D, Malchau H, Lindahl H, et al. High annual surgeon volume reduces the risk of adverse events following primary total hip arthroplasty: a registry-based study of 12,100 cases in Western Sweden. *Acta Orthop* 2019;90:153–8. <https://doi.org/10.1080/17453674.2018.1554418>.
- [77] Ravi B, Jenkinson R, Austin PC, Croxford R, Wasserstein D, Escott B, et al. Relation between surgeon volume and risk of complications after total hip arthroplasty: propensity score matched cohort study. *BMJ* 2014;348:g3284. <https://doi.org/10.1136/bmj.g3284>.
- [78] Eibach S, Krug H, Lobsien E, Hoffmann KT, Kupsch A. Preoperative treatment with Botulinum Toxin A before total hip arthroplasty in a patient with tetraspasticity: case report and review of literature. *NeuroRehabilitation* 2011;28:81–3. <https://doi.org/10.3233/NRE-2011-0635>.
- [79] Mehta S, Vankleunen JP, Booth RE, Lotke PA, Lonner JH. Total knee arthroplasty in patients with Parkinson's disease: impact of early postoperative neurologic intervention. *Am J Orthop (Belle Mead NJ)* 2008;37:513–6.
- [80] Wilkin GP, Ibrahim MM, Smit KM, Beaulé PE. A contemporary definition of hip dysplasia and structural instability: toward a comprehensive classification for acetabular dysplasia. *J Arthroplasty* 2017;32:S20–7. <https://doi.org/10.1016/j.arth.2017.02.067>.
- [81] Shrader MW, Wimberly L, Thompson R. Hip surveillance in children with cerebral palsy. *J Am Acad Orthop Surg* 2019;27:760–8. <https://doi.org/10.5435/JAOS-D-18-00184>.
- [82] Ueoka K, Kabata T, Kajino Y, Yoshitani J, Ueno T, Tsuchiya H. The accuracy of the computed tomography-based navigation system in total hip arthroplasty is comparable with crowe type IV and crowe type I dysplasia: a case-control study. *J Arthroplasty* 2019;34:2686–91. <https://doi.org/10.1016/j.arth.2019.06.002>.
- [83] Lübbecke A, Suvá D, Perneger T, Hoffmeyer P. Influence of preoperative patient education on the risk of dislocation after primary total hip arthroplasty. *Arthritis Rheum* 2009;61:552–8. <https://doi.org/10.1002/art.24340>.
- [84] Zuckerman LM. Parkinson's disease and the orthopaedic patient. *J Am Acad Orthop Surg* 2009;17:48–55. <https://doi.org/10.5435/00124635-200901000-00007>.
- [85] Rezapoor M, Parvizi J. Prevention of periprosthetic joint infection. *J Arthroplasty* 2015;30:902–7. <https://doi.org/10.1016/j.arth.2015.02.044>.
- [86] Huang RC, Malkani AL, Harwin SF, Hozack WJ, Mont MA, Higuera-Rueda CA, et al. Multicenter evaluation of a modular dual mobility construct for revision total hip arthroplasty. *J Arthroplasty* 2019;34:S287–91. <https://doi.org/10.1016/j.arth.2019.03.027>.
- [87] Jones SA. Constrained acetabular liners. *J Arthroplasty* 2018;33:1331–6. <https://doi.org/10.1016/j.arth.2018.01.026>.
- [88] Rowan FE, Benjamin B, Pietrak JR, Haddad FS. Prevention of dislocation after total hip arthroplasty. *J Arthroplasty* 2018;33:1316–24. <https://doi.org/10.1016/j.arth.2018.01.047>.
- [89] Jameson SS, Lees D, James P, Serrano-Pedraza I, Partington PF, Muller SD, et al. Lower rates of dislocation with increased femoral head size after primary total hip replacement: a five-year analysis of NHS patients in England. *J Bone Joint Surg Br* 2011;93:876–80. <https://doi.org/10.1302/0301-620X.93B7.26657>.
- [90] White Jr RE, Forness TJ, Allman JK, Junick DW. Effect of posterior capsular repair on early dislocation in primary total hip replacement. *Clin Orthop Relat Res* 2001;393:163–7. <https://doi.org/10.1097/00003086-200112000-00019>.

Medline/PubMed search strategy.

#1 THR population	"Total hip arthroplasty" OR THA OR "total hip replacement" OR THR OR "arthroplasty, replacement, Hip"[Mesh]
#2 Neurological Population	"cerebral palsy" OR "spina bifida" OR myelomeningocele OR poliomyelitis OR "Parkinson disease" OR "multiple sclerosis" OR stroke OR CVA OR "acquired brain injury" OR Charcot OR "neuropathic arthropathy" OR "neuromuscular disease" OR "spinal injury" OR "paralytic hip" OR "Cerebral Palsy"[Mesh] OR "Spinal Dysraphism"[Mesh] OR "Meningomyelocele"[Mesh] OR "poliomyelitis"[Mesh] OR "Parkinson Disease"[Mesh] OR "Multiple Sclerosis"[Mesh] OR "Stroke"[Mesh] OR "Brain Injuries"[Mesh] OR "Arthropathy, Neurogenic"[Mesh] OR "Neuromuscular Diseases"[Mesh] OR "Spinal Injuries"[Mesh]
#3 Outcome	outcome OR "clinical outcome" OR "patient outcome" OR revision OR mortality OR death OR infection OR complication OR dislocation OR "patient reported outcome measure" OR PROM OR "Treatment Outcome"[Mesh] OR "Patient Reported Outcome Measures"[Mesh] OR "Reoperation"[Mesh] OR "Mortality"[Mesh] OR "Death"[Mesh] OR "Infections"[Mesh] OR "Intraoperative Complications"[Mesh] OR "Postoperative Complications"[Mesh] OR "Joint Dislocations"[Mesh]

Cochrane library search strategy

#1 THR population	"Total hip arthroplasty" OR THA OR "total hip replacement" OR THR OR MeSH descriptor: [Arthroplasty, replacement, hip] explode all trees
#2 Neurological Population	"cerebral palsy" OR "spina bifida" OR myelomeningocele OR poliomyelitis OR "Parkinson disease" OR "multiple sclerosis" OR stroke OR CVA OR "acquired brain injury" OR Charcot OR "neuropathic arthropathy" OR "neuromuscular disease" OR "spinal injury" OR "paralytic hip" OR MeSH descriptor: [Cerebral Palsy] explode all trees OR MeSH descriptor: [Spinal Dysraphism] explode all trees OR MeSH descriptor: [Meningomyelocele] explode all trees OR MeSH descriptor: [Poliomyelitis] explode all trees OR MeSH descriptor: [Parkinson Disease] explode all trees OR MeSH descriptor: [Multiple Sclerosis] explode all trees OR MeSH descriptor: [Stroke] explode all trees OR MeSH descriptor: [Arthropathy, Neurogenic] explode all trees OR MeSH descriptor: [Neuromuscular Disease] explode all trees OR MeSH descriptor: [Spinal Injuries] explode all trees
#3 Outcome	outcome OR "clinical outcome" OR "patient outcome" OR revision OR mortality OR death OR infection OR complication OR dislocation OR "patient reported outcome measure" OR PROM OR MeSH descriptor: [Patient Outcome Assessment] explode all trees OR MeSH descriptor: [Reoperation] explode all trees OR MeSH descriptor: [Mortality] explode all trees OR MeSH descriptor: [Death] explode all trees OR MeSH descriptor: [Infections] explode all trees OR MeSH descriptor: [Intraoperative Complications] explode all trees OR MeSH descriptor: [Postoperative Complications] explode all trees OR MeSH descriptor: [Hip Dislocation] explode all trees

Embase search strategy

#1 THR population	"Total hip arthroplasty" OR THA OR "total hip replacement" OR THR OR 'total hip replacement'/exp
#2 Neurological Population	"cerebral palsy" OR "spina bifida" OR myelomeningocele OR poliomyelitis OR "Parkinson disease" OR "multiple sclerosis" OR stroke OR CVA OR "acquired brain injury" OR Charcot OR "neuropathic arthropathy" OR "neuromuscular disease" OR "spinal injury" OR "paralytic hip" OR 'cerebral palsy'/exp OR 'spinal dysraphism'/exp OR 'meningomyelocele'/exp OR 'poliomyelitis'/exp OR 'Parkinson disease'/exp OR 'multiple sclerosis'/exp OR 'cerebrovascular accident'/exp OR 'acquired brain injury'/exp OR 'neuropathic joint disease'/exp OR 'neuromuscular disease'/exp OR 'spine injury'/exp
#3 Outcome	outcome OR "clinical outcome" OR "patient outcome" OR revision OR mortality OR death OR infection OR complication OR dislocation OR "patient reported outcome measure" OR PROM OR 'treatment outcome'/exp OR 'clinical outcome'/exp OR 'revision arthroplasty'/exp OR 'mortality'/exp OR 'death'/exp OR 'infection'/exp OR 'postoperative complication'/exp OR 'perioperative complication'/exp OR 'hip dislocation'/exp