

9-18-2023

Lateral Approach to the Lumbar Spine: The Utility of an Access Surgeon

Matthew Meade
Thomas Jefferson University

Yunsoo Lee
Thomas Jefferson University

Parker Brush
Thomas Jefferson University

Mark Lambrechts
Thomas Jefferson University

Eleanor Jenkins
Thomas Jefferson University
Follow this and additional works at: <https://jdc.jefferson.edu/orthofp>



Part of the [Orthopedics Commons](#), and the [Surgery Commons](#)

See next page for additional authors

[Let us know how access to this document benefits you](#)

Recommended Citation

Meade, Matthew; Lee, Yunsoo; Brush, Parker; Lambrechts, Mark; Jenkins, Eleanor; Desimone, Cristian; Mccurdy, Michael; Mangan, John; Canseco, Jose; Kurd, Mark; Hilibrand, Alan; Vaccaro, Alexander; Kepler, Christopher; and Schroeder, Gregory, "Lateral Approach to the Lumbar Spine: The Utility of an Access Surgeon" (2023). *Department of Orthopaedic Surgery Faculty Papers*. Paper 208.
<https://jdc.jefferson.edu/orthofp/208>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Orthopaedic Surgery Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Matthew Meade, Yunsoo Lee, Parker Brush, Mark Lambrechts, Eleanor Jenkins, Cristian Desimone, Michael Mccurdy, John Mangan, Jose Canseco, Mark Kurd, Alan Hilibrand, Alexander Vaccaro, Christopher Kepler, and Gregory Schroeder

Lateral approach to the lumbar spine: The utility of an access surgeon

ABSTRACT

Background: Lateral lumbar interbody fusions (LLIFs) utilize a retroperitoneal approach that avoids the intraperitoneal organs and manipulation of the anterior vasculature encountered in anterior approaches to the lumbar spine. The approach was championed by spinal surgeons; however, general/vascular surgeons may be more comfortable with the approach.

Objective: The objective of this study was to compare short-term outcomes following LLIF procedures based on whether a spine surgeon or access surgeon performed the approach.

Materials and Methods: We retrospectively identified all one- to two-level LLIFs at a tertiary care center from 2011 to 2021 for degenerative spine disease. Patients were divided into groups based on whether a spine surgeon or general surgeon performed the surgical approach. The electronic medical record was reviewed for hospital readmissions and complication rates.

Results: We identified 239 patients; of which 177 had approaches performed by spine surgeons and 62 by general surgeons. The spine surgeon group had fewer levels with posterior instrumentation (1.40 vs. 2.00; $P < 0.001$) and decompressed (0.94 vs. 1.25, $P = 0.046$); however, the two groups had a similar amount of two-level LLIFs (29.9% vs. 27.4%, $P = 0.831$). This spine surgeon approach group was found to have shorter surgeries (281 vs. 328 min, $P = 0.002$) and shorter hospital stays Length of Stay (LOS) (3.1 vs. 3.6 days, $P = 0.019$); however, these differences were largely attributed to the shorter posterior fusion construct. On regression analysis, there was no statistical difference in postoperative complication rates whether or not an access surgeon was utilized ($P = 0.226$).

Conclusion: Similar outcomes may be seen regardless of whether a spine or access surgeon performs the approach for an LLIF.

Keywords: Approach, complications, lateral lumbar interbody fusion, minimally invasive, outcomes

INTRODUCTION

With recent advances in lumbar fusion techniques, there is an increasing need for safe surgical access to the lumbar spine. Commonly, vascular or general surgeons have been involved as “access surgeons” for anterior approaches due to their expertise in the peritoneal space.^[1-3] Access surgeons are most often utilized for a direct anterior approach (i.e., anterior lumbar interbody fusions [ALIFs]), in which there is a significant risk with mobilization of the great vessels as well as nonvascular complications like direct injury to the bowel or genitourinary structures.^[3-8] Multiple studies have evaluated the incidence and types of complications in ALIF with and without the use of access surgeons.^[4,9-12] Overall, there is varied support in the literature for general/vascular assistance

MATTHEW H. MEADE, YUNSOO LEE¹, PARKER L. BRUSH¹, MARK J. LAMBRECHTS¹, ELEANOR H. JENKINS¹, CRISTIAN A. DESIMONE¹, MICHAEL A. MCCURDY¹, JOHN J. MANGAN¹, JOSE A. CANSECO¹, MARK F. KURD¹, ALAN S. HILIBRAND¹, ALEXANDER R. VACCARO¹, CHRISTOPHER K. KEPLER¹, GREGORY D. SCHROEDER¹

Department of Orthopaedic Surgery, Jefferson Health – New Jersey, Washington Township, NJ, ¹Department of Orthopaedic Surgery, Rothman Orthopaedic Institute at Thomas Jefferson University, Philadelphia, PA, USA

Address for correspondence: Dr. Yunsoo Lee, 925 Chestnut, Philadelphia, PA 19107, USA. E-mail: yunsoo.a.lee@gmail.com

Submitted: 17-Jul-23


Accepted: 05-Aug-23

Published: 18-Sep-23

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Meade MH, Lee Y, Brush PL, Lambrechts MJ, Jenkins EH, Desimone CA, *et al.* Lateral approach to the lumbar spine: The utility of an access surgeon. *J Craniovert Jun Spine* 2023;14:281-7.

Access this article online	
Website: www.jcvjs.com	Quick Response Code 
DOI: 10.4103/jcvjs.jcvjs_78_23	

with anterior-based approaches to the spine.^[4,10,11,13-15] Other approaches to the spine utilize access surgeons less frequently but are not without their own risks. One such approach is the lateral lumbar interbody fusion (LLIF).

In contrast to the anterior approach where the spine is accessed by moving the intraperitoneal organs and iliac veins to gain access to the anterior lumbar spine, the lateral approach utilizes an entirely retroperitoneal approach and, thus, is associated with a decreased risk of vascular injury.^[16-19] The approach was described and published by Ozgur *et al.* in the early 2000s as a less-invasive alternative to the conventional ALIF and was credited as a novel approach that could be safely performed without a general surgeon for access.^[17,20] It allowed for interbody stabilization with indirect decompression while avoiding the major visceral/vascular risks encountered in the anterior approach to the spine.^[21,22] In addition, compared to the posterior approach, LLIF offers a more extensive discectomy while enabling the preparation of a larger surface area within the disc space for intervertebral graft positioning and avoiding damage to the posterior structures.^[23,24] The major structures at risk while performing an LLIF are neurologic, including the ilioinguinal and lateral femoral cutaneous nerves, as well as the femoral nerve, genitofemoral nerve, and lumbar plexus itself within the psoas musculature. This includes an estimated 4.8% risk of motor femoral nerve palsy after L4-L5 LLIFs.^[18,25-30] Given the decreased risk to vascular and abdominal structures through the lateral approach, access surgeon utilization is less commonly employed.

In contrast to the anterior approach to the spine, there is sparse literature regarding access surgeon utilization and associated outcomes for LLIFs. The current study aimed to address these shortcomings by primarily evaluating postoperative complications and readmissions for patients undergoing an LLIF with and without the utilization of an access surgeon. Secondly, we aim to compare in-hospital opioid use, length of hospital stay (LOS), and surgical characteristics between groups.

MATERIALS AND METHODS

Following the Institutional Review Board Approval, we retrospectively conducted a structured query to identify all patients over 18 years of age who underwent an LLIF over 10 years from 2011 to 2021 by the current procedural terminology code 22558. All patients underwent surgical intervention at a single, tertiary urban academic center. Inclusion criteria consisted of those patients undergoing a single- or two-level LLIF. Patients who underwent surgery for nondegenerative pathologies, including trauma, infection,

and malignancy were excluded. Additional exclusion criteria consisted of patients who received a concomitant interbody fusion through a different approach, prior anterior interbody fusion, and a >4 level concomitant posterior decompression or fusion. The L1-2 level was excluded due to the low sample number and anatomic proximity to the diaphragm. Patients were subsequently stratified into those who underwent LLIF with the utilization of an access surgeon and those who had a spine surgeon perform the entire surgery. Demographic data and patient characteristics were obtained and analyzed through chart review consisting of patient age, sex, race, ethnicity, body mass index (BMI), Charlson Comorbidity Index, and medical comorbidities like diabetes. Surgical information gathered included procedure (LLIF), anatomic location, operative duration, and utilization of access surgeon. Operative duration was calculated as the time in minutes from incision to closure. Additional surgical information analyzed included reported surgical blood loss in mL, utilization of intraoperative blood salvage (cell saver), and need for postoperative blood transfusion. Notably, all LLIFs performed at our institution regardless of the surgeon are performed initially in the lateral decubitus position for interbody cage insertion. Subsequently, the patient is placed prone for posterior fixation.

Cohorts were primarily analyzed for differences in 90-day complications, additional surgeries, and hospital readmissions. Postoperative data were gathered through chart review and encompassed 90 days postoperatively as defined by the Centers for Medicare and Medicaid Services for major procedures. Complications were recorded if they required medical/surgical intervention (Clavien–Dindo grade III and IV) and/or were approach related. Secondly, patients were evaluated for the duration of surgery, LOS, and in-hospital morphine milliequivalents (MMEs) utilized postoperatively.

Data were compared using the independent *t*-tests or Mann–Whitney tests for continuous variables and Chi-squared or Fisher's exact tests for categorical variables. Multivariable regression analysis was performed to identify factors independently associated with postoperative complications, LOS, operative duration, and estimated blood loss (EBL) after an LLIF. $P < 0.05$ was deemed significant, and all statistical analyses were done using RStudio (version 4.1.2, Vienna, Austria).

RESULTS

Demographic information

In total, 1536 patients were identified by our query, and 239 unique patients undergoing LLIF were identified following

operative note review with the remainder of patients having undergone an ALIF. One hundred and seventy-seven (74.1%) patients underwent an LLIF without the utilization of an access surgeon, while the remaining 62 (25.9%) patients underwent an LLIF with an access surgeon. Patients who underwent LLIF with the orthopedic surgeon alone were significantly older compared to those in which an access surgeon was involved (63.9 ± 9.03 vs. 60.5 ± 10.7 , $P = 0.016$). There were no significant differences in medical comorbidities such as BMI (30.4 ± 6.07 vs. 30.1 ± 6.41 , $P = 0.658$), Charlson Comorbidity Index (2.77 ± 1.81 vs. 2.37 ± 1.45 , $P = 0.191$), or presence of diabetes (25.4% vs. 16.1%, $P = 0.186$) between the two groups [Table 1]. In addition, there was no significant difference in other demographic characteristics such as sex, race, or ethnicity between the two groups.

Table 1: Demographics

	Spine surgeon (n=177), n (%)	Access surgeon (n=62), n (%)	P
Age	63.9 (9.93)	60.5 (10.7)	0.016
Sex			
Female	91 (51.4)	37 (59.7)	0.330
Male	86 (48.6)	25 (40.3)	
Race			
White	149 (84.2)	56 (90.3)	0.088
Black	21 (11.9)	2 (3.23)	
Other	7 (3.95)	4 (6.45)	
Ethnicity			
Non-Hispanic	173 (97.7)	59 (95.2)	0.380
Hispanic	4 (2.26)	3 (4.84)	
BMI	30.4 (6.07)	30.1 (6.41)	0.658
CCI	2.77 (1.81)	2.37 (1.45)	0.191
Diabetes			
No	132 (74.6)	52 (83.9)	0.186
Yes	45 (25.4)	10 (16.1)	

Bolded text signifies a significant P (<0.05). BMI - Body mass index; CCI - Charlson Comorbidity Index

Table 2: Surgical information

	Spine surgeon (n=177), n (%)	Access surgeon (n=62), n (%)	P
Number of levels of LLIF			
1	124 (70.1)	45 (72.6)	0.831
2	53 (29.9)	17 (27.4)	
Number of levels fused posteriorly	1.40 (0.97)	2.00 (1.16)	<0.001
Number of levels decompressed	0.94 (1.05)	1.24 (1.13)	0.046
Operative duration (min)	281 (110)	328 (101)	0.002
EBL (mL)	285 (358)	456 (484)	<0.001
Transfusion			
No	111 (96.5)	42 (91.3)	0.227
Yes	4 (3.48)	4 (8.70)	
Cell saver			
No	91 (79.1)	20 (43.5)	<0.001
Yes	24 (20.9)	26 (56.5)	

Bolded text signifies a significant P (<0.05). LLIF - Lateral lumbar interbody fusion; EBL - Estimated blood loss

Surgical data

There were no significant differences in the number of levels fused with lateral interbody cages ($P = 0.831$) between the two cohorts [Table 2]. However, there was a significant difference in the number of levels decompressed and fused posteriorly with the spine surgeon group averaging less levels decompressed (0.94 ± 1.05 vs. 1.24 ± 1.13 , $P = 0.046$) and fused (1.40 ± 0.97 vs. 2.00 ± 1.16 , $P < 0.001$). In addition, the operative duration was significantly longer in the access surgeon group by an average of 47 min (328 ± 101 vs. 281 ± 110 , $P = 0.002$). EBL was significantly higher in operative cases where an access surgeon was utilized (456 ± 484 mL vs. 285 ± 358 mL, $P < 0.001$). This did not result in a significant difference in the need for postoperative transfusion (8.70% vs. 3.48%, $P = 0.227$); however, there was an increased utilization of intraoperative cell salvage in patients in the access surgeon group (56.5% vs. 20.9%, $P < 0.001$).

Inhospital course

The length of hospital stay was significantly different between the cohorts with the access surgeon cohort having a longer LOS (3.59 ± 1.62 days vs. 3.13 ± 1.92 days, $P = 0.019$). Postoperative in-hospital narcotic consumption was lower in patients who had a spine surgeon perform the entire procedure (238 ± 183 MME vs. 334 ± 197 MME, $P = 0.001$). However, this difference was not significant when adjusted for LOS (81.3 MME/day vs. 99.1 MME/day, $P = 0.157$) [Table 3].

Postoperative data

During the 90-day postoperative period, there were no differences in patients requiring additional surgery (2.26% vs. 4.84%, $P = 0.380$) or readmissions among the two groups (2.82% vs. 4.84% $P = 0.431$). Additional surgeries included four incision and drainage procedures, one revision decompression, one interbody removal, and one additional posterior fusion for spondylolisthesis. The

spine surgeon approach group had a lower postoperative complication rate (5.08% vs. 12.9%, $P = 0.048$). Postoperative complications included one death, three infections, seven visits to the emergency department (six for pain control and one for postoperative hypotension), two acute kidney injuries, one interbody dislodgement, one ileus, and one deep venous thrombosis. One patient experienced both an acute kidney injury and a deep venous thrombosis postoperatively [Table 4].

Multivariate regression analysis

The presence of an access surgeon was not associated with postoperative complications (odds ratio [OR]: 1.97; confidence interval [CI]: 0.64–5.92; $P = 0.226$), LOS (effect size: 0.25, CI: -0.26–0.75, $P = 0.338$), operative duration (effect size: 14, CI: -16–45, $P = 0.362$), or EBL (effect size: 40, CI: -76–156, $P = 0.503$). Patients with longer fusion constructs posteriorly were independently more likely to have a postoperative complication (OR = 1.92; CI = 1.18–3.15; $P = 0.008$), increased LOS (effect size: 0.39, CI: 0.16–0.62,

$P = 0.001$), increased operative duration (effect size: 58, CI: 44–76, $P < 0.001$), and more EBL (effect size: 206, CI: 154–257, $P < 0.001$). The number of levels decompressed posteriorly was also independently associated with increased operative duration (effect size: 25, CI: 12–38, $P < 0.001$) and increased EBL (effect size: 51, CI: 2–100, $P = 0.043$), but not postoperative complications ($P = 0.543$) or LOS ($P = 0.296$) [Table 5].

DISCUSSION

Previous literature regarding the utility of access surgeons has focused on the anterior approach to the lumbar spine. The risk profile of LLIF is significantly different from other lumbar spine surgical approaches and as such, understanding the utility of a spinal access surgeon cannot be gleaned from the data on other approaches to the anterior lumbar spine. We report in 239 patients that the use of an access surgeon does not impact inhospital or postoperative outcomes for patients undergoing LLIFs.

Although the lateral approach to the lumbar spine has demonstrated certain benefits compared to the traditional posterior approach, it also comes with its own set of unique risks. Studies have suggested that the lateral approach to the lumbar spine for fusion can shorten the LOS, reduce intraoperative blood loss, and decrease the risk of vascular/nerve injury.^[17,23,31,32] However, a significant risk with the lateral approach is the potential morbidity associated with dissection about the psoas musculature that can result in postoperative hip flexion pain or femoral nerve palsy.^[30,33,34] More specifically, when utilizing the lateral approach, there is an increased risk to vital neurologic structures, including the lumbar plexus and lumbar nerve roots.^[35] As the surgeon moves from cranial to caudal (i.e., L1-2 to L4-5), the psoas musculature and the intimately related lumbar plexus follow a more ventral trajectory posing an increasingly greater risk of injury.^[30] To mitigate the risk of lumbar plexus injury, intraoperative neuromonitoring is commonly utilized. Despite this, the rates of sensory deficit in the form of anterior thigh numbness and motor weakness (decreased hip flexion strength) are reported in 9.6% and 3.2% of patients, respectively, after an 18-month follow-up from LLIF.^[36]

Table 3: Inhospital course

	Spine surgeon (n=177)	Access surgeon (n=62)	P
Length of stay	3.13 (1.92)	3.59 (1.62)	0.019
MME in hospital	238 (183)	334 (197)	0.001
MME/day	81.3 (45.1)	99.1 (60.6)	0.157

Bolded text signifies a significant P (<0.05). MME - Morphine milliequivalent

Table 4: 90-day postoperative complications

	Spine surgeon (n=177), n (%)	Access surgeon (n=62), n (%)	P
Additional surgery			
No	173 (97.7)	59 (95.2)	0.380
Yes	4 (2.26)	3 (4.84)	
Chest tube			
No	177 (100)	61 (98.4)	0.259
Yes	0	1 (1.61)	
Readmission			
No	172 (97.2)	59 (95.2)	0.431
Yes	5 (2.82)	3 (4.84)	
Complication			
No	168 (94.9)	54 (87.1)	0.048
Yes	9 (5.08)	8 (12.9)	

Bolded text signifies a significant P (<0.05)

Table 5: Multivariate regression analysis

Variable	Postoperative complications			Length of stay			Operative duration			EBL		
	OR	CI	P	Effect size	CI	P	Effect size	CI	P	Effect size	CI	P
Access surgeon	1.97	0.64–5.92	0.226	0.25	-0.26–0.76	0.338	14	-16–45	0.362	40	-76–156	0.503
Levels fused posteriorly	1.92	1.18–3.15	0.008	0.39	0.16–0.62	0.001	58	44–76	<0.001	206	154–257	<0.001
Levels decompressed posteriorly	1.16	0.70–1.86	0.543	0.11	-0.10–0.33	0.296	25	12–38	<0.001	51	2–100	0.043

Length of stay is represented by days, length of surgery is represented by minutes, and EBL is represented by mm. Bolded text signifies a significant P (<0.05). OR - Odds ratio; CI - Confidence interval; EBL - Estimated blood loss

In addition to anterior thigh numbness and decreased hip flexion strength, the lateral approach can also affect other vital abdominal structures. One of the major goals in the development of the lateral approach to the lumbar spine was to mitigate risk to vascular structures that are commonly encountered during the anterior approach to the lumbar spine. A recent meta-analysis demonstrated the rates of aortic injury and common iliac lacerations during LLIF to be 0.12% and 0.25%, respectively.^[37] This contrasts the published rates of vascular injury associated with ALIF, which range from 1.9% to 24%.^[2,5,7] The decreased rate of vascular injury reported in LLIF can be partially attributed to the lateral positioning utilized, which results in the critical vascular structures moving away from the surgical corridor.^[38] Despite the overall decreased risk of vascular injury reported during LLIF, if it does occur, repairing it may be difficult due to the patient's lateral decubitus position.^[38-41] Other approach-related complications associated with the lateral approach include bowel injuries, ureteral injuries, and seromas.^[21] Although the retroperitoneal anatomy is not one that spine surgeons often encounter, we found no difference in approach-related complications, such as the instances of bowel, vascular, or ureteral injury. Thus, the addition of an access surgeon does not appear to change outcomes when performing the retroperitoneal approach of an LLIF.

However, like most novel orthopedic procedures, there is a learning curve to perform LLIF that must be considered. In one study, it was shown that complications such as iliopsoas and quadriceps weakness decrease as surgeons obtain more experience with the surgery.^[42] Other studies have echoed this sentiment stating that disadvantages of the LLIF include the learning curve associated with new surgical techniques and the involvement of a regional anatomy unfamiliar to most spine surgeons.^[30] Multiple recent studies have shown a temporally related decrease in postoperative neurologic complications with increased time performing LLIFs. One such study demonstrated a proportional trend for decreasing anterior thigh pain over 6 years.^[43] Similarly, Le *et al.* observed a learning curve with a reduction in the incidence of postoperative thigh numbness during 3 years of performing LLIF (from 26.1% to 10.7%).^[44] However, given that the approach has been utilized for over 20 years and there has been an exponential rise in its utilization and incorporation into training programs, spine surgeon exposure and level of comfort with this approach are likely improving.

Several of the secondary outcomes in this study were found to be significantly different between the two groups; however, these were likely due to confounding factors. The operative duration in the access surgeon group was nearly

one hour longer with a statistically significant difference of 47 min/case (281.0 vs. 328.0 min, $P = 0.002$). This difference is likely attributed to larger surgeries performed in the access surgeon group, and this hypothesis is supported by our regression analysis suggesting that each additional level fused and decompressed posteriorly was associated with an increase in the procedure length by over one hour. Importantly, complications after surgical intervention of the spine directly correlate with operative duration. Regarding LLIFs, Bendersky *et al.* previously recommended that psoas retraction be limited to 20–30 min per operative level.^[23,29] Specifically, they noted that a retraction time less than this decreased postoperative neurologic complications. Ultimately, the effect an access surgeon has on operative duration has yet to be fully elucidated in the literature; however, our study suggests that their impact is negligible for LLIFs.^[45] Similarly, EBL was shown to be significantly greater with an access surgeon (285 mL vs. 456 mL, $P < 0.001$); however, the most likely reason for this difference is in the greater number of levels fused and decompressed posteriorly in the access surgeon cohort as seen on multiple regression. Another explanation for this significant difference is that “cell saver” or intraoperative cell salvage was more likely to be used in cases with an access surgeon. Therefore, there were more objective data to estimate intraoperative blood loss compared to operative cases where a cell saver was not used. However, most importantly, despite the greater reported blood loss in the access surgeon cohort, this did not result in a significant difference in transfusion rates between the two groups.

Although we found that postoperative LOS may be longer in the access surgeon group, this may also be due to confounding factors. Postoperative LOS was significantly longer by 0.5 days in the access surgeon group (3.1 days vs. 3.6 days; $P = 0.019$). However, the utilization of an access surgeon was not independently associated with increased LOS on regression analysis. Our LOS is consistent with other studies, which report similar lengths of hospital stay following an LLIF (3.16 days for inpatient cases and 3.3 days for cases that include a posterior spinal fusion).^[46,47] Moreover, although we found that patients with access surgeons consumed a greater number of MMEs (238 vs. 334 MME; $P = 0.001$) during their hospital stay, when examining the MME/day, the amounts were not statistically different (81.3 vs. 99.1 MME/day; $P = 0.157$).

This study is not without limitations. First, it is retrospective in nature and has inherent weaknesses due to this method of evaluation with incomplete/incorrect documentation and unknown selection bias in potentially missed patients. In

addition, this study incorporated several spine surgeons with extensive experience performing LLIF. All spine surgeons performing these operations had undergone training in the lateral approach and completed a spinal surgery fellowship. As such, this cohort may not be generalizable to a cohort of surgeons with less experience. In addition, all spinal surgeons at our institution perform the lateral approach in the lateral decubitus position and then transition to a prone position for posterior instrumentation. The results of this study may not be generalizable to surgeons who do not transition to a prone position for the latter portion of the surgical intervention. Moreover, the differences in the baseline and surgical characteristics between the two groups confound the results on short-term outcomes. We attempt to control for this difference with a multivariate regression analysis; however, the size of the cohort limits the number of variables that can be controlled for through this analysis.

CONCLUSION

To date, there is limited evaluation on whether an access surgeon is beneficial when performing a lateral lumbar approach to the spine for LLIF. Our study demonstrates that similar results may be obtained with and without utilizing an access surgeon with the lateral approach to the lumbar spine for LLIF procedures. This study supports the notion that lateral access surgery to the lumbar spine can be performed safely by spinal surgeons and access surgeons. Further study is needed to understand the utility of access surgeons in revision cases and its potential benefit.

Financial support and sponsorship

Nil.

Conflicts of interest

MHM – None; YL – None; PB – None; ML - BMC Musculoskeletal Disorders: Editorial or governing board, Wolters Kluwer Health - Lippincott Williams and Wilkins: Editorial or governing board; EJ – None; CD – None; MAM – None; North American Spine Society: Board or committee member; JC - Accelus: Research support, Cervical Spine Research Society: Board or committee member, PathKeeper Surgical: Stock or stock Options; Unpaid consultant, Wolters Kluwer Health - Lippincott Williams and Wilkins: Editorial or governing board; MK - Camber Spine: Paid consultant, DuraStat LLC: Stock or stock Options, K2M Spine: Paid consultant, Spinal Elements: IP royalties; Paid consultant, Stryker: IP royalties; AH - Biomet: IP royalties, CTLAmedica: IP royalties, Paradigm spine: Stock or stock Options; AV - Accellus: Other financial or material support; Stock or stock Options, Advanced Spinal Intellectual Properties: Stock or stock Options, AO Spine: Other financial or material support, Atlas Spine: IP

royalties; Stock or stock Options, Avaz Surgical: Stock or stock Options, AVKN Patient Driven Care: Stock or stock Options, Cytonics: Stock or stock Options, Deep Health: Stock or stock Options, Dimension Orthotics LLC: Stock or stock Options, Electrocore: Stock or stock Options, Elsevier: Publishing royalties, financial or material support, Flagship Surgical: Stock or stock Options, FlowPharma: Stock or stock Options, Globus Medical: IP royalties; Stock or stock Options, Harvard Medtech: Stock or stock Options, Innovative Surgical Design: Stock or stock Options, Jaypee: Publishing royalties, financial or material support, Jushi (Haywood): Stock or stock Options, Medtronic: IP royalties, National Spine Health Foundation: Board or committee member, Nuvasive: Stock or stock Options, Orthobullets: Stock or stock Options, Parvizi Surgical Innovation: Stock or stock Options, Progressive Spinal Technologies: Stock or stock Options, Sentryx: Other financial or material support; Stock or stock Options, Spinal Elements: IP royalties, SpineWave: IP royalties, Stout Medical: Stock or stock Options, Stryker: IP royalties, Taylor Francis/Hodder and Stoughton: Publishing royalties, financial or material support, Thieme: Publishing royalties, financial or material support, ViewFi Health: Stock or stock Options; CK - Clinical Spine Surgery: Editorial or governing board, Curetiva: IP royalties, Regeneration Technologies, Inc.: IP royalties; GD – Advance, Medical: Paid consultant, AO Spine: Board or committee member, AOSpine: Other financial or material support, Bioventus: Paid consultant. Cerapedics: Research support, Cervical Spine Research Society: Board or committee member, DePuy, A Johnson and Johnson Company: Research support, Medtronic Sofamor Danek: Research support, Surgalign: Paid consultant, Wolters Kluwer Health - Lippincott Williams and Wilkins: Editorial or governing board.

REFERENCES

1. Asha MJ, Choksey MS, Shad A, Roberts P, Imray C. The role of the vascular surgeon in anterior lumbar spine surgery. *Br J Neurosurg* 2012;26:499-503.
2. Chiriano J, Abou-Zamzam AM Jr., Urayeneza O, Zhang WW, Cheng W. The role of the vascular surgeon in anterior retroperitoneal spine exposure: Preservation of open surgical training. *J Vasc Surg* 2009;50:148-51.
3. Garg J, Woo K, Hirsch J, Bruffey JD, Dilley RB. Vascular complications of exposure for anterior lumbar interbody fusion. *J Vasc Surg* 2010;51:946-50.
4. Jarrett CD, Heller JG, Tsai L. Anterior exposure of the lumbar spine with and without an “access surgeon”: Morbidity analysis of 265 consecutive cases. *J Spinal Disord Tech* 2009;22:559-64.
5. Brau SA, Delamarter RB, Schiffman ML, Williams LA, Watkins RG. Vascular injury during anterior lumbar surgery. *Spine J* 2004;4:409-12.
6. Sasso RC, Best NM, Mummaneni PV, Reilly TM, Hussain SM. Analysis of operative complications in a series of 471 anterior lumbar interbody fusion procedures. *Spine (Phila Pa 1976)* 2005;30:670-4.
7. Mobbs RJ, Phan K, Daly D, Rao PJ, Lennox A. Approach-related

- complications of anterior lumbar interbody fusion: Results of a combined spine and vascular surgical team. *Glob Spine J* 2016;6:147-54.
8. Smith MW, Rahn KA, Shugart RM, Belschner CD, Stout KS, Cheng I. Comparison of perioperative parameters and complications observed in the anterior exposure of the lumbar spine by a spine surgeon with and without the assistance of an access surgeon. *Spine J* 2011;11:389-94.
 9. Menezes CM, Alamin T, Amaral R, Carvalho AD, Diaz R, Guiroy A, et al. Need of vascular surgeon and comparison of value for anterior lumbar interbody fusion (ALIF) in lateral decubitus: Delphi consensus. *Eur Spine J* 2022;31:2270-8.
 10. Phan K, Xu J, Scherman DB, Rao PJ, Mobbs RJ. Anterior lumbar interbody fusion with and without an “access surgeon”: A systematic review and meta-analysis. *Spine (Phila Pa 1976)* 2017;42:E592-601.
 11. Quraishi NA, Konig M, Booker SJ, Shafafy M, Boszczyk BM, Grevitt MP, et al. Access related complications in anterior lumbar surgery performed by spinal surgeons. *Eur Spine J* 2013;22 Suppl 1:S16-20.
 12. Bateman DK, Millhouse PW, Shahi N, Kadam AB, Maltenfort MG, Koerner JD, et al. Anterior lumbar spine surgery: A systematic review and meta-analysis of associated complications. *Spine J* 2015;15:1118-32.
 13. Gumbs AA, Shah RV, Yue JJ, Sumpio B. The open anterior paramedian retroperitoneal approach for spine procedures. *Arch Surg* 2005;140:339-43.
 14. Holt RT, Majd ME, Vadva M, Castro FP. The efficacy of anterior spine exposure by an orthopedic surgeon. *J Spinal Disord Tech* 2003;16:477-86.
 15. Bianchi C, Ballard JL, Abou-Zamzam AM, Teruya TH, Abu-Assal ML. Anterior retroperitoneal lumbosacral spine exposure: Operative technique and results. *Ann Vasc Surg* 2003;17:137-42.
 16. Härtl R, Joeris A, McGuire RA. Comparison of the safety outcomes between two surgical approaches for anterior lumbar fusion surgery: Anterior lumbar interbody fusion (ALIF) and extreme lateral interbody fusion (ELIF). *Eur Spine J Off Publ Eur Spine Soc Eur Spinal Deform Soc Eur Sect Cerv Spine Res Soc* 2016;25:1484-521.
 17. Ozgur BM, Aryan HE, Pimenta L, Taylor WR. Extreme lateral interbody fusion (XLIF): A novel surgical technique for anterior lumbar interbody fusion. *Spine J* 2006;6:435-43.
 18. Ozgur BM, Agarwal V, Nail E, Pimenta L. Two-year clinical and radiographic success of minimally invasive lateral transposas approach for the treatment of degenerative lumbar conditions. *SAS J* 2010;4:41-6.
 19. Deukmedjian AR, Le TV, Dakwar E, Martinez CR, Uribe JS. Movement of abdominal structures on magnetic resonance imaging during positioning changes related to lateral lumbar spine surgery: A morphometric study: Clinical article. *J Neurosurg Spine* 2012;16:615-23.
 20. Lehmen JA, Gerber EJ. MIS lateral spine surgery: A systematic literature review of complications, outcomes, and economics. *Eur Spine J* 2015;24 Suppl 3:287-313.
 21. Epstein NE. Extreme lateral lumbar interbody fusion: Do the cons outweigh the pros? *Surg Neurol Int* 2016;7:S692-700.
 22. Rodgers WB, Gerber EJ, Patterson J. Intraoperative and early postoperative complications in extreme lateral interbody fusion: An analysis of 600 cases. *Spine (Phila Pa 1976)* 2011;36:26-32.
 23. Winder MJ, Gambhir S. Comparison of ALIF versus XLIF for L4/5 interbody fusion: Pros, cons, and literature review. *J Spine Surg* 2016;2:2-8.
 24. Tatsumi R, Lee YP, Khajavi K, Taylor W, Chen F, Bae H. *In vitro* comparison of endplate preparation between four mini-open interbody fusion approaches. *Eur Spine J* 2015;24 Suppl 3:372-7.
 25. Guérin P, Obeid I, Bourghli A, Masquefa T, Luc S, Gille O, et al. The lumbosacral plexus: Anatomic considerations for minimally invasive retroperitoneal transposas approach. *Surg Radiol Anat* 2012;34:151-7.
 26. Berjano P, Gautschi OP, Schils F, Tessitore E. Extreme lateral interbody fusion (XLIF®): How I do it. *Acta Neurochir (Wien)* 2015;157:547-51.
 27. Dakwar E, Val FL, Uribe JS. Trajectory of the main sensory and motor branches of the lumbar plexus outside the psoas muscle related to the lateral retroperitoneal transposas approach. *J Neurosurg Spine* 2011;14:290-5.
 28. Hrabalek L, Adamus M, Gryga A, Wanek T, Tucek P. A comparison of complication rate between anterior and lateral approaches to the lumbar spine. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2014;158:127-32.
 29. Bendersky M, Solá C, Muntadas J, Gruenberg M, Calligaris S, Mereles M, et al. Monitoring lumbar plexus integrity in extreme lateral transposas approaches to the lumbar spine: A new protocol with anatomical bases. *Eur Spine J* 2015;24:1051-7.
 30. Kwon B, Kim DH. Lateral lumbar interbody fusion: Indications, outcomes, and complications. *J Am Acad Orthop Surg* 2016;24:96-105.
 31. Rabau O, Navarro-Ramirez R, Aziz M, Teles A, Mengxiao Ge S, Quillo-Olvera J, et al. Lateral lumbar interbody fusion (LLIF): An update. *Glob Spine J* 2020;10:17S-21S.
 32. Sharma AK, Kepler CK, Girardi FP, Cammisa FP, Huang RC, Sama AA. Lateral lumbar interbody fusion: Clinical and radiographic outcomes at 1 year: A preliminary report. *J Spinal Disord Tech* 2011;24:242-50.
 33. Lee YP, Regev GJ, Chan J, Zhang B, Taylor W, Kim CW, et al. Evaluation of hip flexion strength following lateral lumbar interbody fusion. *Spine J* 2013;13:1259-62.
 34. Knight RQ, Schwaegler P, Hanscom D, Roh J. Direct lateral lumbar interbody fusion for degenerative conditions: Early complication profile. *J Spinal Disord Tech* 2009;22:34-7.
 35. Houten JK, Alexandre LC, Nasser R, Wollowick AL. Nerve injury during the transposas approach for lumbar fusion: Report of 2 cases. *J Neurosurg Spine* 2011;15:280-4.
 36. Lykissas MG, Aichmair A, Hughes AP, Sama AA, Lebl DR, Taher F, et al. Nerve injury after lateral lumbar interbody fusion: A review of 919 treated levels with identification of risk factors. *Spine J* 2014;14:749-58.
 37. Hijji FY, Narain AS, Bohl DD, Ahn J, Long WW, DiBattista JV, et al. Lateral lumbar interbody fusion: A systematic review of complication rates. *Spine J* 2017;17:1412-9.
 38. Mobbs RJ, Phan K, Malham G, Seex K, Rao PJ. Lumbar interbody fusion: Techniques, indications and comparison of interbody fusion options including PLIF, TLIF, MI-TLIF, OLIF/ATP, LLIF and ALIF. *J Spine Surg* 2015;1:2-18.
 39. Arnold P, Anderson K, McGuire R. The lateral transposas approach to the lumbar and thoracic spine: A review. *Surg Neurol Int* 2012;3:198.
 40. Barbagallo GM, Albanese V, Raich AL, Dettori JR, Sherry N, Balsano M. Lumbar lateral interbody fusion (LLIF): Comparative effectiveness and safety versus PLIF/TLIF and predictive factors affecting LLIF outcome. *Evid Based Spine Care J* 2014;5:28-37.
 41. Lee YS, Park SW, Kim YB. Direct lateral lumbar interbody fusion: Clinical and radiological outcomes. *J Korean Neurosurg Soc* 2014;55:248-54.
 42. Jacob KC, Patel MR, Prabhu MC, Vanjani NN, Pawlowski H, Munim MA, et al. Lateral lumbar interbody fusion: Single surgeon learning curve. *World Neurosurg* 2022;164:e411-9.
 43. Aichmair A, Lykissas MG, Girardi FP, Sama AA, Lebl DR, Taher F, et al. An institutional six-year trend analysis of the neurological outcome after lateral lumbar interbody fusion: A 6-year trend analysis of a single institution. *Spine (Phila Pa 1976)* 2013;38:E1483-90.
 44. Le TV, Burkett CJ, Deukmedjian AR, Uribe JS. Postoperative lumbar plexus injury after lumbar retroperitoneal transposas minimally invasive lateral interbody fusion. *Spine (Phila Pa 1976)* 2013;38:E13-20.
 45. Feeley A, Feeley I, Clesham K, Butler J. Is there a variance in complication types associated with ALIF approaches? A systematic review. *Acta Neurochir (Wien)* 2021;163:2991-3004.
 46. Smith WD, Wohns RN, Christian G, Rodgers EJ, Rodgers WB. Outpatient minimally invasive lumbar interbody: Fusion predictive factors and clinical results. *Spine (Phila Pa 1976)* 2016;41 Suppl 8:S106-22.
 47. Youssef JA, McAfee PC, Patty CA, Raley E, DeBauche S, Shucosky E, et al. Minimally invasive surgery: Lateral approach interbody fusion: Results and review. *Spine (Phila Pa 1976)* 2010;35:S302-11.