

2-19-2024

Risk Factors Associated With Poor Outcomes After Quadriceps Tendon Repair

Carlo Coadonato
Thomas Jefferson University


Thomas Jefferson University

Neel K. Patel
Thomas Jefferson University

John Hayden Sonnier
Thomas Jefferson University

Gregory Connors

Follow this and additional works at: https://jdc.jefferson.edu/rothman_institute
See next page for additional authors

 Part of the [Orthopedics Commons](#), and the [Surgery Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Coadonato, Carlo; ; Patel, Neel K.; Sonnier, John Hayden; Connors, Gregory; Sabitsky, Matthew; Johnson, Emma; Mazur, Donald W.; Brahmabhatt, Shyam; and Freedman, Kevin, "Risk Factors Associated With Poor Outcomes After Quadriceps Tendon Repair" (2024). *Rothman Institute Faculty Papers*. Paper 252.
https://jdc.jefferson.edu/rothman_institute/252

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Rothman Institute Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Carlo Coadonato, , Neel K. Patel, John Hayden Sonnier, Gregory Connors, Matthew Sabitsky, Emma Johnson, Donald W. Mazur, Shyam Brahmabhatt, and Kevin Freedman

Risk Factors Associated With Poor Outcomes After Quadriceps Tendon Repair

Carlo Coladonato,* MS, Adeeb Jacob Hanna,* BS, Neel K. Patel,* MD, John Hayden Sonnier,* MD, Gregory Connors,† BS, Matthew Sabitsky,‡ BS, Emma Johnson,* MD, Donald W. Mazur,* MD, Shyam Brahmabhatt,* MD, and Kevin B. Freedman,*§ MD

Investigation performed at Rothman Orthopaedic Institute at the Sidney Kimmel Medical College at Thomas Jefferson University, Philadelphia, Pennsylvania, USA

Background: Ruptures of the quadriceps tendon present most frequently in older adults and individuals with underlying medical conditions.

Purpose: To examine the relationship between patient-specific factors and tear characteristics with outcomes after quadriceps tendon repair.

Study Design: Case-control study; Level of evidence, 3.

Methods: A retrospective review was conducted on all patients who underwent quadriceps tendon repair between January 1, 2016, and January 1, 2021, at a single institution. Patients <18 years and those with chronic quadriceps tendon tears (>6 weeks to surgery) were excluded. Information was collected regarding patient characteristics, presenting symptoms, tear characteristics, physical examination findings, and postoperative outcomes. Poor outcome was defined as a need for revision surgery, complications, postoperative range of motion of (ROM) <110° of knee flexion, and extensor lag of >5°.

Results: A total of 191 patients met the inclusion criteria. Patients were aged 58.5 ± 13.2 years at the time of surgery, were predominantly men (90.6%), and had a mean body mass index (BMI) of 32.2 ± 6.3 kg/m². Patients underwent repair with either suture anchors (15.2%) or transosseous tunnels (84.8%). Postoperatively, 18.5% of patients experienced knee flexion ROM of <110°, 11.3% experienced extensor lag of >5°, 8.5% had complications, and 3.2% underwent revision. Increasing age (odds ratio [OR], 1.03 [95% CI, 1.004-1.07]) and female sex (OR, 3.82 [95% CI, 1.25-11.28]) were significantly associated with postoperative knee flexion of <110°, and increasing age (OR, 1.08 [95% CI, 1.04-1.14]) and greater BMI (OR, 1.14 [95% CI, 1.05-1.23]) were significantly associated with postoperative extensor lag of >5°. Current smoking status (OR, 15.44 [95% CI, 3.97-65.90]) and concomitant retinacular tears (OR, 9.62 [95% CI, 1.67-184.14]) were associated with postoperative complications, and increasing age (OR, 1.05 [95% CI, 1.02-1.08]) and greater BMI (OR, 1.08 [95% CI, 1.02-1.14]) were associated with risk of acquiring any poor outcome criteria.

Conclusion: Patient-specific characteristics—such as increasing age, greater BMI, female sex, retinacular involvement, and current smoking status—were found to be risk factors for poor outcomes after quadriceps tendon repair. Further studies are needed to identify potentially modifiable risk factors that can be used to set patient expectations and improve outcomes.

Keywords: outcomes; prognostic; quadriceps tendon repair

Ruptures of the quadriceps tendon are potentially devastating injuries that can lead to significant morbidity. Although relatively uncommon, they present most

frequently in older adults and those with underlying medical conditions.¹² Specifically, spontaneous bilateral ruptures have been shown to be associated with gout, diabetes, and steroid use.¹⁶ However, acute traumatic tears classically present after a fall, with the patient unable to extend the knee.²⁶ While some tears can be treated conservatively, surgical repair is usually indicated to

The Orthopaedic Journal of Sports Medicine, 12(2), 23259671241229105
DOI: 10.1177/23259671241229105
© The Author(s) 2024

This open-access article is published and distributed under the Creative Commons Attribution - NonCommercial - No Derivatives License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits the noncommercial use, distribution, and reproduction of the article in any medium, provided the original author and source are credited. You may not alter, transform, or build upon this article without the permission of the Author(s). For article reuse guidelines, please visit SAGE's website at <http://www.sagepub.com/journals-permissions>.

fully restore the functionality of the extensor mechanism of the knee.¹⁷ A majority of quadriceps tears are avulsions from the proximal pole of the patella, and the repair of the quadriceps tendon is usually performed using either suture anchors or transosseous tunnels through the patella to restore the bone-tendon interface.³⁷

Patient-specific factors have been shown to be predictive of poor outcomes after surgical treatment of several other knee conditions.²⁸ For example, a body mass index (BMI) of >30 kg/m² has been shown to increase the risk of complications after multiligamentous knee injuries and to be predictive of an inferior range of motion (ROM) after knee arthroplasty.^{10,33} However, there is a paucity of literature regarding the effect of risk factors—such as age, BMI, and smoking—on outcomes of quadriceps tendon repair.

This study aimed to examine the effect of different patient and intraoperative factors on the outcome after quadriceps tendon repair. It was hypothesized that older age, higher BMI, and a history of smoking would be associated with worse outcomes after quadriceps tendon repair.

METHODS

A retrospective review was conducted on patients who underwent quadriceps tendon repair between January 1, 2016, and January 1, 2021, at a single institution, searching the patient database using Current Procedural Terminology codes 27835 and 27836. Patients with definitive documentation of a quadriceps tendon tear within the operative report were included in the study. Those patients with (1) multiligamentous knee injury, (2) age <18 years, or (3) previous ipsilateral quadriceps tendon tear were excluded. Information was collected regarding patient characteristics, presenting symptoms, tear characteristics, physical examination findings, and postoperative outcomes. This study was considered exempt from institutional review board approval.

Preoperative Characteristics

Preoperative descriptive data and intraoperative findings listed in Table 1 were collected. Moreover, the presence of potential risk factors and comorbidities that affect soft tissue healing was determined in each patient—including

TABLE 1
Patient Characteristics (N = 191)^a

Characteristic	Value
Age, y	58.53 ± 13.17
Sex	
Male	173 (90.58)
Female	18 (9.42)
BMI, kg/m ² (n = 189)	32.18 ± 6.32
Tear thickness (n = 190)	
Partial	72 (38.1)
Full	117 (61.9)
Time from injury to surgery, days	15.01 ± 8.03
Concomitant surgery	
Yes	20 (10.47)
No	171 (89.53)
Fixation method	
Transosseous tunnel	162 (84.82)
Suture anchor	29 (15.18)
Preoperative Tegner score (n = 25)	1.88 ± 2.23
Preoperative ROM, deg (n = 99)	80.96 ± 30.19
Preoperative extensor lag $>5^\circ$ (n = 183)	
Yes	167 (91.26)
No	16 (8.74)
Hypertension (n = 189)	
Yes	89 (47.09)
No	100 (52.91)
Hyperlipidemia (n = 190)	
Yes	71 (37.37)
No	119 (62.63)
Endocrine disorders (n = 190)	
None	149 (78.42)
Diabetes	32 (16.84)
Hypothyroid	6 (3.16)
Hyperthyroid	1 (0.53)
Other	2 (1.05)
Chronic kidney disease (n = 190)	
Yes	2 (1.05)
No	188 (98.95)
Smoking status (n = 186)	
Nonsmoker	143 (76.88)
Former smoker	22 (11.83)
Current smoker	21 (11.29)
Fluoroquinolone/steroid use	
Yes	12 (6.28)
No	179 (93.72)

^aData are reported as mean ± SD or n (%). BMI, body mass index.

[§]Address correspondence to Kevin B. Freedman, MD, Rothman Orthopaedics, 825 Old Lancaster Road, Bryn Mawr, PA 19010, USA (email: Kevin.Freedman@rothmanortho.com).

*Rothman Orthopaedic Institute at the Sidney Kimmel Medical College at Thomas Jefferson University, Philadelphia, Pennsylvania, USA.

[†]Drexel University College of Medicine, Philadelphia, Pennsylvania, USA.

[‡]Sidney Kimmel Medical College at Thomas Jefferson University, Philadelphia, Pennsylvania, USA.

Final revision submitted August 11, 2023; accepted August 21, 2023.

One or more of the authors has declared the following potential conflict of interest or source of funding: N.K.P. has received a grant from DJO and education payments from Eclipse Technology Solutions, Paladin Technology Solutions, Smith & Nephew, and Mid-Atlantic Surgical Systems. D.W.M. has received education payments from Arthrex and Liberty Surgical and nonconsulting fees from Paladin Technology Solutions. S.B. has received nonconsulting fees from Arthrex and hospitality payments from Wright Medical Technology and Smith & Nephew. K.B.F. has received a grant from Vericel; education payments from Liberty Surgical; consulting fees from Vericel, Innocoll, and Medical Device Business Services; nonconsulting fees from Vericel; and honoraria from Vericel. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.

Ethical approval for this study was waived by Philadelphia University and Thomas Jefferson University (ref No. 152).

body mass index (BMI), hypertension, hyperlipidemia, diabetes mellitus, thyroid dysfunction, chronic kidney disease, history of steroid or fluoroquinolone use, and smoking history (eg, current, former, never). For this study, age and BMI were analyzed as continuous variables.

Radiographic Evaluation

Magnetic resonance imaging (MRI) was utilized to determine tear thickness, location, concomitant pathology, amount of retraction, retinacular involvement, tendon degeneration, and presence of arthritic changes, which were collected as well. Retinacular involvement was defined as edema and thickening of the retinaculum compatible with a partial tear. On MRI, tendon degeneration was defined as a thickened tendon with signal heterogeneity instead of the usual low signal intensity associated with a healthy tendon. The amount of retraction was calculated by measuring the distance from the tendon stump to the retracted muscle belly tendon.

Surgical Technique and Postoperative Rehabilitation

The quadriceps tendon repair techniques used were either fixation through transosseous patellar tunnels or with suture anchors in the proximal pole of the patella. Regardless of the fixation method, Fiberwire (Arthrex) was passed through the medial and lateral halves of the quadriceps tendon in a Krakow fashion. For the transosseous tunnel group, 3 bone tunnels were drilled in the patella from the proximal pole of the patella to the distal pole and through the patellar tendon. Then, the free limbs of the FiberWire were passed through the tunnels and tied over the bony bridge on the distal pole of the patella. For the suture anchor group, the medial and lateral free limbs of the FiberWire were loaded into 2 SwiveLock anchors (Arthrex), respectively, and those anchors were then placed in the proximal pole of the patella. Both fixation constructs are secured with the knee in full extension.

The initial postoperative rehabilitation protocol included a 6-week period in a hinged knee brace locked in extension for ambulation with a ROM of 0° to 30° at 2 weeks postoperatively, 0° to 70° at 4 weeks postoperatively, 0° to 90° at 6 weeks postoperatively, and full ROM at 8 weeks postoperatively. Weightbearing was allowed immediately postoperatively with a hinged knee brace locked in full extension for the first 6 weeks. After 6 weeks, the brace was unlocked for 2 weeks and then discontinued at 8 weeks. Patients were allowed to return to full activities by 4 months postoperatively.

Postoperative Outcome Measures

Poor outcomes were defined as follows: (1) requiring revision tendon repair/reconstruction; (2) reported complications; (3) postoperative knee flexion ROM <110° as measured passively with a goniometer; and (4) extensor lag of >5° as measured using active motion with

a goniometer. Complications that were considered poor outcomes included postoperative infection, quadriceps retear, arthrofibrosis, deep vein thrombosis, flexion contracture, septic arthritis, and persistent gapping of the repair site. The final ROM was recorded at the 6-month postoperative visit.

Statistical Analysis

A 2-tailed *t* test or the Mann-Whitney *U* test (continuous variables) and the chi-square test or the Fisher exact test (categorical variables) were used to determine significant risk factors for poor outcomes. Statistical significance was set at *P* < .05. Variables with a significant value for bivariate comparison were then analyzed for prediction potential using a multivariate logistic regression model for each category of poor outcome. Because of the limited occurrences within certain poor outcome cohorts, only variables with the most significant *P* value on bivariate analysis were chosen for multivariate regression to minimize overfitting the regression model. All statistical analyses used R Studio Version 4.1.2 (R Project for Statistical Computing).

RESULTS

The study cohort comprised 191 patients (173 men and 18 women), with a mean age of 58.5 ± 13.2 years (Table 1). The mean age of women in this cohort was 58 ± 13.2 years as compared with men, which was 59 ± 13.2 years. The mean BMI for this cohort was 32.2 ± 6.3 kg/m², with no differences in baseline characteristics between sexes. The most common surgical procedure was transosseous fixation (*n* = 162), followed by suture anchors (*n* = 29). Preoperative Tegner scores were not significantly associated with the poor outcomes examined. The mean time to surgery for this cohort was 15 ± 8 days. The time to surgery was not significantly associated with any of the predetermined poor outcomes (Table 2).

Radiographic Findings

Tear location was assessed via magnetic resonance imaging (MRI) and predominantly distal (98.4%), with a large portion of patients experiencing concomitant pathology (77.5%), such as meniscal injury, anterior cruciate ligament tear, and extra articular ligamentous injury (Table 3). A significant difference was found in those with MRI evidence of retinacular involvement that subsequently developed complications (OR, 9.62 [95% CI, 1.67-184.14]).

Loss of Knee Flexion ROM

Of the 191 patients within this cohort, 178 had postoperative ROM data. The mean postoperative knee flexion for the cohort was 124.3° ± 18°. On postoperative physical

TABLE 2
Radiographic Findings in the Overall Cohort^a

Finding	Value
Tear thickness (n = 190)	
Partial	72 (38.1)
Full	117 (61.9)
Tear location (n = 189)	
Distal	186 (98.41)
Central	1 (0.53)
Superficial/medial portion of distal	1 (0.53)
Lateral	1 (0.53)
Presence of concomitant pathology (n = 191)	
Yes	148 (77.49)
No	43 (22.51)
Amount of retraction, cm (n = 124)	2.1 ± 1.18
Retraction present (n = 130)	
Yes	124 (95.38)
No	6 (4.62)
Retinacular involvement (n = 179)	
Yes	107 (59.78)
No	72 (40.22)
Tendon degeneration (n = 188)	
Yes	50 (26.6)
No	138 (73.4)
Arthritis (n = 189)	
Yes	39 (20.63)
No	150 (79.37)

^aData are reported as mean ± SD or n (%).

TABLE 3
Postoperative Outcomes in the Overall Cohort^a

Outcome	Value
Surgical complications (n = 188)	
Yes	16 (8.51)
No	172 (91.49)
Postoperative ROM (n = 178)	122.79 ± 18.92
<110°	33 (18.54)
Normal	145 (81.46)
Extensor lag (n = 186)	
>5°	21 (11.29)
Normal	165 (88.71)
Revision surgery (n = 188)	
Yes	6 (3.19)
No	182 (96.81)

^aData are reported as mean ± SD or n (%). ROM, range of motion.

examination, 18.54% of patients experienced loss of ROM categorized as <110° of knee flexion, with a mean ROM of 122.79° ± 18.92°. Bivariate analysis demonstrated that sex ($P = .020$), endocrine disorders ($P = .005$), and chronic kidney disease ($P = .032$) were associated with decreased postoperative ROM. Logistical regression showed that increased age (odds ratio [OR], 1.03 [95% CI, 1.004-1.07]) and female sex (OR, 3.82 [95% CI, 1.25-11.28]) were associated with postoperative knee flexion of <110° (Table 4). Although significant in bivariate analysis, endocrine

TABLE 4
Multivariate Analysis for Poor Outcomes
(Postoperative Knee Flexion ROM <110°)^a

Predictor (n = 178 observations)	OR (95% CI)	<i>P</i>
Age at surgery, y	1.03 (1.004-1.07)	.031
Female sex	3.82 (1.25-11.28)	.015

^aBoldface *P* values indicate statistical significance ($P < .05$). OR, odds ratio; ROM, range of motion.

TABLE 5
Multivariate Analysis for Poor Outcomes
(Postoperative Extensor Lag >5°)^a

Predictors (n = 186 observations)	OR (95% CI)	<i>P</i>
Age at surgery, y	1.08 (1.04-1.14)	.001
BMI, kg/m ²	1.14 (1.05-1.23)	.001
Smoking status		
Nonsmoker (reference)		
Former smoker	2.52 (0.65-8.79)	.158
Current smoker	3.91 (0.89-15.48)	.055

^aBoldface *P* values indicate statistical significance ($P < .05$). BMI, body mass index; OR, odds ratio.

disorders and chronic kidney disease were not analyzed in the regression because of the limited occurrences within the total cohort. In addition, when subanalyzed by surgical technique, there were no significant differences in rates of loss of ROM between transosseous (16.7%) and suture anchor repair (20.6%) ($P = .585$).

Postoperative Extensor Lag

Of the 191 patients assessed, 186 patients had documentation for knee extension strength postoperatively. On the postoperative physical examination, 11.3% experienced an extensor lag >5°. Bivariate analysis reported significance for age ($P = .017$), preoperative Tegner ($P = .072$), and smoking status ($P = .044$). Logistic regression demonstrated that a postoperative extensor lag of >5° was significantly associated with an increase in age (OR, 1.08 [95% CI, 1.04-1.14]) and BMI (OR, 1.14 [95% CI, 1.05-1.23]). Patient smoking status did not significantly affect postoperative extension lag (Table 5). In addition, in the subanalysis according to surgical technique, there were no significant differences in the rates of postoperative extensor lag between transosseous (10.5%) and suture anchor repair (13.8%) ($P = .515$).

Complications

Within this cohort, 8.51% had any postoperative complications, including postoperative infection (n = 2), quadriceps retear (n = 6), arthrofibrosis (n = 3), deep vein thrombosis (n = 1), septic arthritis (n = 2), and persistent

gapping of the repair site ($n = 2$). Bivariate analysis showed a significant association with evidence of retinacular involvement ($P = .008$), tendon degeneration ($P = .037$), and smoking status ($P < .001$). Multivariate logistical regression showed that postoperative complications were significantly associated with current smoking status (OR, 15.44 [95% CI, 3.97-65.90]) and retinacular involvement assessed via MRI (OR, 9.62 [95% CI, 1.67-184.14]) (Table 6). Former smoking status and female sex were not significantly associated with an increased risk of postoperative complications. In addition, when subanalyzed by surgical technique, there were no significant differences in the rates of complications between transosseous (9.3%) and suture anchor repair (3.4%) ($P = .474$).

Need for Revision Surgery

Revision quadriceps repair was required in 3.14% ($n = 6$) of patients in this cohort. When subanalyzed by surgical technique, there were no differences in rates of revision between transosseous (3.7%) and suture anchor repair (0%) ($P = .593$). Given the small sample size, regression analysis was not completed for this subgroup.

Overall Risk for Poor Outcomes

Patients who experienced any of the 4 specified poor outcomes were grouped into a “poor outcomes” cohort. Of the 191 patients, 54 were designated as poor outcome patients. Bivariate analysis according to outcomes is shown in Table 7. Logistic regression demonstrated that patients who were older (OR, 1.05 [95% CI, 1.02-1.08]) and had a greater BMI (OR, 1.08 [95% CI, 1.02-1.14]) were significantly more likely to experience 1 or more of the poor outcomes examined. Female sex did not have significance when accounting for all combined poor outcome variables. ($P = .1334$) (Table 8). In addition, the subanalysis of surgical techniques did not demonstrate any significant difference in rates of combined poor outcomes between transosseous repair (27.7%) and suture anchor repair (31%) ($P = .893$).

DISCUSSION

The main findings of this study suggest that advanced age, greater BMI, female sex, evidence of retinacular involvement, and current smokers were significantly associated with poor outcomes such as loss of ROM, postoperative extensor lag, and complications. These findings are consistent with several previous studies on various orthopaedic procedures.^{1-3,13-15} Although not seen in the current study, other variables—such as hypertension and chronic kidney disease—have been implicated in previous studies. Lewis et al examined risk factors for infection after quadriceps tendon repair and found that hypertension and hypothyroidism were an independent risk factor for reoperation.¹⁸ Similarly, hypertension has been shown to delay superficial wound healing and increases the risk of readmission

TABLE 6
Multivariate Analysis for Poor Outcomes (Complications)^a

Predictors ($n = 188$ observations)	OR (95% CI)	<i>P</i>
Female sex	1.48 (0.24-7.42)	.645
Smoking status		
Nonsmoker (reference)		
Former smoker	3.60 (3.97-65.90)	.170
Current smoker	15.44 (3.97-65.90)	<.001
Retinacular involvement	9.62 (1.67-184.14)	.038

^aBoldface *P* values indicate statistical significance ($P < .05$). OR, odds ratio.

in orthopaedic surgeries.^{1,5} In our study, bivariate comparisons of patients with hypertension, and all poor outcomes were not significantly associated.

The incidence of quadriceps tendon injury is classically associated with middle-aged men, peaking in the seventh decade and are regarded as uncommon in women of all age groups.¹² Many of the previous studies examining quadriceps tendon repair have an overwhelmingly large ratio of male to female participants, thereby leaving much to be known regarding outcome differences between sexes.^{7,8,24,31} Similarly, the cohort in this study was predominantly men. The present study found that women were significantly more likely to experience a decreased postoperative ROM ($P = .015$). Outcome differences between sexes are well documented in other areas of sports medicine, and authors have hypothesized that sex-specific risk factors—such as anatomic and hormonal differences—may be to blame.^{20,22,34} In the present study, baseline characteristics—such as age and BMI—were no different between men and women ($P = .978$ and $P = .076$, respectively) Similarly, there was no difference in time to surgery between men and women ($P = .123$). Given that the female sex was not statistically associated with the risk of combined poor outcomes, further studies are warranted with larger populations of female patients to discern whether differences exist between the sexes.

Spontaneous quadriceps tendon rupture usually occurs due to the degeneration of tendons in patients caused by systematic diseases such as end-stage renal disease, rheumatoid arthritis, diabetes mellitus, hyperparathyroidism, and long-term steroid use.^{4,29} An epidemiological study found that patients >60 years had a 3-fold increase in the incidence of spontaneous quadriceps tendon rupture compared with patients <40 years.³² In the present cohort, there was no association between quadriceps tendon repair and hypertension, chronic kidney disease, or hyperlipidemia. Of note, the number of patients with chronic kidney disease (1.05%) was quite low in this cohort, and, as a result, there may not have been enough of a sample to observe an impact. Further, endocrine disorders were found to be significantly associated with combined poor outcomes on bivariate analysis. However, limited occurrences in the population did not support regression analysis.

Age at the time of surgery plays a significant role in our ability to recover after injury. Herron et al¹⁵ found that

TABLE 7
Bivariate Analysis According to Patients With and Without Poor Outcomes^a

Characteristic	Poor Outcomes (n = 54)	Without Poor Outcomes (n = 137)	P	Characteristic	Poor Outcomes (n = 54)	Without Poor Outcomes (n = 137)	P
Age, years (n = 191)	62.3 ± 14.3	57 ± 12.4	.034	Chronic kidney disease (n = 190)			.077
Sex (n = 191)			.061	No	51 (96.2)	137 (100)	
Male	45 (83.3)	128 (93.4)		Yes	2 (3.77)	0 (0)	
Female	9 (16.7)	9 (6.57)		Smoking status (n = 186)			.157
BMI, kg/m ² (n = 189)	33.8 ± 7.12	31.6 ± 5.89	.018	Nonsmoker	34 (68)	109 (80.1)	
Tear thickness (n = 190)			.370	Former smoker	7 (14)	15 (11)	
Partial	17 (32.1)	55 (40.4)		Current smoker	9 (18)	12 (8.82)	
Full	36 (67.9)	81 (59.6)		Fluoroquinolone/steroid use (n = 191)			.743
Time from injury to surgery, days (n = 191)	15.6 ± 7.21	14.8 ± 8.33	.250	No	50 (92.6)	129 (94.2)	
Concomitant surgery (n = 191)			0.545	Yes	4 (7.41)	8 (5.84)	
No	50 (92.6)	121 (88.3)		Tear location (n = 189)			>.99
Yes	4 (7.41)	16 (11.7)		Distal	54 (100)	132 (97.8)	
Fixation method (n = 191)			.893	Central	0 (0)	1 (0.74)	
Transosseous tunnel	45 (83.3)	117 (85.4)		Superficial medial portion	0 (0)	1 (0.74)	
Suture anchors	9 (16.7)	20 (14.6)		Lateral	0 (0)	1 (0.74)	
Preoperative Tegner score (n = 25)	1.43 ± 2.30	2.06 ± 2.24	.333	Concomitant pathology (n = 191)			.307
Preoperative extensor lag (n = 183)			>.99	No	9 (16.7)	34 (24.8)	
No	4 (7.69)	12 (9.16)		Yes	45 (83.3)	103 (75.2)	
Yes	48 (92.3)	119 (90.8)		Amount of retraction, cm (n = 130)	2.09 ± 1.02	2.09 ± 1.24	.675
Hypertension (n = 189)			.102	Retraction present (n = 130)			>.99
No	22 (42.3)	78 (56.9)		No	2 (5.13)	4 (4.40)	
Yes	30 (57.7)	59 (43.1)		Yes	37 (94.9)	87 (95.6)	
Hyperlipidemia (n = 190)			.217	Retinacular involvement (n = 179)			.309
No	29 (54.7)	90 (65.7)		No	17 (33.3)	55 (43)	
Yes	24 (45.3)	47 (34.3)		Yes	34 (66.7)	73 (57)	
Endocrine disorders (n = 190)			.035	Tendon degeneration (n = 188)			.960
None	37 (68.5)	112 (81.8)		No	39 (72.2)	99 (73.9)	
Diabetes	11 (20.4)	20 (14.6)		Yes	15 (27.8)	35 (26.1)	
Diabetes, hypothyroid	0 (0)	1 (0.73)		Arthritis (n = 189)			.182
Hypothyroid	4 (7.41)	2 (1.46)		No	39 (72.2)	111 (82.2)	
Hyperthyroid	1 (1.85)	0 (0)		Yes	15 (27.8)	24 (17.8)	
Other	0 (0)	2 (1.46)					

^aData are reported as mean ± SD or n (%). Boldface P values indicate statistically significant differences between groups (P < .05). BMI, body mass index.

orthopaedic trauma in older people is associated with a greater risk of mortality than for similar fractures in the young.¹⁵ However, when it comes to outcomes after quadriceps tendon repair, a recent study has shown that patient reported outcomes in patients <40 years are equivalent to historical controls aged >40 years.³¹ Notably, a considerable percentage of patients <40 years showed significant long-term sequelae, such as pain and stiffness, despite their youth.³¹ As quadricep tendon injuries are most commonly seen in middle-aged men, our cohort aligns with the historical demographic of patients who undergo quadricep tendon repair. In contrast to previous findings, our study reveals a significant risk for decreased postoperative knee flexion, extensor lag, and poor overall outcomes as age increases. However, it is unclear whether the equivalent outcomes shown in previous studies truly capture the impact of age after quadriceps tendon repair or may be a consequence of higher demands and expectations of younger patients.

TABLE 8
Multivariate Analysis for Combined Poor Outcomes^a

Predictors (n = 191 observations)	OR (95% CI)	P
Age, y	1.05 (1.02-1.08)	.002
Female sex	2.24 (0.77-6.52)	.1334
BMI, kg/m ²	1.08 (1.02-1.14)	.010

^aBoldface P values indicate statistical significance (P < .05). BMI, body mass index; OR, odds ratio.

Increased BMI predisposes patients to a range of musculoskeletal and metabolic disorders. In a study by Cardoso et al, investigators found that obese patients had significantly worse long-term outcomes (namely, pain, poor function, and difficulties with activities of daily living) after an operatively treated ankle fracture with a follow-up of over a decade.⁹ Similarly, in a recent systematic review by

Macchi et al,²¹ obesity was reported to increase the risk of tendinopathy, tendon tear, rupture, and postoperative complications after tendon surgery compared with the non-obese population.²¹ Several studies that have examined soft tissue injuries of the knee have shown the negative impact of obesity on ROM, which is consistent with the findings of this study.^{6,19}

Smoking has been implicated widely across medicine as associated with poor wound healing and an array of pathology. The documented effects of byproduct toxic compounds found within cigarette smoke—such as nicotine, carbon monoxide, and hydrogen cyanide—suggest possible mechanisms for impaired wound healing.²³ Yoshikawa and Katada³⁸ found that smoking could be associated with a risk of poor postoperative outcomes irrespective of the type of surgery performed. While this study does not directly investigate the effect of smoking cessation, a previous review showed that longer periods of smoking cessation are more effective for reducing the risk of complications after surgical procedures but with no threshold of time established.³⁶ Thus, there may still be a benefit to emphasizing smoking cessation in the perioperative period, even though quadriceps tendon tears are typically treated within a short period from initial injury. The efficacy of smoking cessation in the perioperative period is unclear, but physicians' awareness of high-risk patients can help set expectations for outcomes.

This study did not find a significant association between time from injury to surgery and poor outcomes. However, in a previous systematic review by Ciriello et al,¹¹ worse outcomes were reported in delayed repairs after quadricep rupture. The mean time to surgery for this cohort was higher than reported in previous studies.^{27,31,35} Overall, all patients were treated within 6 weeks of injury at a mean of 15.01 ± 8.03 days. As the delay from injury to surgical repair increases, the complexity of the surgical repair also increases because of the development of scar contraction and soft tissue retraction. However, successful repair has been described up to 8 years after injury.³⁰ Further studies are needed with larger cohorts to determine the true effect of the timing of surgical repair on overall outcomes.

Limitations

Our study is not without limitations. First, the retrospective nature of the study limits the data collected to recorded information in the patient's electronic medical record. While 5 risk factors were found to significantly affect outcomes after quadriceps tendon repair, the cohort size was relatively small. Thus, there may be additional factors that affect outcomes that were not captured in this study. For example, risk factors such as fluoroquinolone and corticosteroid usage, which have been shown to increase the risk of any tendon rupture, may not have been found to be significant in this study because only 12 patients in the cohort were exposed to either steroids or fluoroquinolones in the time adjacent to the surgery.²⁵ Similarly, the limited data on female patients did not allow for a robust subanalysis of sex-based differences and thus

may not adequately capture the extent of clinical differences. Another limitation of our study was the lack of reporting of contralateral uninjured leg ROM on postoperative testing. Although the threshold was set low at 110° to mitigate overstating patients identified as having poor outcomes in terms of ROM, this did not allow for a baseline comparison of ROM.

CONCLUSION

Patient-specific characteristics—such as increasing age, greater BMI, female sex, retinacular involvement, and smoking status—were risk factors for poor outcomes after quadriceps tendon repair. Further studies are needed to identify potentially modifiable risk factors that can be used to set patient expectations and improve outcomes.

REFERENCES

- Ahmed AA, Moar PA, Kleiner M, Torg JS, Miyamoto CT. Hypertensive patients show delayed wound healing following total hip arthroplasty. *PLOS ONE*. 2011;6(8):e23224. doi:10.1371/journal.pone.0023224
- Alvi HM, Mednick RE, Krishnan V, Kwasny MJ, Beal MD, Manning DW. The effect of BMI on 30 day outcomes following total joint arthroplasty. *J Arthroplasty*. 2015;30(7):1113-1117. doi:10.1016/j.arth.2015.01.049
- Başdelioğlu K. Effects of body mass index on outcomes of total knee arthroplasty. *Eur J Orthop Surg Traumatol Orthop Traumatol*. 2021;31(3):595-600. doi:10.1007/s00590-020-02829-6
- Basic-Jukic N, Juric I, Racki S, Kes P. Spontaneous tendon ruptures in patients with end-stage renal disease. *Kidney Blood Press Res*. 2009;32(1):32-36. doi:10.1159/000201792
- Bhashyam N, Ramos RD la G, Nakhla J, et al. Thirty-day readmission and reoperation rates after single-level anterior cervical discectomy and fusion versus those after cervical disc replacement. *Neurosurg Focus*. 2017;42(2):E6. doi:10.3171/2016.11.FOCUS16407
- Bin Abd Razak HR, Chong HC, Tan HCA. Obesity is associated with poorer range of motion and Tegner scores following hamstring autograft anterior cruciate ligament reconstruction in Asians. *Ann Transl Med*. 2017;5(15):304. doi:10.21037/atm.2017.06.04
- Boudissa M, Roudet A, Rubens-Duval B, Chaussard C, Saragaglia D. Acute quadriceps tendon ruptures: a series of 50 knees with an average follow-up of more than 6 years. *Orthop Traumatol Surg Res OTSR*. 2014;100(2):213-216. doi:10.1016/j.otsr.2013.09.014
- Brossard P, Le Roux G, Vasse B. Acute quadriceps tendon rupture repaired by suture anchors: outcomes at 7 years' follow-up in 25 cases. *Orthop Traumatol Surg Res*. 2017;103(4):597-601. doi:10.1016/j.otsr.2017.02.013
- Cardoso DV, Paccaud J, Dubois-Ferrière V, et al. The effect of BMI on long-term outcomes after operatively treated ankle fractures: a study with up to 16 years of follow-up. *BMC Musculoskelet Disord*. 2022;23(1):317. doi:10.1186/s12891-022-05247-3
- Cho WS, Park JH, Kim JM, Hwang WY, Nam TS. Factors affecting range of motion after total knee arthroplasty. *J Korean Orthop Assoc*. 2003;38(7):683. doi:10.4055/jkoa.2003.38.7.683
- Ciriello V, Gudipati S, Tosounidis T, Soucacos PN, Giannoudis PV. Clinical outcomes after repair of quadriceps tendon rupture: a systematic review. *Injury*. 2012;43(11):1931-1938. doi:10.1016/j.injury.2012.08.044
- Clayton RAE, Court-Brown CM. The epidemiology of musculoskeletal tendinous and ligamentous injuries. *Injury*. 2008;39(12):1338-1344. doi:10.1016/j.injury.2008.06.021
- Divi SN, Goyal DKC, Galetta MS, et al. How does body mass index influence outcomes in patients after lumbar fusion? *Spine*. 2020;45(8):555-561. doi:10.1097/BRS.0000000000003313

14. He Y, Omar M, Feng X, Neunaber C, Jagodzinski M. Impact of smoking on the incidence and post-operative complications of total knee arthroplasty: a systematic review and meta-analysis of cohort studies. *Bosn J Basic Med Sci.* 2022;22(3):353-365. doi:10.17305/bjms.2021.6538
15. Herron J, Hutchinson R, Lecky F, et al. The impact of age on major orthopaedic trauma: an analysis of the United Kingdom Trauma Audit Research Network database. *Bone Jt J.* 2017;99-B(12):1677-1680. doi:10.1302/0301-620X.99B12.BJJ-2016-1140.R2
16. Ilan DI, Tejwani N, Keschner M, Leibman M. Quadriceps tendon rupture: *J Am Acad Orthop Surg.* 2003;11(3):192-200. doi:10.5435/00124635-200305000-00006
17. Konrath GA, Chen D, Lock T, et al. Outcomes following repair of quadriceps tendon ruptures: *J Orthop Trauma.* 1998;12(4):273-279. doi:10.1097/00005131-199805000-00010
18. Lewis DC, Athoff AD, Kamalopathy P, Yarboro SR, Miller MD, Werner BC. Risk factors for infection and revision surgery following patellar tendon and quadriceps tendon repairs: an analysis of 3,442 patients. *J Knee Surg.* 2022;35(13):1495-1502. doi:10.1055/s-0041-1727113
19. Lian J, Patel NK, Nickoli M, et al. Obesity is associated with significant morbidity after multiligament knee surgery. *J Knee Surg.* 2020;33(6):525-530. doi:10.1055/s-0039-1681027
20. Lin CY, Casey E, Herman DC, Katz N, Tenforde AS. Sex differences in common sports injuries. *PM&R.* 2018;10(10):1073-1082.
21. Macchi M, Spezia M, Elli S, Schiaffini G, Chisari E. Obesity increases the risk of tendinopathy, tendon tear and rupture, and postoperative complications: a systematic review of clinical studies. *Clin Orthop.* 2020;478(8):1839-1847. doi:10.1097/CORR.0000000000001261
22. Magnuson JA, Wolf BR, Cronin KJ. Sex-related differences in patients undergoing surgery for shoulder instability: a Multicenter Orthopaedic Outcomes Network (MOON) Shoulder Instability cohort study. *J Shoulder Elbow Surg.* 2019;28(6):1013-1021.
23. McDaniel JC, Browning KK. Smoking, chronic wound healing, and implications for evidence-based practice. *J Wound Ostomy Cont Nurs.* 2014;41(5):415-E2. doi:10.1097/WON.0000000000000057
24. Mille F, Adam A, Aubry S, et al. Prospective multicentre study of the clinical and functional outcomes following quadriceps tendon repair with suture anchors. *Eur J Orthop Surg Traumatol.* 2016;26(1):85-92. doi:10.1007/s00590-015-1710-6
25. Morales DR, Slattery J, Pacurariu A, Pinheiro L, McGettigan P, Kurz X. Relative and absolute risk of tendon rupture with fluoroquinolone and concomitant fluoroquinolone/corticosteroid therapy: population-based nested case-control study. *Clin Drug Investig.* 2019;39(2):205-213. doi:10.1007/s40261-018-0729-y
26. Nori S. Quadriceps tendon rupture. *J Fam Med Prim Care.* 2018;7(1):257. doi:10.4103/jfmpc.jfmpc_341_16
27. O'Shea K, Kenny P, Donovan J, Condon F, McElwain JP. Outcomes following quadriceps tendon ruptures. *Injury.* 2002;33(3):257-260. doi:10.1016/S0020-1383(01)00110-3
28. Patel NK, Lian J, Nickoli M, et al. Risk factors associated with complications after operative treatment of multiligament knee injury. *Orthop J Sports Med.* 2021;9(3):2325967121994203. doi:10.1177/2325967121994203
29. Pei YC, Hsieh PC, Huang LZ, Chiang CK. Simultaneous bilateral quadriceps tendon rupture in a uremic patient. *Formos J Musculoskelet Disord.* 2011;1(2):35-39. doi:10.1016/j.fjmd.2010.12.005
30. Pocock CAJ, Trikha SP, Bell JSP. Delayed reconstruction of a quadriceps tendon. *Clin Orthop.* 2008;466(1):221-224. doi:10.1007/s11999-007-0002-9
31. Rao S, Johnson EE, D'Amore T, Szeto S, Otlans P, Cohen SB. Outcomes after repair of quadriceps tendon rupture in patients aged 40 years and younger. *Orthop J Sports Med.* 2022;10(5):232596712210971. doi:10.1177/23259671221097107
32. Reito A, Paloneva J, Mattila VM, Launonen AP. The increasing incidence of surgically treated quadriceps tendon ruptures. *Knee Surg Sports Traumatol Arthrosc Off J ESSKA.* 2019;27(11):3644-3649. doi:10.1007/s00167-019-05453-y
33. Ridley TJ, Cook S, Bollier M, et al. Effect of body mass index on patients with multiligamentous knee injuries. *Arthrosc J Arthrosc Relat Surg Off Publ Arthrosc Assoc N Am Int Arthrosc Assoc.* 2014;30(11):1447-1452. doi:10.1016/j.arthro.2014.05.035
34. Shi H, Jiang Y, Ren S, Hu X, Huang H, Ao Y. Sex differences in the knee orthopaedic injury patterns among recreational alpine skiers. *BMC Sports Sci Med Rehabil.* 2020;12(1):74. doi:10.1186/s13102-020-00224-6
35. Siwek CW, Rao JP. Ruptures of the extensor mechanism of the knee joint. *J Bone Joint Surg Am.* 1981;63(6):932-937.
36. Theadom A, Cropley M. Effects of preoperative smoking cessation on the incidence and risk of intraoperative and postoperative complications in adult smokers: a systematic review. *Tob Control.* 2006;15(5):352-358. doi:10.1136/tc.2005.015263
37. Yepes H, Tang M, Morris SF, Stanish WD. Relationship between hypovascular zones and patterns of ruptures of the quadriceps tendon. *J Bone Joint Surg Am.* 2008;90(10):2135-2141. doi:10.2106/JBJS.G.01200
38. Yoshikawa R, Katada J. Effects of active smoking on postoperative outcomes in hospitalised patients undergoing elective surgery: a retrospective analysis of an administrative claims database in Japan. *BMJ Open.* 2019;9(10):e029913. doi:10.1136/bmjopen-2019-029913