

8-14-2021

## Psychological Problems Among Patients With Chronic Medical Disorders During the COVID-19 Pandemic

Ali Akbar Asadi-Pooya  
*Thomas Jefferson University*

Seyed Ali Nabavizadeh  
*Shiraz University of Medical Sciences*

Saeid Sadeghian  
*Jundishapur University of Medical Sciences*

Mina Shahisavandi  
*Shiraz University of Medical Sciences*

Zohreh Barzegar  
Follow this and additional works at: <https://jdc.jefferson.edu/neurologyfp>  
*Shiraz University of Medical Sciences*

 Part of the [Neurology Commons](#), [Psychiatry Commons](#), and the [Psychiatry and Psychology Commons](#)

**[Let us know how access to this document benefits you](#)**

*See next page for additional authors*

---

### Recommended Citation

Asadi-Pooya, Ali Akbar; Nabavizadeh, Seyed Ali; Sadeghian, Saeid; Shahisavandi, Mina; Barzegar, Zohreh; Nezafat, Abdullah; Ashjazadeh, Nahid; Bazrafshan Drissi, Hamed; and Sahraian, Ali, "Psychological Problems Among Patients With Chronic Medical Disorders During the COVID-19 Pandemic" (2021). *Department of Neurology Faculty Papers*. Paper 254.  
<https://jdc.jefferson.edu/neurologyfp/254>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Neurology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

---

**Authors**

Ali Akbar Asadi-Pooya, Seyed Ali Nabavizadeh, Saeid Sadeghian, Mina Shahisavandi, Zohreh Barzegar, Abdullah Nezafat, Nahid Ashjazadeh, Hamed Bazrafshan Drissi, and Ali Sahraian

# Psychological Problems Among Patients With Chronic Medical Disorders During the COVID-19 Pandemic

Ali Akbar Asadi-Pooya<sup>1,2</sup>, Seyed Ali Nabavizadeh<sup>1</sup>, Saeid Sadeghian<sup>3</sup>, Mina Shahisavandi<sup>1</sup>, Zohreh Barzegar<sup>1</sup>, Abdullah Nezafat<sup>1</sup>, Nahid Ashjazadeh<sup>1</sup>, Hamed Bazrafshan Drissi<sup>4</sup>, Ali Sahraian<sup>1</sup>

<sup>1</sup> Epilepsy Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

<sup>2</sup> Department of Neurology, Jefferson Comprehensive Epilepsy Center, Thomas Jefferson University, Philadelphia, USA

<sup>3</sup> Department of Pediatric Neurology, Golestan Medical, Educational, and Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

<sup>4</sup> Department of Cardiovascular, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Received: 02 Dec. 2020; Accepted: 18 Jun. 2021

**Abstract-** The aim of the current study was to investigate the rates of stress, anxiety, and depression among people in south Iran (a group from the general population without a history of any chronic medical problems, and cohorts of patients were recruited from epilepsy, diabetes, and cardiac disease clinics). We surveyed a sample of people during September 2020: a group of the general population without a history of any chronic medical problems, people with epilepsy, people with diabetes mellitus (DM), and people with cardiac problems. The survey included four general questions and two COVID-19 specific questions [contracting COVID-19, relatives with COVID-19]. Furthermore, the survey included the DASS (Depression-Anxiety-Stress Scale)-21 questionnaire. 487 people were surveyed (154 people with epilepsy, 127 patients with DM, 98 people with cardiac problems, and 108 healthy individuals). Among people without a history of any chronic medical illnesses, 14% had any psychological problems. The highest rates of depression and anxiety were observed among patients with DM (52% and 57%, respectively), and the highest rate of increased stress was observed among people with cardiac problems (40%). The existence of any underlying medical problem was significantly associated with higher rates of depression, anxiety, and stress among the participants. While many patients with underlying chronic medical conditions suffer from depression, anxiety, and stress during the COVID-19 pandemic, we cannot establish a cause and effect relationship between the COVID-19 pandemic and increased psychological problems among these patients.

© 2021 Tehran University of Medical Sciences. All rights reserved.

*Acta Med Iran* 2021;59(6):322-326.

**Keywords:** Coronavirus disease 2019 (COVID-19); Anxiety; Depression; Epilepsy; Seizure; Stress

## Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has caused a catastrophic pandemic (COVID-19) since late 2019 (1). This virus is highly contagious and has a high potential for human-to-human transmission. It may cause a severe and fatal illness characterized by acute respiratory distress syndrome, multi-organ failure, and death (1). However, in addition to the physical health hazards of this viral infection, the psychological impacts of the COVID-19 pandemic on

different groups of people should also be taken into account seriously (2,3). Mental health issues such as stress, anxiety, depression, frustration, and uncertainty have emerged progressively during the COVID-19 outbreak. Similarly, studies performed on the psychological impacts of previous infectious outbreaks, such as the severe acute respiratory syndrome (SARS), have found significant psychological problems (e.g., anxiety, depression, or stress) among the public during the outbreak (4).

Major risk factors for a more severe illness due to

**Corresponding Author:** A.A. Asadi-Pooya

Epilepsy Research Center, Shiraz University of Medical Sciences, Shiraz, Iran  
Tel: +98 9352274990, Fax: +98 7136125841 E-mail address: aliasadipooya@yahoo.com

Copyright © 2021 Tehran University of Medical Sciences. Published by Tehran University of Medical Sciences

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited

COVID-19 have been identified as older age and underlying chronic medical disorders, among others (1,5,6). Cerebrovascular diseases (odds ratio [OR]: 3.66), chronic obstructive pulmonary disease (OR: 2.39), cardiovascular diseases (OR: 2.84), diabetes mellitus (DM) (OR: 2.78), and hypertension (OR: 2.24) have been associated with severe COVID-19 (5). Therefore, it is plausible to assume that people with chronic medical problems are also at higher risk of experiencing undue stress, anxiety, and depression compared with that in the general public due to the fear of contracting a more severe illness and its potential consequences.

The aim of the current study was to investigate the rates of stress, anxiety, and depression among people in south Iran. We investigated four groups of people in order to clarify the psychological impacts of the COVID-19 pandemic on different groups of patients with underlying medical problems; a group from the general population without a history of any chronic medical problems, patients with epilepsy, patients with DM, and patients with cardiac problems. We hypothesized that the rates of stress, anxiety, and depression are significantly higher among patients with the chronic medical condition compared with those among healthy individuals.

## Materials and Methods

### Participants

In this cross-sectional study, we surveyed four groups of people during September 2020: a group of the general population from Shiraz (a major city in south Iran) without a history of any chronic medical problems selected randomly, patients with epilepsy (referring consecutively to the neurology clinic at Shiraz University of Medical Sciences), patients with DM (referring consecutively to the DM clinic at Shiraz University of Medical Sciences), and patients with chronic cardiac problems (referring consecutively to the cardiology clinic at Shiraz University of Medical Sciences). The inclusion criteria were adults ( $\geq 18$  years) and literacy ( $> 5$  years of education). The exclusion criteria included intellectual disability and unwillingness to participating in the study.

### Instruments

The survey included four general questions [age, sex, education (school vs. college), and medical/psychiatric problems]. It also included two COVID-19 specific questions [contracting COVID-19 (self-declared), relatives (i.e., spouse, children, siblings, parents) with COVID-19 (self-declared)]. Furthermore, the survey included the DASS (Depression-Anxiety-Stress Scale)-

21 questionnaire (Appendix 1). In this short version of the DASS questionnaire, each item (depression, anxiety, and stress) is assessed by seven questions. The DASS-21 scale has been translated and validated in the Farsi language (7).

### Statistical analyses

Statistical analyses were performed using the independent *t*-test, Fisher's Exact test, Pearson chi-square test, and Bonferroni correction test. Moderate, severe, or extremely severe depression/anxiety/stress was considered as the existence of a psychological burden (vs. normal and mild scores). Variables with a  $P < 0.1$  in univariate tests were assessed in a logistic regression model. Odds ratio (OR) and 95% confidence interval (CI) were calculated. A  $P$  (2-sided) less than 0.05 was considered significant.

## Results

The total number of the participants was 487 people [154 patients with epilepsy (66 men and 88 women), 127 patients with DM (40 men and 88 women), 98 patients with cardiac problems (54 men and 44 women), and 108 healthy individuals (66 men and 88 women)]. The mean age of the participants was 37 years (standard deviation: 16 years) (range: 18 to 97 years). The mean age of the groups differed significantly ( $27 \pm 9$  years in controls,  $33 \pm 12$  years in the epilepsy group,  $43 \pm 18$  years among patients with DM, and  $50 \pm 15$  years among those with cardiac problems;  $P = 0.0001$ ). They included 271 females and 216 males. The sex ratios of the groups were not significantly different ( $P = 0.07$ ).

Moderate, severe, or extremely severe depression was observed among 33% of the participants (160 people). Moderate, severe, or extremely severe anxiety was observed among 41% of the participants (201 people). Moderate, severe, or extremely severe stress was observed among 29% of the participants (139 people). Table 1 shows the factors associated with depression/anxiety/stress in univariate analysis. The existence of an underlying medical disorder was consistently associated with higher rates of depression, anxiety, and stress among the studied participants (compared with that in healthy individuals). Among people without a history of any chronic medical illnesses, these rates were 14% for all three psychological problems (depression/anxiety/stress). The highest rates of depression and anxiety were observed among patients with DM (52% and 57%, respectively); patients with epilepsy had a risk less than those with DM and more than

patients with cardiac problems. The highest rate of increased stress was observed among patients with cardiac problems (40%); patients with epilepsy had a risk lower than those with DM and also patients with cardiac problems.

We then analyzed the association between depression/anxiety/stress and variables with a  $P < 0.1$  in a binary logistic regression model. The models that were generated by regression analysis were significant

( $P = 0.0001$ ). Table 2 shows the results of these analyses. The existence of an underlying medical problem was significantly associated with higher rates of depression, anxiety, and stress among the participants (with robust ORs). Female sex was significantly associated with higher rates of depression and anxiety (with modest ORs). A history of relatives with COVID-19 was significantly associated with higher rates of depression (with a modest OR) (Table 2).

**Table 1. Factors associated with depression/anxiety/stress in univariate analysis**

	Depression		P	df
	With (N=160)	Without (N=327)		
Sex (Female: Male)	100: 60	171: 156	0.041	1
Mean age ± Standard deviation (years)	40 ± 16	36 ± 16	0.038	485
Education (college)	58 (36%)	131 (40%)	0.428	1
Medical problem (none, epilepsy, diabetes, cardiac)	15 (14%), 50 (32%), 66 (52%), 29 (30%)	93 (86%), 104 (68%), 61 (48%), 69 (70%)	0.0001	3
Relatives with COVID-19	36 (23%)	50 (15%)	0.058	1
History of COVID-19	13 (8%)	29 (9%)	0.865	1
<b>Anxiety</b>				
	With (N=201)	Without (N=286)	P	df
Sex (Female: Male)	121: 80	150: 136	0.096	1
Mean age ± Standard deviation (years)	39 ± 16	36 ± 16	0.073	485
Education (college)	72 (36%)	117 (41%)	0.257	1
Medical problem (none, epilepsy, diabetes, cardiac)	15 (14%), 78 (51%), 73 (57%), 35 (36%)	93 (86%), 76 (49%), 54 (43%), 63 (64%)	0.0001	3
Relatives with COVID-19	41 (20%)	45 (16%)	0.187	1
History of COVID-19	14 (7%)	28 (10%)	0.326	1
<b>Stress</b>				
	With (N=139)	Without (N=348)	P	df
Sex (Female: Male)	84: 55	187: 161	0.190	1
Mean age ± Standard deviation (years)	41 ± 16	36 ± 16	0.003	485
Education (college)	44 (32%)	145 (42%)	0.039	1
Medical problem (none, epilepsy, diabetes, cardiac)	15 (14%), 38 (25%), 47 (37%), 39 (40%)	93 (86%), 116 (75%), 80 (63%), 59 (60%)	0.0001	3
Relatives with COVID-19	31 (22%)	55 (16%)	0.114	1
History of COVID-19	12 (8%)	30 (9%)	1.000	1

\*After Bonferroni correction, a significant predictive value is 0.008. The significant P is in bold. df: degree of freedom

**Table 2. Factors associated with depression/anxiety/stress in logistic regression models (compared with healthy individuals)**

	Depression		P	df
	Odds Ratio	95% confidence interval		
Medical problem (epilepsy/ diabetes/ cardiac)	3.13/ 6.96/ 2.89	1.63-6.03/ 3.49-13.93/ 1.32-6.30	0.001/ 0.0001/ 0.007	1
Sex (Female)	1.72	1.14- 2.59	0.010	1
Relatives with COVID-19	1.71	1.03-2.85	0.038	1
Age			0.731	1
	Anxiety			
Medical problem (epilepsy/ diabetes/ cardiac)	6.52/ 8.48/ 3.56	3.44-12.36/ 4.27-16.85/ 1.67-7.58	0.0001/ 0.0001/ 0.001	1
Sex (Female)	1.51	1.02- 2.36	0.038	1
Age			0.756	1
	Stress			
Medical problem (epilepsy/ diabetes/ cardiac)	1.90/ 3.25/ 3.56	0.95-3.77/ 1.61-6.57/ 1.64-7.69	0.066/ 0.001/ 0.001	1
Age				0.611
Education (college)				0.525

## Discussion

In this cross-sectional study, we observed that many patients with various underlying chronic medical conditions suffer from depression, anxiety, and stress during the COVID-19 pandemic. In a national face-to-face household survey from Iran that was published five years ago, the most common category of psychiatric disorders was any anxiety disorder (15.6%), and the most prevalent particular disorder was major depressive disorder (12.7%) (8). These rates are comparable with what we observed among the healthy individuals in the current study (14% for both depression and anxiety). Furthermore, in studies from the pre-COVID era in Iran, the prevalence of depression among Iranian patients with DM was estimated to be 62% (95% confidence interval [CI]: 57-67), and the prevalence of anxiety among Iranian patients with DM was estimated to be 65% (95% CI: 42-82) (9). The prevalence of depression among cardiovascular patients in Iran was estimated to be 47% (95% CI: 38-56) in one study (10). Finally, in one cross-sectional hospital-based study of 74 adults with epilepsy, 26 (35%) patients had symptoms of depression (11). These rates are also comparable with what we observed among patients with DM, cardiovascular disorders, and epilepsy in the current study.

Therefore, while we observed that many patients with underlying chronic medical conditions (i.e., epilepsy, DM, and cardiovascular disorders) suffer from depression, anxiety, and stress during the COVID-19 pandemic, we cannot establish or even hypothesize a cause and effect relationship between COVID-19 pandemic and increased psychological burden and problems among these patients based on the current study. However, the prevalence of depressive and anxiety symptoms was reported to be significantly higher in the UK, relative to pre-pandemic epidemiological data (12). This cross-cultural difference should be studied more in the future.

We can provide the following speculation to explain our counterintuitive findings described above. The general public in Iran has already been under tremendous psychological pressure due to various reasons (e.g., economic, political, etc.) (13). On top of that, patients with chronic medical problems (e.g., epilepsy, DM, etc.) in Iran may suffer from psychological disorders as a comorbidity to their underlying illness (as any other person elsewhere in the world) (14) or because of the existing undue pressure due to social reasons (e.g., shortage of their drugs due to economic sanctions) (13).

The interactions between chronic diseases and psychological problems have been described in the literature (15). However, why the COVID-19 pandemic has not (significantly) worsened the already high rates of psychological problems in Iran is an intriguing finding that should be explored in future studies.

This study has some limitations. This was a cross-sectional study, and we did not have the pre-pandemic epidemiological data or mental health histories of the participants. In addition, the groups differed with respect to their age, and this might have affected the results. Furthermore, we did not investigate the drug history (e.g., antidepressants) of the patients. Finally, the timing between getting ill with COVID-19 (in the affected people) and completing these questionnaires was not recorded.

We can conclude that many patients with various underlying chronic medical conditions suffer from depression, anxiety, and stress during the COVID-19 pandemic. The observed high rates of depression, anxiety, and stress in Iran (during the COVID-19 pandemic or at any other time), particularly among patients with underlying medical problems, are alarming and should be taken into account seriously. These psychological problems (depression, anxiety, and stress) are not only associated with impaired quality of life (16) but also may affect the health status of the people and even increase the risk of premature death (17). There are various interventions available that may help reduce or prevent depression, anxiety, and stress among the affected people and also those at risk; health promotion interventions, exercise, and yoga, are some examples (18-20). Having said that, the key element is to screen the at risk populations and detect any psychological problem as soon as possible.

## References

1. Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *J Autoimmun* 2020;109:102433.
2. Luo M, Guo L, Yu M, Wang H. The Psychological and Mental Impact of Coronavirus Disease 2019 (COVID-19) on Medical Staff and General Public—A Systematic Review and Meta-analysis. *Psychiatry Res* 2020;291:113190.
3. Vahedian-Azimi A, Moayed MS, Rahimibashar F, Shojaei S, Ashtari S, Pourhoseingholi MA. Comparison of the severity of psychological distress among four groups of an Iranian population regarding COVID-19 pandemic. *BMC*

- Psychiatry 2020;20:402.
4. Maunder R, Hunter J, Vincent L, Bennett J, Peladeau N, Leszcz M. The immediate psychological and occupational impact of the 2003 SARS outbreak in a teaching hospital. *CMAJ* 2003;168:1245-51.
  5. Del Sole F, Farcomeni A, Loffredo L, Carnevale R, Menichelli D, Vicario T, et al. Features of severe COVID-19: A systematic review and meta-analysis. *Eur J Clin Invest* 2020;50:e13378.
  6. Romero Starke K, Petereit-Haack G, Schubert M, Kämpf D, Schliebner A, Hegewald J, et al. The Age-Related Risk of Severe Outcomes Due to COVID-19 Infection: A Rapid Review, Meta-Analysis, and Meta-Regression. *Int J Environ Res Public Health* 2020;17:5974.
  7. Samani S, Jowkar B. Validity of Farsi version of DASS (Depression-Anxiety-Stress Scale). *Human Soc Sci* 2006;26:65-77.
  8. Sharifi V, Amin-Esmaeili M, Hajebi A, Motevalian A, Radgoodarzi R, Hefazi M, et al. Twelve-month prevalence and correlates of psychiatric disorders in Iran: the Iranian Mental Health Survey, 2011. *Arch Iran Med* 2015;18:76-84.
  9. Khalighi Z, Badfar G, Mahmoudi L, Soleymani A, Azami M, Shohani M. The prevalence of depression and anxiety in Iranian patients with diabetes mellitus: A systematic review and meta-analysis. *Diabetes Metab Syndr* 2019;13:2785-94.
  10. Ghaemmohamadi MS, Behzadifar M, Ghashghae A, Mousavinejad N, Ebadi F, Saedi Shahri SS, et al. Prevalence of depression in cardiovascular patients in Iran: A systematic review and meta-analysis from 2000 to 2017. *J Affect Disord* 2018;227:149-55.
  11. Foroughipour M, Mokhber N, Azarpajoo MR, Taghavi M, Modarres Gharavi M, Akbarzadeh F, et al. Coping mechanisms: depression and suicidal risk among patients suffering from idiopathic epilepsy. *Int J High Risk Behav Addict* 2013;1:178-82.
  12. Pieh C, Budimir S, Delgadillo J, Barkham M, Fontaine JRJ, Probst T. Mental health during COVID-19 lockdown in the United Kingdom. *Psychosom Med* 2021;83:328-37.
  13. Asadi-Pooya AA, Azizimalamiri R, Badv RS, Yarali B, Asadollahi M, Homayoun M, et al. Impacts of the international economic sanctions on Iranian patients with epilepsy. *Epilepsy Behav* 2019;95:166-8.
  14. Asadi-Pooya AA, Kanemoto K, Kwon OY, Taniguchi G, Dong Z, Chinvarun Y, et al. Depression in people with epilepsy: How much do Asian colleagues acknowledge it? *Seizure* 2018;57:45-9.
  15. Conversano C. Psychological common factors in chronic diseases. *Frontiers Psychol* 2019;10:2727.
  16. Michaelis R, Tang V, Wagner JL, Modi AC, LaFrance WC Jr, Goldstein LH, et al. Cochrane systematic review and meta-analysis of the impact of psychological treatments for people with epilepsy on health-related quality of life. *Epilepsia* 2018;59:315-32.
  17. Keller A, Litzelman K, Wisk LE, Maddox T, Cheng ER, Creswell PD, et al. Does the perception that stress affects health matter? The association with health and mortality. *Health Psychol* 2012;31:677-84.
  18. James-Palmer A, Anderson EZ, Zucker L, Kofman Y, Daneault JF. Yoga as an Intervention for the Reduction of Symptoms of Anxiety and Depression in Children and Adolescents: A Systematic Review. *Front Pediatr* 2020;8:78.
  19. Morris L, Stander J, Ebrahim W, Eksteen S, Meaden OA, Ras A, et al. Effect of exercise versus cognitive behavioural therapy or no intervention on anxiety, depression, fitness and quality of life in adults with previous methamphetamine dependency: a systematic review. *Addict Sci Clin Pract* 2018;13:4.
  20. Goodarzi A. A comprehensive review on COVID-19 infection and comorbidities of various organs. *Acta Med Iran* 2021;59:4-14.