

7-16-2022

Predictive Factors and Outcomes for Ibrutinib in Relapsed/ refractory Marginal Zone Lymphoma: A multicenter Cohort Study

Narendranath Epperla
The Ohio State University

Qihong Zhao
The Ohio State University

Sayan Mullick Chowdhury

The Ohio State University works at: <https://jdc.jefferson.edu/medfp>

Part of the Hematology Commons, and the Oncology Commons

Washington University in St. Louis

[Let us know how access to this document benefits you](#)

Tamara K Moyo

Atrium Health
Recommended Citation

Epperla, Narendranath; Zhao, Qihong; Chowdhury, Sayan Mullick; Shea, Lauren; Moyo, Tamara K; Reddy, Nishitha; Sheets, Julia; Weiner, David M; Geethakumari, Praveen Ramakrishnan; Kandarpa, Malathi; Bruno, Ximena Jordan, Thomas, Collin, Churnetski, Michael C; Hsu, Andrew; Zurbruggen, Luke; Tan, Cherie; Lindsey, Kathryn; Maakaron, Joseph; Caimi, Paolo F; Torka, Pallawi; Bello, Celeste; Ayyappan, Sabarish; Karmali, Reem; Kim, Seo-Hyun; Kress, Anna; Kothari, Shalin; Sawalha, Yazeed; Christian, Beth; David, Kevin A; Greenwell, Irl Brian; Janakiram, Murali; Kenkre, Vaishalee P; Olszewski, Adam J; Cohen, Jonathon B; Palmisiano, Neil D.; Umyarova, Elvira; Wilcox, Ryan A; Awan, Farrukh T; Alderuccio, Juan Pablo; Barta, Stefan K; Grover, Natalie S; Ghosh, Nilanjan; Bartlett, Nancy L; Herrera, Alex F; and Shouse, Geoffrey, "Predictive Factors and Outcomes for Ibrutinib in Relapsed/refractory Marginal Zone Lymphoma: A multicenter Cohort Study" (2022). *Department of Medicine Faculty Papers*. Paper 365.
<https://jdc.jefferson.edu/medfp/365>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Medicine Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Narendranath Epperla, Qihong Zhao, Sayan Mullick Chowdhury, Lauren Shea, Tamara K Moyo, Nishitha Reddy, Julia Sheets, David M Weiner, Praveen Ramakrishnan Geethakumari, Malathi Kandarpa, Ximena Jordan Bruno, Colin Thomas, Michael C Churnetski, Andrew Hsu, Luke Zurbriggen, Cherie Tan, Kathryn Lindsey, Joseph Maakaron, Paolo F Caimi, Pallawi Torka, Celeste Bello, Sabarish Ayyappan, Reem Karmali, Seo-Hyun Kim, Anna Kress, Shalin Kothari, Yazeed Sawalha, Beth Christian, Kevin A David, Irl Brian Greenwell, Murali Janakiram, Vaishalee P Kenkre, Adam J Olszewski, Jonathon B Cohen, Neil D. Palmisiano, Elvira Umyarova, Ryan A Wilcox, Farrukh T Awan, Juan Pablo Alderuccio, Stefan K Barta, Natalie S Grover, Nilanjan Ghosh, Nancy L Bartlett, Alex F Herrera, and Geoffrey Shouse

CORRESPONDENCE

Open Access



Predictive factors and outcomes for ibrutinib in relapsed/refractory marginal zone lymphoma: a multicenter cohort study

Narendranath Epperla^{1*}, QiuHong Zhao¹, Sayan Mullick Chowdhury¹, Lauren Shea², Tamara K. Moyo³, Nishitha Reddy⁴, Julia Sheets⁵, David M. Weiner⁶, Praveen Ramakrishnan Geethakumari⁷, Malathi Kandarpa⁸, Ximena Jordan Bruno⁹, Colin Thomas¹⁰, Michael C. Churnetski¹¹, Andrew Hsu¹², Luke Zurbriggen¹³, Cherie Tan¹⁴, Kathryn Lindsey¹⁵, Joseph Maakaron¹⁶, Paolo F. Caimi¹⁷, Pallawi Torka¹⁸, Celeste Bello¹⁹, Sabarish Ayyappan²⁰, Reem Karmali²¹, Seo-Hyun Kim²², Anna Kress²³, Shalin Kothari²³, Yazeed Sawalha¹, Beth Christian¹, Kevin A. David¹⁴, Irl Brian Greenwell¹⁵, Murali Janakiram¹⁶, Vaishalee P. Kenkre¹³, Adam J. Olszewski¹², Jonathon B. Cohen¹¹, Neil Palmisiano¹⁰, Elvira Umyarova⁹, Ryan A. Wilcox⁸, Farrukh T. Awan⁷, Juan Pablo Alderuccio²⁴, Stefan K. Barta⁶, Natalie S. Grover⁵, Nilanjan Ghosh³, Nancy L. Bartlett², Alex F. Herrera^{25†} and Geoffrey Shouse^{25†}

Abstract

Ibrutinib is effective in the treatment of relapsed/refractory (R/R) marginal zone lymphoma (MZL) with an overall response rate (ORR) of 48%. However, factors associated with response (or lack thereof) to ibrutinib in R/R MZL in clinical practice are largely unknown. To answer this question, we performed a multicenter (25 US centers) cohort study and divided the study population into three groups: “ibrutinib responders”—patients who achieved complete or partial response (CR/PR) to ibrutinib; “stable disease (SD)”; and “primary progressors (PP)” —patients with progression of disease as their best response to ibrutinib. One hundred and nineteen patients met the eligibility criteria with 58%/17% ORR/CR, 29% with SD, and 13% with PP. The median PFS and OS were 29 and 71.4 months, respectively, with no difference in PFS or OS based on the ibrutinib line of therapy or type of therapy before ibrutinib. Patients with complex cytogenetics had an inferior PFS (HR = 3.08, 95% CI 1.23–7.67, $p = 0.02$), while those with both complex cytogenetics (HR = 3.00, 95% CI 1.03–8.68, $p = 0.04$) and PP (HR = 13.94, 95% CI 5.17–37.62, $p < 0.001$) had inferior OS. Only primary refractory disease to first-line therapy predicted a higher probability of PP to ibrutinib (RR = 3.77, 95% CI 1.15–12.33, $p = 0.03$). In this largest study to date evaluating outcomes of R/R MZL treated with ibrutinib, we show that patients with primary refractory disease and those with PP on ibrutinib are very high-risk subsets and need to be prioritized for experimental therapies.

Keywords: Marginal zone lymphoma, MZL, Ibrutinib, Relapsed, Refractory

[†]Alex F. Herrera and Geoffrey Shouse co-senior authors

*Correspondence: Narendranath.Epperla@osumc.edu

¹ Division of Hematology, Department of Medicine, The Ohio State University, Columbus, OH 43210, USA

Full list of author information is available at the end of the article

To the editor

Marginal zone lymphomas (MZL) are the third most common B cell non-Hodgkin lymphoma (NHL) comprising 7% of all NHLs [1–3]. Ibrutinib was FDA-approved for relapsed or refractory (R/R) MZL based on phase II clinical trial that showed an overall response rate (ORR)



of 48% [4]. In the recently updated long-term follow-up of this study, the ORR was 58% with a median duration of response (DOR) of 27.6 months [5]. However, factors associated with response (or lack thereof) to ibrutinib in R/R MZL in clinical practice are largely unknown. Hence, we sought to evaluate characteristics predictive of ibrutinib failure in R/R MZL and describe the outcomes of patients on ibrutinib therapy in a “real-world” setting.

In this multicenter retrospective cohort study, we included adult patients (18 years or older) with R/R MZL who received ibrutinib monotherapy between 2010 and 2019 at 25 US medical centers. The study population was divided into 3 groups: “ibrutinib responders”—patients who achieved a complete response (CR) or partial response (PR) to ibrutinib as their best response; “stable disease (SD)”; and “primary progressors (PP)”—patients with progression of disease as their best response to ibrutinib. The primary objective of the study was to evaluate the *real-world* efficacy outcomes of ibrutinib in R/R MZL including response rates, duration of response (DOR), progression-free survival (PFS), and overall survival (OS). Secondary objectives included the evaluation of factors predictive of PP, PFS, and OS. See supplementary appendix for definitions and statistical analysis.

A total of 119 patients met the inclusion criteria. Sixty-nine patients achieved a disease response (ORR 58% with a CR rate of 17%), 35 (29%) had SD, and 15 (13%) had PP. Table 1 shows the baseline characteristics of the patient population. Among the 69 patients who achieved CR/PR, median DOR was 36.8 months (95% CI 25.5–NR) (Additional file 1: Fig. S1A). When stratified by CR or PR status (Additional file 1: Fig. S1B), median DOR was not reached (NR) (95% CI 32–NR) in patients who achieved CR compared to 26 months (95% CI 20.2–NR) in those achieving PR ($p=0.057$). Median PFS and OS for the entire group ($n=119$) were 29 months (Additional file 1: Fig. S2A) and 71.4 months (Additional file 1: Fig. S2B), respectively. The 1-year and 2-year PFS and OS rates were 66% and 55%, and 87% and 85%, respectively. When stratified by the ibrutinib line of therapy (second line vs. third line vs. fourth line and beyond), there was no difference in PFS (median PFS in similar order, 28.5 months vs. 28.2 months vs. 39.8 months, respectively, $p=0.89$,

Additional file 1: Fig. S3A) or OS (median OS in similar order, NR vs. 71.4 months vs. 44.5 months, respectively, $p=0.37$, Additional file 1: Fig. S3B). Among the factors evaluated to determine the predictors of PFS and OS (see Additional file 1: Tables S1 and S2), complex cytogenetics portended inferior PFS (HR=3.08, 95% CI 1.23–7.67, $p=0.02$), while both complex cytogenetics (HR=3.00, 95% CI 1.03–8.68, $p=0.04$) and PP (HR=13.94, 95% CI 5.17–37.62, $p<0.001$) were associated with poor OS. Among the factors evaluated for association with PP (Table 2), only primary refractory disease (to first-line therapy) predicted a higher probability of PP to ibrutinib (RR=3.77, 95% CI 1.15–12.33, $p=0.03$). Lastly, the prior line of therapy (Additional file 1: Table S3) was not associated with differences in outcomes associated with ibrutinib therapy (Additional file 1: Figs. S4 and S5).

In this multicenter retrospective study, we made several important observations. First, the ORR to ibrutinib was 58% with predominantly PRs (41%), which is in line with the results of the phase 2 registration trial [4, 5]. Second, the median DOR was 36.8 months and was longer in those achieving CR compared to PR (although not statistically significant). Third, patients with primary refractory disease had a significantly higher probability of progression on ibrutinib. Fourth, there was no difference in the PFS, or OS based on the number or type of prior therapies. This is in contrast to the data in mantle cell lymphoma, wherein the greatest benefit from ibrutinib was noted in patients receiving ibrutinib in earlier lines of therapy (especially second-line therapy) [6]. Fifth, the presence of complex cytogenetics was predictive of inferior PFS and OS.

The ORR and DOR with ibrutinib in R/R MZL patients in our study were in line with the results of the phase 2 registration trial [4, 5]. The median PFS, however, was longer in the current study (29 months) compared to the previously published results (15.7 months) [5]. One plausible explanation could be the receipt of rituximab monotherapy prior to ibrutinib, which was higher in the current study (49% vs. 27% in the phase 2 trial), as the median PFS was 30.4 months in the recipients of rituximab monotherapy in the clinical trial [5]. Another possible explanation is that in clinical trials routine scans

Table 1 Baseline characteristics

	All patients N = 119 (%)	IB CR + PR N = 69 (%)	IB SD N = 35 (%)	IB PD N = 15 (%)	p value
Median age at diagnosis in years (range)	64 (23–90)	66 (23–90)	63 (40–86)	64 (38–89)	0.67
Median age at ibrutinib therapy in years (range)	68 (27–91)	69 (27–90)	67 (42–86)	65 (41–91)	0.74
Gender					0.89
Male	55 (46)	33 (48)	15 (43)	7 (47)	
Female	64 (54)	36 (52)	20 (57)	8 (53)	
BMI					0.95
< 30	71 (71)	43 (72)	20 (69)	8 (73)	
≥ 30	29 (29)	17 (28)	9 (31)	3 (27)	
Missing	19	9	6	4	
ECOG PS at diagnosis					0.53
0	46 (46.5)	25 (42)	13 (50)	8 (61)	
1	47 (47.5)	32 (53)	11 (42)	4 (31)	
≥ 2	6 (6)	3 (5)	2 (8)	1 (8)	
Missing	20	9	9	2	
MZL subtype					0.97
NMZL	50 (42)	28 (41)	17 (49)	5 (33)	
SMZL	29 (24)	17 (25)	8 (23)	4 (27)	
EMZL	40 (34)	24 (34)	10 (28)	6 (40)	
Stage at diagnosis					0.95
1–2	19 (17)	11 (16)	6 (18)	2 (14)	
3–4	96 (83)	56 (84)	28 (82)	12 (86)	
Missing	4	2	1	1	
B symptoms at diagnosis					0.62
No	81 (74)	44 (70)	25 (78)	12 (80)	
Yes	29 (26)	19 (30)	7 (22)	3 (20)	
Missing	9	6	3	0	
LDH higher than institutional baseline					0.80
No	70 (71)	43 (71)	20 (74)	7 (64)	
Yes	29 (29)	18 (29)	7 (26)	4 (36)	
Missing	20	8	8	4	
Albumin at diagnosis					0.75
Normal	80 (81)	49 (80)	22	9	
Low	19 (19)	12 (20)	4	3	
Missing	20	8	9	3	
Monoclonal protein at diagnosis					0.05
No	49 (56)	30 (54)	17 (74)	2 (25)	
Yes	38 (44)	26 (46)	6 (26)	6 (75)	
Missing	32	13	12	7	
BM involvement at diagnosis					0.72
No	32 (32)	17 (30)	10 (33)	5 (42)	
Yes	67 (68)	40 (70)	20 (67)	7 (58)	
Not done	20	11	5	3	
TP53 mutation/17p deletion (n = 67)*					0.99
No	19 (28)	10 (23)	7 (47)	2 (25)	
Yes	10 (15)	7 (16)	2 (13)	1 (13)	
Unavailable/not tested	38 (57)	27 (61)	6 (40)	5 (62)	
Complex cytogenetics (n = 67)*					0.16
No	57 (85)	37 (90)	17 (85)	6 (67)	
Yes	10 (15)	4 (10)	3 (15)	3 (33)	

Table 1 (continued)

	All patients N = 119 (%)	IB CR + PR N = 69 (%)	IB SD N = 35 (%)	IB PD N = 15 (%)	p value
Primary refractory disease**					0.07
No	89 (75)	56 (81)	25 (71)	8 (53)	
Yes	30 (25)	13 (19)	10 (29)	7 (47)	
First-line therapy					0.47
Rituximab	58 (49)	35 (51)	19 (54)	4 (27)	
BR	30 (25)	16 (23)	9 (26)	5 (33)	
R-CVP	11 (9)	7 (10)	1 (3)	3 (20)	
R-CHOP	9 (8)	5 (7)	2 (6)	2 (13)	
Others	11 (9)	6 (9)	4 (11)	1 (7)	
Receipt of maintenance R					0.27
No	88 (74)	47 (68)	29 (83)	12 (80)	
Yes	31 (26)	22 (32)	6 (17)	3 (20)	
Line of ibrutinib therapy					0.40
Second line	54 (45)	31 (45)	16 (46)	7 (47)	
Third line	41 (35)	27 (39)	9 (26)	5 (33)	
Fourth line and beyond	24 (20)	11 (16)	10 (28)	3 (20)	
Median f/up in months (range)^	23 (1–75)	23 (1–72)	26 (3–75)	6 (3–22)	

CR complete response, PR partial response, SD stable disease, PD progressive disease, BMI body mass index, ECOG PS Eastern Cooperative Oncology Group performance status, MZL marginal zone lymphoma, LDH lactate dehydrogenase, BM bone marrow, BR bendamustine rituximab, R-CVP rituximab, cyclophosphamide, vincristine, prednisone, R-CHOPrituximab, cyclophosphamide, Adriamycin, vincristine, prednisone, f/up follow-up

*Only among those who had bone marrow involvement. Complex karyotype was defined as the presence of at least three chromosomal aberrations in at least two cells

**Primary refractory disease: defined as progression of disease at the end of induction therapy or within 6 months of treatment completion. Among these 30 patients, 13 received rituximab, 9 received BR, 4 received R-CHOP, 3 received R-CVP, 1 received other

^Among those who are alive

are performed at frequent intervals and radiologic but asymptomatic relapses are picked up. Scheduled surveillance scans are typically less frequent outside of a clinical trial, and as a result, asymptomatic progressions may not be identified until a patient experiences clinical evidence of progression, thus making the PFS appear longer.

We did not capture the information on the toxicity and dose modification of ibrutinib (dose interruption or discontinuation) in the current study. Other limitations

include the lack of data on CD5, Ki-67 expression, and *MYD88* mutation status precluding our ability to study the impact of these variables on response and survival.

In conclusion, in this first and the largest study to date to report the *real-world* outcomes of R/R MZL treated with ibrutinib, we show that patients with primary refractory disease and those with PP on ibrutinib are very high-risk subsets and need to be prioritized for experimental and cellular therapies.

Table 2 Modeling on risk of progression on ibrutinib

Variable	PP versus CR/PR			SD versus CR/PR		
	RR	95% CI	p value	RR	95% CI	p value
Age at diagnosis	1.02	0.96–1.08	0.61	0.99	0.96–1.03	0.65
Gender						
Male	Referent					
Female	1.05	0.34–3.22	0.93	1.22	0.54–2.78	0.63
BMI						
< 30	Referent					
≥ 30	0.95	0.22–4.04	0.94	1.14	0.43–3.01	0.79
ECOG PS at diagnosis						
0	Referent					
1	0.39	0.10–1.46	0.16	0.66	0.25–1.73	0.40
≥ 2	1.04	0.09–11.61	0.97	1.28	0.19–8.75	0.80
MZL subtype						
NMZL	Referent					
SMZL	1.32	0.31–5.63	0.71	0.78	0.27–2.19	0.63
EMZL	1.40	0.38–5.20	0.61	0.69	0.26–1.79	0.44
Stage at diagnosis						
1–2	Referent					
3–4	1.18	0.23–6.06	0.84	0.92	0.31–2.75	0.88
B symptoms at diagnosis						
No	Referent					
Yes	0.58	0.15–2.30	0.44	0.65	0.24–1.76	0.40
LDH higher than institutional baseline						
No	Referent					
Yes	1.37	0.35–5.28	0.65	0.84	0.30–2.33	0.73
Monoclonal protein at diagnosis						
No	Referent					
Yes	3.46	0.64–18.84	0.15	0.41	0.14–1.19	0.10
BM involvement at diagnosis						
No	Referent					
Yes	0.60	0.16–2.15	0.43	0.85	0.33–2.20	0.74
TP53 mutation/17p deletion						
No	Referent					
Yes	0.76	0.08–7.25	0.81	0.81	0.15–4.48	0.81
Complex cytogenetics						
No	Referent					
Yes	4.63	0.81–26.35	0.08	1.63	0.32–8.21	0.55
Primary refractory disease*						
No	Referent					
Yes	3.77	1.15–12.33	0.03	1.72	0.66–4.47	0.26
Line of ibrutinib therapy						
Second line	Referent					
Third line	0.82	0.23–2.90	0.76	0.65	0.24–1.70	0.38
Fourth line and beyond	1.21	0.26–5.54	0.81	1.76	0.62–5.04	0.29

CR complete response, PR partial response, SD stable disease, PD progressive disease, BMI body mass index, ECOG PS Eastern Cooperative Oncology Group performance status, MZL marginal zone lymphoma, LDH lactate dehydrogenase, BM bone marrow

*Primary refractory disease: defined as progression of disease at the end of induction therapy or within 6 months of treatment completion

Abbreviations

MZL: Marginal zone lymphoma; R/R: Relapsed/refractory; ORR: Overall response rate; CR: Complete remission; PFS: Progression-free survival; OS: Overall survival.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13045-022-01316-1>.

Additional file 1. Supplemental Appendix.

Acknowledgements

None.

Author contributions

Narendranath Epperla contributed to conception and design and collection and assembly of data. Qihong Zhao and Narendranath Epperla analyzed the data. All authors interpreted the data. In manuscript writing, the first draft was prepared by Narendranath Epperla. All authors provided critical scientific input and provided the final approval of manuscript.

Funding

None.

Availability of data and materials

Please contact author for data request.

Declarations**Ethics approval and consent to participate**

The study was approved by the institutional review boards at all the participating sites and performed in compliance with the Declaration of Helsinki.

Consent for publication

Not applicable.

Competing interests

NE: Speakers Bureau for Incyte, Honoraria/consulting/ad boards for TG Therapeutics, Pharmacyclics, BeiGene, Seattle Genetics, and Novartis; Research funding BeiGene. TKM ad board for Seattle Genetics. NR: Consultancy KITE, AbbVie, BMS, Celgene; Research funding Genentech, BMS. PRG: Consultancy to Kite Pharma, Rafael Pharma, Pharmacyclics LLC, and BMS. PC: Research funding ADC Therapeutics, Genentech; Ad board: ADC Therapeutics, Genentech, Bayer, Verastem, KITE; Speakers Bureau Celgene. PT: Consulting fees from ADC Therapeutics, TG Therapeutics, Kura Oncology, and Genentech. SA: Consulting/Ad board for TG Therapeutics, Seattle Genetics, BeiGene, Intellisphere, Fate Therapeutics, AstraZeneca. RK: Advisory Board: Celgene Corporation, Gilead Sciences, Juno Therapeutics, Kite Pharma, Janssen, Karyopharm, Pharmacyclics, MorphoSys, Epizyme, Genentech/Roche, EUSA; Grants/Research Support: Celgene Corporation/Juno Therapeutics/BMS, Takeda, BeiGene, Gilead Sciences/Kite. Speakers Bureau: AstraZeneca, BeiGene, Gilead Sciences, MorphoSys. SK Consultation with Karyopharm and Incyte. YS: Research funding from BMS, Celgene, TG Therapeutics, and BeiGene and has consulted for TG Therapeutics and Epizyme. BC: Research funding: Acerta, Celgene, Genentech, Merck, Millennium, MorphoSys, Roche, Triphase; Ad board: Verastem, Seattle Genetics, AstraZeneca. MJ: Research funding Takeda, Fate, Nektar. AJO: Research funding Genentech, Spectrum Pharmaceuticals, TG Therapeutics, Adaptive Biotech. JCB: Consulting/Advisory board—Janssen, BeiGene, AstraZeneca, Loxo/Lilly, Aptitude Health, Kite/Gilead, HutchMed; Research Funding: AstraZeneca, BMS/Celgene, Genentech, Loxo/Lilly, Takeda, Novartis, HutchMed, Biinvent. NP: Research funding from Genentech and AbbVie. FTA: Consultancy to Genentech, AstraZeneca, AbbVie, Janssen, Pharmacyclics, Gilead Sciences, Kite Pharma, Celgene, Karyopharm, MEI Pharma, Verastem, Incyte, BeiGene, Johnson and Johnson, Dava Oncology, BMS, Merck, Cardinal Health, ADCT therapeutics, Epizyme. JPA: Consulting fees and research funding from ADC Therapeutics. SKB: Consultancy Monsanto; Honoraria: Atara, Seattle Genetics, Janssen, Pfizer. NSG: Research funding Genentech; honoraria/consulting/ad boards Kite, ADC, Novartis. NG: Consulting/advisory role for

Seattle Genetics, TG Therapeutics, AstraZeneca, Pharmacyclics, BMS, Gilead, BeiGene, Incyte, Karyopharm, Roche/Genentech, Novartis, Loxo/Lilly, Genmab, Adaptive Biotech. NLB: Research funding: ADC Therapeutics, Autolus, BMS, Celgene, Forty Seven, Genentech, Immune Design, Janssen, Merck, Millennium, Pharmacyclics, Affirmed Therapeutics, Dynavax, Gilead, MedImmune, Novartis; Consulting/Ad board: Kite Pharma, Pfizer, ADC Therapeutics, Roche/Genentech, Seattle Genetics, BTG, Acerta. GS: Honoraria from Kite Pharmaceuticals and BeiGene. AFH: Consultancy: BMS, Merck, Gilead, Adaptive Biotech, Seattle Genetics, Karyopharm; Research funding: Merck, Genentech, Gilead, Seattle Genetics, Immune Design, AstraZeneca, Pharmacyclics, ADCT Therapeutics. QZ, SMC, LS, JS, DMW, MK, XJB, CT, MCC, AH, LZ, CT, KL, KM, CB, SHK, AK, KAD, IBG, VPK, EU, and RAW have no relevant COI.

Author details

¹Division of Hematology, Department of Medicine, The Ohio State University, Columbus, OH 43210, USA. ²Washington University, St. Louis, MO, USA. ³Levine Cancer Center, Atrium Health, Charlotte, NC, USA. ⁴Vanderbilt University Medical Center, Nashville, TN, USA. ⁵University of North Carolina, Chapel Hill, NC, USA. ⁶Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA, USA. ⁷Harold C. Simmons Comprehensive Cancer Center, University of Texas Southwestern Medical Center, Dallas, TX, USA. ⁸Rogel Cancer Center, University of Michigan, Ann Arbor, MI, USA. ⁹University of Vermont, Burlington, VT, USA. ¹⁰Thomas Jefferson University, Philadelphia, PA, USA. ¹¹Winship Cancer Institute, Emory University Medical Center, Atlanta, GA, USA. ¹²Brown University, Providence, RI, USA. ¹³University of Wisconsin, Madison, WI, USA. ¹⁴Cancer Institute of New Jersey, New Brunswick, NJ, USA. ¹⁵Medical University of South Carolina, Charleston, SC, USA. ¹⁶University of Minnesota, Minneapolis, MN, USA. ¹⁷University Hospitals Seidman Cancer Center, Cleveland, OH, USA. ¹⁸Roswell Park Cancer Institute, Buffalo, NY, USA. ¹⁹H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL, USA. ²⁰University of Iowa, Iowa City, IA, USA. ²¹Northwestern University, Chicago, IL, USA. ²²Rush University, Chicago, IL, USA. ²³Yale University, New Haven, CT, USA. ²⁴University of Miami, Miami, FL, USA. ²⁵City of Hope, Duarte, CA, USA.

Received: 30 May 2022 Accepted: 8 July 2022

Published online: 16 July 2022

References

1. Teras LR, DeSantis CE, Cerhan JR, Morton LM, Jemal A, Flowers CR. 2016 US lymphoid malignancy statistics by World Health Organization subtypes. *CA Cancer J Clin.* 2016;66(6):443–59.
2. Swerdlow SH, Campo E, Pileri SA, Harris NL, Stein H, Siebert R, et al. The 2016 revision of the World Health Organization classification of lymphoid neoplasms. *Blood.* 2016;127(20):2375–90.
3. Cerhan JR, Habermann TM. Epidemiology of marginal zone lymphoma. *Ann Lymphoma.* 2021;5:1. <https://doi.org/10.21037/aol-20-28>.
4. Noy A, de Vos S, Thieblemont C, Martin P, Flowers CR, Morschhauser F, et al. Targeting Bruton tyrosine kinase with ibrutinib in relapsed/refractory marginal zone lymphoma. *Blood.* 2017;129(16):2224–32.
5. Noy A, de Vos S, Coleman M, Martin P, Flowers CR, Thieblemont C, et al. Durable ibrutinib responses in relapsed/refractory marginal zone lymphoma: long-term follow-up and biomarker analysis. *Blood Adv.* 2020;4(22):5773–84.
6. Rule S, Dreyling M, Goy A, Hess G, Auer R, Kahl B, et al. Ibrutinib for the treatment of relapsed/refractory mantle cell lymphoma: extended 3.5-year follow up from a pooled analysis. *Haematologica.* 2019;104(5):e211–4.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.