

8-9-2022

## GRK2 Regulates ADP Signaling in Platelets Via P2Y1 and P2Y12.

Xuefei Zhao

*Thomas Jefferson University*

Matthew Cooper

*Thomas Jefferson University*

James V Michael

*Thomas Jefferson University*

Yanki Yarman


*Thomas Jefferson University*

Aiden Baltz

*Thomas Jefferson University*

Follow this and additional works at: <https://jdc.jefferson.edu/medfp>

See next page for additional authors

 Part of the [Hematology Commons](#), and the [Translational Medical Research Commons](#)

[Let us know how access to this document benefits you](#)

---

### Recommended Citation

Zhao, Xuefei; Cooper, Matthew; Michael, James V; Yarman, Yanki; Baltz, Aiden; Chuprun, J Kurt; Koch, Walter J; McKenzie, Steven E.; Tomaiuolo, Maurizio; Stalker, Timothy J; Zhu, Li; and Ma, Peisong, "GRK2 Regulates ADP Signaling in Platelets Via P2Y1 and P2Y12." (2022). *Department of Medicine Faculty Papers*. Paper 370.

<https://jdc.jefferson.edu/medfp/370>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Medicine Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

---

**Authors**

Xuefei Zhao, Matthew Cooper, James V Michael, Yanki Yarman, Aiden Baltz, J Kurt Chuprun, Walter J Koch, Steven E. McKenzie, Maurizio Tomaiuolo, Timothy J Stalker, Li Zhu, and Peisong Ma

# GRK2 regulates ADP signaling in platelets via P2Y<sub>1</sub> and P2Y<sub>12</sub>

Xuefei Zhao,<sup>1,2</sup> Matthew Cooper,<sup>1</sup> James V. Michael,<sup>1</sup> Yanki Yarman,<sup>1</sup> Aiden Baltz,<sup>1</sup> J. Kurt Chuprun,<sup>3</sup> Walter J. Koch,<sup>3</sup> Steven E. McKenzie,<sup>1</sup> Maurizio Tomaiuolo,<sup>4</sup> Timothy J. Stalker,<sup>1</sup> Li Zhu,<sup>2,\*</sup> and Peisong Ma<sup>1,\*</sup>

<sup>1</sup>Cardeza Foundation for Hematologic Research, Department of Medicine, Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, PA; <sup>2</sup>Cyrus Tang Hematology Center, Soochow University, Suzhou, China; <sup>3</sup>Center for Translational Medicine, Department of Pharmacology, Lewis Katz School of Medicine, Temple University, Philadelphia, PA; and <sup>4</sup>Vickie and Jack Farber Vision Research Center, Wills Eye Hospital, Philadelphia, PA

## Key Points

- GRK2 regulates the hemostatic response by limiting ADP P2Y<sub>1</sub>- and P2Y<sub>12</sub>-mediated signaling.
- Maintaining GRK2 activity in platelets may be beneficial for prevention of thrombotic diseases.

The critical role of G protein-coupled receptor kinase 2 (GRK2) in regulating cardiac function has been well documented for >3 decades. Targeting GRK2 has therefore been extensively studied as a novel approach to treating cardiovascular disease. However, little is known about its role in hemostasis and thrombosis. We provide here the first evidence that GRK2 limits platelet activation and regulates the hemostatic response to injury. Deletion of GRK2 in mouse platelets causes increased platelet accumulation after laser-induced injury in the cremaster muscle arterioles, shortens tail bleeding time, and enhances thrombosis in adenosine 5'-diphosphate (ADP)-induced pulmonary thromboembolism and in FeCl<sub>3</sub>-induced carotid injury. GRK2<sup>-/-</sup> platelets have increased integrin activation, P-selectin exposure, and platelet aggregation in response to ADP stimulation. Furthermore, GRK2<sup>-/-</sup> platelets retain the ability to aggregate in response to ADP restimulation, indicating that GRK2 contributes to ADP receptor desensitization. Underlying these changes in GRK2<sup>-/-</sup> platelets is an increase in Ca<sup>2+</sup> mobilization, RAS-related protein 1 activation, and Akt phosphorylation stimulated by ADP, as well as an attenuated rise of cyclic adenosine monophosphate levels in response to ADP in the presence of prostaglandin I<sub>2</sub>. P2Y<sub>12</sub> antagonist treatment eliminates the phenotypic difference in platelet accumulation between wild-type and GRK2<sup>-/-</sup> mice at the site of injury. Pharmacologic inhibition of GRK2 activity in human platelets increases platelet activation in response to ADP. Finally, we show that GRK2 binds to endogenous G<sub>βγ</sub> subunits during platelet activation. Collectively, these results show that GRK2 regulates ADP signaling via P2Y<sub>1</sub> and P2Y<sub>12</sub>, interacts with G<sub>βγ</sub>, and functions as a signaling hub in platelets for modulating the hemostatic response to injury.

## Introduction

Blood platelets are small, anucleate cells that play a central role in hemostasis, thrombosis, and inflammation. Prior to injury, platelets circulate in a discoid resting state. When challenged by vascular injury, platelets rapidly activate and aggregate with each other, forming a plug on the vessel wall to prevent vascular leakage. Platelet activation needs to be regulated so that platelets can be prepared for rapid activation after vascular injury to prevent bleeding while at the same time avoiding an overly exuberant response to

Submitted 7 January 2022; accepted 10 June 2022; prepublished online on *Blood Advances* First Edition 17 June 2022; final version published online 4 August 2022. DOI 10.1182/bloodadvances.2022007007.

\*L.Z. and P.M. contributed equally to this study.

The data sets and protocols in this study are available to other investigators from the corresponding author without unreasonable restrictions (e-mail: peisong.ma@jefferson.edu).

The full-text version of this article contains a data supplement.

© 2022 by The American Society of Hematology. Licensed under Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

injury that may lead to ischemia, heart attack, or stroke. Although initial adhesion of platelets to the damaged vessel wall is driven by collagen, subsequent recruitment of additional platelets into a growing thrombus requires mediators such as thrombin, thromboxane A<sub>2</sub> (TxA<sub>2</sub>), and adenosine 5'-diphosphate (ADP), all of which act through G protein-coupled receptors (GPCRs).<sup>1,2</sup> Mechanisms that provide negative feedback toward activated GPCRs to limit platelet activation and thrombus formation are thus key to achieving an optimal response to vascular injury. Emerging studies from our group and others strongly suggest that GPCR kinases (GRKs) are critical negative regulators for platelet activation.<sup>3-7</sup>

GRKs are a group of serine/threonine kinases that typically terminate agonist-occupied GPCR signaling in a phosphorylation-dependent manner. In cells other than platelets, GRKs phosphorylate agonist-bound GPCRs, triggering arrestin binding followed by clathrin-mediated endocytosis.<sup>8</sup> We have been able to detect the expression of GRK2, GRK5, and GRK6 in platelets.<sup>3</sup> Over the last 30 years, the role of GRK2 in cardiovascular disease has been widely studied, and it has been reported that GRK2 messenger RNA and its activity increase in patients with cardiac ischemia and hypertrophy.<sup>9,10</sup> Conversely, the impact of GRK2 on hemostasis and thrombosis remains unknown. Using a transfected cell assay, Hardy et al<sup>11</sup> suggest that GRK2 is involved in ADP P2Y<sub>12</sub> receptor-mediated platelet desensitization. However, little is known about the negative feedback mechanisms governing ADP P2Y receptor-mediated platelet activation.

Here, we show that GRK2<sup>-/-</sup> mice display greater platelet activation and platelet accumulation at the site of the injury. The increased responsiveness to agonists in GRK2<sup>-/-</sup> platelets is limited to ADP P2Y<sub>1</sub>- and P2Y<sub>12</sub>-mediated signaling. Absence of GRK2 in platelets increases thrombosis in ADP-induced pulmonary thromboembolism and in FeCl<sub>3</sub>-induced carotid injury. P2Y<sub>12</sub> antagonist treatment eliminates the phenotypic difference in platelet accumulation between wild-type (WT) and GRK2<sup>-/-</sup> mice in response to injury. Pharmacologic inhibition of GRK2 in human platelets causes an increase in platelet activation in response to 2-methylthioadenosine diphosphate (2MesADP). Collectively, these observations show for the first time that GRK2 functions as a critical negative regulator in hemostasis and thrombosis.

## Methods

### Vascular injury: platelet accumulation and fibrin generation

GRK2<sup>-/-</sup> (GRK2<sup>fl/fl</sup>/Pf4-Cre<sup>+</sup>) male mice with matched WT control littermates (GRK2<sup>fl/fl</sup>/Pf4-Cre<sup>-</sup>) aged 8 to 12 weeks were used. Briefly, Alexa Fluor 568-labeled anti-CD41 antibody F(ab)<sub>2</sub> fragments, Alexa Fluor 488-labeled anti-P-selectin, and Alexa Fluor 647-labeled fibrinogen were administered via the retro-orbital plexus. Arterioles in the cremaster muscle of 30 to 50 μm in diameter were studied. Vascular injury was induced by using a laser ablation system (Ablate! photoablation system; Intelligent Imaging Innovations [3i]). Thrombus formation was observed for 3 minutes and analyzed as previously described.<sup>3</sup> We also examined laser-induced thrombus formation in WT and GRK2<sup>-/-</sup> mice treated with a P2Y<sub>12</sub> antagonist, cangrelor. Cangrelor (0.75 μg before each injury) or vehicle (saline) was administered via IV infusion through the jugular vein.

The studies involved in the research reported in this article have been approved by the Thomas Jefferson University Institutional Review Board and Institutional Animal Care and Use Committee. The study was conducted in accordance with the Declaration of Helsinki.

### Pulmonary thromboembolism

Pulmonary thromboembolism assay was performed as we previously described.<sup>6</sup> Briefly, Alexa Fluor 750-labeled anti-GPIX was administered IV into anesthetized WT and GRK2<sup>-/-</sup> mice. After 20 minutes, the anesthetized mice were dosed with ADP (0.3 mg/g) IV. The lung was imaged by using the Licor imaging system.

### Statistical analysis

Results are presented as mean ± standard error of the mean. Data were analyzed by using the *t* test or analysis of variance followed by appropriate post hoc test. *P* < .05 was considered statistically significant. Additional statistical analysis and methods are described in the supplemental Figures 3, 4, 8, 9, and 10 and Figure legends.

## Results

### Generation and characterization of GRK2<sup>-/-</sup> mice


GRK2<sup>-/-</sup> mice die during gestation due to retardation of cardiac development.<sup>12,13</sup> Thus, we took advantage of a GRK2 conditional knockout mouse model. We crossed "floxed" GRK2 mice, containing two loxP sites in intron 2 and intron 6 of the GRK2 locus,<sup>13</sup> with Pf4-Cre mice expressing the Cre recombinase under the control of the megakaryocyte-specific platelet factor-4 (Pf4) promoter. This allowed for deletion of GRK2 specifically in megakaryocytes and platelets (Figure 1A; supplemental Figure 1).<sup>14</sup> GRK2<sup>lox/lox</sup>/Pf4-Cre<sup>-</sup> are henceforth referred to as WT, and GRK2<sup>lox/lox</sup>/Pf4-Cre<sup>+</sup> are further referred to as GRK2<sup>-/-</sup> in this study.

GRK2<sup>-/-</sup> mice grew and developed normally. This platelet-specific deletion of GRK2 in mice did not affect the expression of the other 2 major GRKs (GRK5 and GRK6) in platelets (Figure 1B; supplemental Figure 2A). The platelet number, mean platelet volume, red blood cell count, and white blood cell count of GRK2<sup>-/-</sup> mice were comparable to those of littermate WT control mice (supplemental Table 1). The surface expression of integrin is normal on GRK2<sup>-/-</sup> platelets (supplemental Figure 2B).

### GRK2<sup>-/-</sup> mice display increased platelet accumulation at the outer layer of the thrombus in response to injury

Platelet function in vivo was assessed with real-time confocal fluorescence microscopy following a laser-induced injury in the cremaster muscle arterioles.<sup>15</sup> Hemostatic thrombi that form after penetrating injuries in this setting have a characteristic core/shell architecture in which the extent of platelet activation is determined by the distribution and concentration of agonists in the immediate environment of each platelet. In the core region of the thrombus, thrombin mediates platelet activation with minimal requirement of ADP and TxA<sub>2</sub>. In contrast, ADP and TxA<sub>2</sub> signaling are critical for formation of the outer shell region.<sup>15-17</sup>

Representative images of hemostatic plugs formed in WT and GRK2<sup>-/-</sup> mice at 1.5 minutes after injury are shown in Figure 2Ai. Platelet accumulation occurred at the same initial rate in GRK2<sup>-/-</sup>



**Figure 1. Generation and characterization of platelet-specific deletion of GRK2 in mouse.** (A) Schematic map of loxP sites inserted in the mouse GRK2 gene. Two loxP sites flanking exons III through VI of mouse GRK2 are shown as triangles. (B) Western blot images of platelet lysate from PF4-Cre<sup>-</sup> and PF4-Cre<sup>+</sup> mice probed with the GRK2-, GRK5-, and GRK6-specific antibodies. Deletion of GRK2 in platelets does not affect the expression of GRK5 and GRK6. *n* = 4. Supplemental Figure 2 features the full blots. ns, not significant.

mice as in WT control mice. However, after reaching 40 seconds, there was a decline in CD41 fluorescence area in control mice that did not occur in the GRK2<sup>-/-</sup> mice (Figure 2Aii). Total platelet accumulation in the GRK2<sup>-/-</sup> mice was approximately one-third greater than in the control mice, as measured by area under the curve (Figure 2Aiii; supplemental Figure 3A). This increase was due to an expansion of the P-selectin (-) shell region, rather than the P-selectin (+) core region (Figure 2Aiv; supplemental Figure 3B). Fibrin accumulation was unaffected in GRK2<sup>-/-</sup> mice (Figure 2B; supplemental Figure 3C).

### Deletion of GRK2 in mouse platelets increases the response to ADP stimulation ex vivo

Next, we compared platelet integrin activation and  $\alpha$ -granule secretion (P-selectin exposure) between GRK2<sup>-/-</sup> and WT mice to further examine the effect of GRK2 deletion on platelet activity. Using flow cytometry to measure the activated integrin  $\alpha_{IIb}\beta_3$  and P-selectin in highly diluted platelets with their specific antibodies (Jon/A and anti-P-selectin, respectively), we observed significantly increased integrin  $\alpha_{IIb}\beta_3$  activation and P-selectin exposure in GRK2<sup>-/-</sup> platelets in response to ADP stimulation. There was a slightly increased P-selectin exposure in GRK2<sup>-/-</sup> platelets in response to U46619 (TxA<sub>2</sub> analogue) (Figure 3A; supplemental Figure 4). Platelet activation in response to PAR4 thrombin receptor agonist peptide (PAR4-AP), AYPGKF, and convulxin (glycoprotein

VI ligand) was normal (Figure 3A). In addition, no increase in basal P-selectin exposure or integrin activation on GRK2<sup>-/-</sup> platelets was noted, suggesting that the GRK2<sup>-/-</sup> platelets are not circulating in a preactivated state.


A platelet aggregation assay further verified this result, showing that GRK2<sup>-/-</sup> mouse platelets had an increased response to ADP stimulation (Figure 3B). Because other agonists rely on feedback from ADP signaling for maximal effect, we next performed platelet aggregation in response to PAR4-AP stimulation in the absence or presence of apyrase. There was an increase in platelet aggregation in response to PAR4-AP stimulation alone (Figure 3B; supplemental Figure 5A). However, the increased aggregation in GRK2<sup>-/-</sup> mouse platelets was abolished when secreted ADP was removed by treatment with 1 U/mL apyrase (Figure 3B; supplemental Figure 5B). Furthermore, the response to PAR4-AP stimulation was shifted to higher concentrations in the presence of apyrase, which is consistent with previous reports.<sup>18-20</sup> Conversely, GRK2<sup>-/-</sup> platelet aggregation was normal in response to U46619 or convulxin stimulation (supplemental Figure 5C-D). We also used single end point luminescence assays<sup>6</sup> to measure ATP release in platelets. There was no difference in ATP release between WT and GRK2<sup>-/-</sup> platelets in response to ADP, indicating that deletion of GRK2 in platelets did not affect dense granule secretion after ADP stimulation (supplemental Figure 6). Together, these observations suggest the specific targeting of GRK2 to the ADP-dependent signaling pathway.

### GRK2 affects ADP signaling by receptor desensitization

GRK2 is known to modulate GPCR signaling pathways at the receptor level via phosphorylation-dependent  $\beta$ -arrestin recruitment and receptor endocytosis.<sup>8</sup> To determine if GRK2 regulates ADP signaling through receptor desensitization, we measured platelet aggregation in response to ADP restimulation in WT and GRK2<sup>-/-</sup> mouse platelets under incubation or stirring conditions. After pre-incubation with 5  $\mu$ M ADP for 3 minutes, stimulation with 5  $\mu$ M ADP induced a greater aggregation response in GRK2<sup>-/-</sup> platelets than in WT platelets (Figure 3Ci). The second stimulation with 5  $\mu$ M ADP after pre-challenge also induced a greater aggregation response in GRK2<sup>-/-</sup> mouse platelets than in WT platelets (Figure 3Cii). ADP rechallenge caused a significant decrease in the maximum aggregation in WT platelets, whereas there was only a small decrease in GRK2<sup>-/-</sup> platelets after ADP restimulation (supplemental Figure 7). Together, these data suggest that GRK2 may regulate ADP receptors through the canonical mechanism of desensitization.

### GRK2 deletion in mouse platelets affects P2Y<sub>1</sub> receptor signaling pathways ex vivo

Human and mouse platelets express 2 distinct ADP receptors, P2Y<sub>1</sub> and P2Y<sub>12</sub>. Coactivation of these 2 receptors is essential for ADP-induced platelet activation.<sup>21</sup> P2Y<sub>1</sub> couples to G<sub>q</sub> and P2Y<sub>12</sub> couples to G<sub>i</sub> family members, primarily G<sub>i2</sub>. P2Y<sub>1</sub> activation leads to the release of Ca<sup>2+</sup> from intracellular stores followed by the influx of extracellular Ca<sup>2+</sup>.<sup>22,23</sup> P2Y<sub>1</sub>-dependent signaling mediates platelet shape change and transient, rapidly reversible aggregation induced by ADP. P2Y<sub>12</sub> is critically involved in integrin activation, inhibition of cyclic adenosine monophosphate (cAMP) formation, and stabilization of platelet aggregation.<sup>24</sup>



**Figure 2. Increased platelet accumulation at the site of vascular injury in GRK2<sup>-/-</sup> mice.** (Ai) Representative images of hemostatic plugs formed in WT and GRK2<sup>-/-</sup> mice 1.5 minutes after injury. (Aii) Confocal intravital fluorescence microscopy was performed to follow platelet accumulation (CD41) and P-selectin expression over 3 minutes. (Aiii) Platelet accumulation at the site of injury as measured by CD41 (area under the curve [AUC]) for 61 injuries in 7 WT mice and 41 injuries in 7 GRK2<sup>-/-</sup> mice. (Aiv) P-selectin positivity expressed as AUC. Forty-three injuries in 7 WT mice and 35 injuries in 7 GRK2<sup>-/-</sup> mice. (Bi) Fibrin deposition was examined after making small penetrating injuries in the cremaster muscle arterioles with a laser in GRK2<sup>-/-</sup> mice and littermate control mice. Thirty injuries in 7 WT mice and 29 injuries in 7 GRK2<sup>-/-</sup> mice. (Bii) Fibrin accumulation as measured by the mean fibrin area at the end of the 3-minute interval. Data sets were compared by using a two-by-two unbalanced analysis of variance. Data are presented as mean  $\pm$  standard error of the mean.

To assess the role of GRK2 in P2Y<sub>1</sub>-mediated platelet activation, we first studied activation of WT platelets vs GRK2<sup>-/-</sup> platelets in response to MRS2365, an agonist specific to P2Y<sub>1</sub>, by flow cytometry.<sup>25,26</sup> There was an increase in integrin activation and an increasing trend in P-selectin exposure in GRK2<sup>-/-</sup> platelets compared with WT platelets after stimulation with MRS2365 (Figure 4A; supplemental Figure 8). We next measured intracellular Ca<sup>2+</sup> release in WT and GRK2<sup>-/-</sup> mouse platelets in

response to ADP or MRS2365. It is worth noting that Ca<sup>2+</sup> mobilization in response to ADP is a readout for P2Y<sub>1</sub>- and not for P2Y<sub>12</sub>-mediated signaling.<sup>3</sup> There was a greater increase in intracellular Ca<sup>2+</sup> concentration in the GRK2<sup>-/-</sup> platelets relative to WT platelets, after stimulation with ADP or MRS2365 (Figure 4B). Taken together, these results suggest that GRK2 negatively regulates P2Y<sub>1</sub>-mediated signaling during platelet activation.




Figure 3.

## GRK2 deletion in mouse platelets affects P2Y<sub>12</sub> receptor signaling pathways *ex vivo*

To determine the role of GRK2 in P2Y<sub>12</sub> receptor signaling, we investigated several P2Y<sub>12</sub>-dependent signaling pathways in WT and GRK2<sup>-/-</sup> platelets. First, we studied the activation of the small GTPase RAS-related protein 1 (RAP1) in WT and GRK2<sup>-/-</sup> platelets. RAP1 activation is one of the critical steps leading to integrin  $\alpha_{IIb}\beta_3$  activation in platelets.<sup>27,28</sup> P2Y<sub>12</sub>-mediated signaling plays a critical role in maintaining sustained RAP1 activation.<sup>29</sup> Our data showed no increase in basal RAP1-GTP level in GRK2<sup>-/-</sup> platelets compared with that of WT mice. In contrast, there was a significantly greater increase in RAP1-GTP levels in response to ADP stimulation in GRK2<sup>-/-</sup> platelets than in controls (Figure 5A).

P2Y<sub>12</sub> signaling in platelets triggers G<sub>i</sub>-mediated activation of the serine-threonine kinases Akt1 and Akt2.<sup>30,31</sup> Thus, we compared phosphorylation of Akt (p-Akt) between ADP-stimulated WT and GRK2<sup>-/-</sup> mouse platelets in the presence or absence of MRS2500 (P2Y<sub>1</sub> antagonist) or cangrelor (P2Y<sub>12</sub> antagonist). GRK2<sup>-/-</sup> platelets showed a significant increase in p-Akt upon ADP stimulation (Figure 5B), and this difference persisted when blocking P2Y<sub>1</sub> receptor with MRS2500. However, the p-Akt level in both WT and GRK2<sup>-/-</sup> platelets was abolished when platelets were incubated with the P2Y<sub>12</sub> receptor antagonist cangrelor. Interestingly, in the presence of MRS2500, there was a small reduction in p-Akt after ADP stimulation, which may be due to indirect contributions from P2Y<sub>12</sub>-independent pathways.

Endothelium-derived prostaglandin (PGI<sub>2</sub>) targets the prostacyclin receptor, resulting in an increase in cAMP to prevent premature platelet activation. During platelet activation, cAMP production is suppressed by ADP-triggered P2Y<sub>12</sub> signaling via G<sub>i2 $\alpha$</sub> . To examine whether GRK2 deletion affects P2Y<sub>12</sub> inhibition of cAMP accumulation, we measured the cAMP level in both WT and GRK2<sup>-/-</sup> platelets under resting and PGI<sub>2</sub> conditions with or without ADP stimulation. These data show that there was no difference in basal and PGI<sub>2</sub>-induced cAMP formation in WT controls vs GRK2<sup>-/-</sup> platelets in the absence of ADP stimulation. In contrast, a slight but significant reduction in cAMP formation was observed in GRK2<sup>-/-</sup> platelets compared with the controls after ADP stimulation (Figure 5C). Collectively, these results show that GRK2 negatively regulates P2Y<sub>12</sub>-mediated signaling pathway in platelets.

## GRK2 regulates P2Y<sub>12</sub> signaling during hemostatic plug formation *in vivo*

Our prior studies show the importance of ADP/P2Y<sub>12</sub> signaling for platelet recruitment and/or retention in the outer shell region of hemostatic plugs.<sup>15,16</sup> We therefore hypothesize that blocking P2Y<sub>12</sub> signaling would eliminate the observed increased platelet accumulation at the outer layer of the thrombus in GRK2<sup>-/-</sup> mice.

To test this hypothesis, we examined thrombus formation after laser-induced injury to the cremaster muscle arterioles in WT and GRK2<sup>-/-</sup> mice treated with a P2Y<sub>12</sub> antagonist (cangrelor). Cangrelor or vehicle (saline) was administered IV immediately before each laser injury, as described in the Materials and methods. Platelet accumulation in saline-treated groups displayed a pattern similar to that shown in Figure 2Aii. Cangrelor treatment resulted in a reduction in total platelet accumulation at the site of vascular injury (Figure 6A; supplemental Figure 9). The rates of platelet accumulation in WT vs GRK2<sup>-/-</sup> in the presence of cangrelor were virtually identical. Treatment with cangrelor did not attenuate formation of the hemostatic plug core region of P-selectin-positive platelets in both WT and GRK2<sup>-/-</sup> mice (supplemental Figure 10A). These results indicate that blocking P2Y<sub>12</sub> receptors with cangrelor eliminates the difference in platelet accumulation between GRK2<sup>-/-</sup> mice and control mice at the site of vascular injury.

In addition, the hemostatic response was measured by using a tail bleeding assay.<sup>32</sup> GRK2<sup>-/-</sup> mice had a significantly shorter bleeding time compared with control mice (Figure 6B). Collectively, these data suggest that GRK2 regulates hemostatic response *in vivo*.


## Absence of GRK2 accelerates pulmonary thromboembolism and promotes thrombosis in FeCl<sub>3</sub>-induced carotid artery

Next, we examined the effects of GRK2 deficiency on platelet function using a pulmonary thromboembolism mouse model, a thrombosis model in which platelet activation is the most prominent feature.<sup>33</sup> To determine whether deletion of GRK2 might increase the incidence of formation of pulmonary microthrombi induced by platelet activation, GRK2<sup>-/-</sup> and their WT control littermates were injected IV with 0.3 mg/g ADP. ADP has been previously shown to induce disseminated pulmonary thromboembolism.<sup>34,35</sup> The result indicates that GRK2<sup>-/-</sup> mice had more platelet accumulation in the lungs than observed in the WT control mice (Figure 6Ci-ii). There was no difference in mortality rate between WT and GRK2<sup>-/-</sup> mice (supplemental Figure 10B). Blocking P2Y<sub>12</sub> receptors with clopidogrel abrogated ADP-induced platelet activation and abolished the difference in platelet accumulation in the lungs (Figure 6C; supplemental Figure 10C). These results suggest that GRK2 plays an important role in limiting platelet accumulation in the lung.

To further determine the role of GRK2 in thrombosis, we performed a FeCl<sub>3</sub>-induced carotid artery injury model.<sup>36</sup> Five of 6 GRK2<sup>-/-</sup> mice reached stable occlusion after 7.5% FeCl<sub>3</sub>-induced vascular injury. The WT control mice either did not occlude at all or had an unstable occlusion during the 30 minutes of the experiment (Figure 6D). These results indicate that GRK2 plays an important role in regulating thrombosis.

**Figure 3 (continued) Increased platelet integrin activation,  $\alpha$ -granule secretion, aggregation, and decreased receptor desensitization in GRK2<sup>-/-</sup> mice under ADP stimulation.** (A) Platelets from GRK2<sup>-/-</sup> and littermate control (WT) mice were stained with fluorophore-conjugated antibodies to either activated integrin  $\alpha_{IIb}\beta_3$  (Jon/A antibody) or P-selectin and measured by flow cytometry. Platelets were stimulated with ADP, TxA<sub>2</sub> mimetic (U46619), PAR4-AP (AYPGKF), or convulxin at the concentrations indicated (n = 4). (B) Platelet aggregation in response to a PAR4-AP or ADP at the concentrations indicated (n  $\geq$  3). (C) Receptor desensitization measured by platelet aggregation in response to ADP. (Ci) Platelets from GRK2<sup>-/-</sup> and littermate WT control mice were stimulated with 5  $\mu$ M ADP after pre-incubation with 5  $\mu$ M ADP. (Cii) Platelets from GRK2<sup>-/-</sup> and littermate WT control mice were stimulated with 5  $\mu$ M ADP first and restimulated with 5  $\mu$ M ADP after a 40-minute interval. n = 4. P  $\leq$  .05 was considered statistically significant. MFI, mean fluorescence intensity; ns, not significant.






**Figure 4. Increased platelet activation and calcium mobilization downstream of P2Y<sub>1</sub>-dependent signaling in GRK2<sup>-/-</sup> platelets.** (A) Platelet activation. Platelets from GRK2<sup>-/-</sup> and littermate control mice (WT) were stained with fluorophore-conjugated antibodies to either activated integrin  $\alpha_{IIb}\beta_3$  (Jon/A antibody) or P-selectin and measured by flow cytometry. Platelets were stimulated with MRS2365 (MRS).  $n = 4$ . (B) Calcium mobilization. Mouse platelets were loaded with fura-2 AM (10  $\mu$ M) and stimulated with ADP or MRS at the concentrations indicated in the absence of extracellular Ca<sup>2+</sup>. Representative measurements are shown. The results of 6 experiments, ADP (upper), and the results of 4 experiments, MRS (lower), are summarized (mean  $\pm$  standard error of the mean). MFI, mean fluorescence intensity; ns, not significant.

### GRK2 does not bind with the PAR1 receptor in thrombin-activated human platelets

To examine whether GRK2 regulates thrombin signaling through a canonical interaction with PAR1, we attempted to immunoprecipitate GRK2 with anti-PAR1 antibody from either resting or thrombin-stimulated human platelets. In contrast to the increased binding of

GRK6 to PAR1 as we previously reported (supplemental Figure 11A),<sup>3</sup> PAR1 receptor does not bind GRK2 in activated platelets (supplemental Figure 11B). This finding is consistent with the functional readouts from flow cytometry and platelet aggregation assays in which deletion of GRK2 in mouse platelets does not affect PAR-dependent signaling.




**Figure 5. Increased RAP1 activation, Akt phosphorylation, and reduced cAMP formation downstream of P2Y<sub>12</sub>-dependent signaling in GRK2<sup>-/-</sup> platelets.**

(A) RAP1 activation. Ai: RAP1 protein expression in WT and GRK2<sup>-/-</sup> platelets. Aii: RAP1-GTP levels were measured in platelets that were either unstimulated or stimulated with ADP for 5 minutes (mean ± standard error of the mean; n = 4). (B) Akt phosphorylation. Gel-filtered platelets from GRK2<sup>-/-</sup> mice or matched WT controls were lysed directly or incubated with 10 μM ADP for 5 minutes in the presence or absence of the P2Y<sub>1</sub> antagonist MRS2500 (50 mM) or the P2Y<sub>12</sub> antagonist cangrelor (100 nM), as indicated. (Bi) Lysates were probed with anti-pAkt (S473) and reprobbed with anti-Akt antibody. (Bii) The p-Akt signal was normalized to the Akt loading control and is represented as signal relative to that of the WT with or without antagonist MRS2500 (mean ± standard error of the mean; n = 6). (C) cAMP formation. cAMP levels in resting platelets and platelets stimulated with PGI<sub>2</sub> in the presence or absence of ADP at the final concentrations indicated (mean ± standard error of the mean; n = 3). ns, not significant.


### Endogenous GRK2 in human platelets interacts with G<sub>βγ</sub> subunits but not G<sub>qα</sub>

As shown previously in nucleated cells, the RGS domain of GRK2 can bind with G<sub>qα</sub> subunits, but not G<sub>iα</sub> subunits, in vitro and therefore inhibit G<sub>q</sub>-mediated signaling.<sup>37</sup> We observed the interaction between GRK2 and G<sub>qα</sub> in human and mouse platelets using an in vitro glutathione S-transferase-GRK2 pull-down assay (Figure 7Ai). There was no binding of GRK2 to G<sub>i2α</sub>. We also

tested whether GRK2 interacts with endogenous G<sub>qα</sub> in platelets. GRK2 was precipitated from resting and thrombin-activated human platelets with an anti-GRK2 antibody, and samples were probed for G<sub>qα</sub>. No binding of GRK2 to endogenous G<sub>qα</sub> was detected by immunoprecipitation (supplemental Figure 12A), even in the presence AIF<sub>4</sub><sup>-</sup> to mimic the GTP hydrolysis transition state of G<sub>qα</sub> (Figure 7Aii), suggesting that G<sub>qα</sub> is not a direct target of negative regulation by GRK2 in platelets. Furthermore, membrane association



**Figure 6. Deletion of GRK2 in platelets causes increased platelet accumulation in the shell region, shortens tail bleeding time, and enhances thrombosis. P2Y<sub>12</sub> inhibition eliminates the phenotypic difference at the site of injury.** (A) Laser-induced thrombus formation in WT and GRK2<sup>-/-</sup> mice treated with the P2Y<sub>12</sub> antagonist (cangrelor). Cangrelor or vehicle (saline) was administered by IV immediately before each of the laser injuries. (Ai) Confocal intravital fluorescence microscopy was then performed to follow platelet accumulation (CD41) over 3 minutes. (Aii) Platelet accumulation at the site of injury as measured by CD41 of 46 injuries in 3 WT mice treated with saline, 52 injuries in 4 GRK2<sup>-/-</sup> mice treated with saline, 44 injuries in 4 WT mice treated with cangrelor, and 39 injuries in 4 GRK2<sup>-/-</sup> mice treated with cangrelor. Data sets were compared by using an unbalanced two-way analysis of variance with Tukey's post hoc test. (B) Hemostatic response was measured by tail bleeding. Tails of anesthetized mice were transected and the time to complete arrest of bleeding (as defined by no bleeding recurring within 2 minutes) was recorded for each mouse.  $n = 9$  mice per genotype. (C) Representative images of anti-glycoprotein IX (GPIX)-labeled thrombi in lungs harvested from WT and GRK2<sup>-/-</sup> mice treated with ADP in the absence or presence of clopidogrel (P2Y<sub>12</sub> antagonist). (Cii and Ciii) Thrombosis scores, representing the mean thrombus area and number of thrombi, of 5 WT mice vs 5 GRK2<sup>-/-</sup> mice are summarized in each group.  $P \leq .05$  was considered statistically significant. (D) The carotid arteries of WT and GRK2<sup>-/-</sup> mice were injured by 7.5% FeCl<sub>3</sub>. Flow rates through the carotid artery were measured with a Doppler flow probe after vascular injury. (Di) Quantitation of the percentage of mice that formed a stable clot, unstable clot, or no occlusion. A  $\chi^2$  test was conducted, producing a  $P$  value of .011. (Dii) The probability of forming unstable occlusion at a respective time interval for WT or GRK2<sup>-/-</sup> mice. Experiments were terminated at 30 minutes. Kaplan-Meier curves were analyzed by using the log-rank test.  $n = 6$  mice per genotype. Data are presented as mean  $\pm$  standard error of the mean. AUC, area under the curve. ns, not significant.



**Figure 7.**  $G_{\beta}$  interacts with endogenous GRK2 in platelets ex vivo but  $G_{\alpha q}$  does not. (Ai) GST-GRK2 was immunoprecipitated from human or mouse platelets, and samples were probed for  $G_{\alpha q}$  or  $G_{i\alpha 2}$ . Glutathione *S*-transferase (GST) reprobe indicated the equal loading.  $n = 2$ . (Aii) Human platelet lysate from resting or 1 U/mL thrombin-stimulated platelets in the presence or absence of GDP+ $AlF_4^-$  were subjected to immunoprecipitation of GRK2 and probed for  $G_{\alpha q}$ .  $n = 3$ . (B) GRK2 from resting and thrombin-stimulated (1 U/mL) human platelet lysate was immunoprecipitated with the specific mouse anti-GRK2 antibody and probed for  $G_{\beta}$ . GRK2 was reprobated with an anti-GRK2 antibody.  $n = 6$ . A marker line was excised as indicated by the vertical line.

of GRK2 is mediated by binding to  $G_{\beta\gamma}$  via its pleckstrin homology domain in nucleated cells, and this binding is critical for the activation of GRK2.<sup>38,39</sup> We observed that GRK2 binds to subunits of  $G_{\beta\gamma}$  during platelet activation in response to thrombin stimulation but not to 2MesADP (Figure 7B; supplemental Figure 12B).

### Pharmacologic inhibition of GRK2 in human platelets

We next investigated the effect of inhibiting GRK2 activity on platelet activation using a GRK2-specific inhibitor, Compound 101 (Cmpd101). Cmpd101 is a potent and selective GRK2 inhibitor and has been widely used in studying the roles of GRK2 in GPCR desensitization.<sup>40</sup> The 50% inhibitory concentration value of Cmpd101 to inhibit GRK2-mediated bovine rod outer segment (bROS) phosphorylation is 290 nM. Treatment of human platelets with Cmpd101 results in an increase in platelet activation in response to 2MesADP (supplemental Figure 13), suggesting that GRK2 activity is required in ADP-mediated signaling in human platelets.

### Discussion

Dysregulated platelet activation and thrombus instability can lead to venous thrombosis, myocardial infarction, and stroke.<sup>41</sup> Considerable efforts have been made to treat thrombotic disorders by selectively inhibiting the pathways downstream of the ADP-P2Y<sub>12</sub> signaling pathway.<sup>42-44</sup> To avoid the risk of premature platelet activation or excess platelet accumulation at sites of injury, several regulatory events are in place to limit the rate and/or extent of platelet activation, including regulators that can limit P2Y<sub>12</sub>-mediated platelet activation.<sup>29</sup> Although GRK2 is a critical regulator of GPCR signaling in the heart, its contribution to platelet activation and its role in GPCR-dependent signaling in platelets are still poorly understood.

In this study, we asked if GRK2 functions as a checkpoint to limit the intensity and duration of platelet activation downstream of GPCRs. Taking advantage of a GRK2<sup>-/-</sup> mouse line in which GRK2 is specifically deleted in megakaryocyte/platelets, we have identified a unique regulatory role of GRK2 in platelets downstream of ADP signaling. Among all the agonists studied in platelet functional assays, only ADP stimulation causes an increase in GRK2<sup>-/-</sup> platelet activation over WT, suggesting a dispensable role of GRK2

in thrombin/PAR4- or TxA<sub>2</sub>/TP-mediated signaling. Human platelets express 150 copies of P2Y<sub>1</sub> receptor and 400 copies of P2Y<sub>12</sub> receptor per cell, respectively.<sup>45</sup> In comparison, mouse platelets express 1000 copies of P2Y<sub>1</sub> receptor and 2700 copies of P2Y<sub>12</sub> receptor per cell.<sup>46</sup> Both human and mouse platelets express higher levels of P2Y<sub>12</sub> than that of P2Y<sub>1</sub>. Our studies show that deletion of GRK2 in mouse platelets leads to an increase in ADP-induced platelet activation, suggesting a critical role for GRK2 in ADP P2Y<sub>12</sub>-dependent signaling. These observations are consistent with the previous findings by Hardy et al<sup>11</sup> that desensitization of P2Y<sub>12</sub> is mediated by GRK2. In addition, they show that desensitization of P2Y<sub>1</sub> receptor in 1321N1 human astrocytoma cells is mainly regulated by protein kinase C but not by GRK2. In contrast to their findings in 1321N1 cells, our studies show that knocking out GRK2 in platelets causes an increase in P2Y<sub>1</sub>-dependent platelet activation. These findings indicate that GRK2 negatively regulates P2Y<sub>1</sub>-mediated signaling in platelets. It is worth noting that 1321N1 cells express similar levels of P2Y<sub>1</sub> and P2Y<sub>12</sub> receptors,<sup>11</sup> indicating that the regulation of P2Y<sub>1</sub> by protein kinase C in nucleated cells is presumably not operative in anucleate platelets. Our results are consistent with a recent report showing that platelets expressing a mutant P2Y<sub>1</sub> receptor that does not desensitize have increased platelet activation in response to ADP and MRS2365 *in vitro*. Interestingly, impaired P2Y<sub>1</sub> receptor desensitization had no effect on *in vivo* hemostasis or thrombosis (Paul et al., ISTH 2021).<sup>47</sup> In addition, recent studies using platelets deficient in arrestin-3 indicate that arrestin-3 negatively regulates platelet activation downstream of P2Y receptors, which is consistent with what we have observed about the regulatory roles of GRK2 in ADP/P2Y signaling.<sup>48,49</sup>

To further test whether the loss of GRK2 produces a pathway-selective gain of function *in vivo*, we investigated the consequence of GRK2 deficiency on thrombus formation using several injury models. In the laser-induced cremaster arteriole injury model, thrombin is the primary driver for platelet activation in the core of hemostatic thrombi, whereas ADP and TxA<sub>2</sub> are the main drivers for platelet accumulation in the shell.<sup>15</sup> Hemostatic plugs formed in GRK2<sup>-/-</sup> mice have a larger shell region, with no increase in the size of the core region. In addition, GRK2<sup>-/-</sup> mice have more platelet accumulation in the lungs after ADP stimulation than observed in the WT control mice. Pharmacologic inhibition of P2Y<sub>12</sub> signaling with cangrelor or clopidogrel abrogates the phenotypic difference between WT control and GRK2<sup>-/-</sup> mice. These findings suggest that the enhanced platelet accumulation observed in GRK2<sup>-/-</sup> mice is primarily due to loss of GRK2-mediated regulation of P2Y<sub>12</sub> signaling. However, we are unable to examine direct interaction between GRK2 and P2Y<sub>12</sub> due to the lack of an effective P2Y<sub>12</sub> antibody, either for western blot analysis or for immunoprecipitation.

The multifunctionality of GRK2 has recently gained attention due to a constellation of effects, including both its canonical (GRK2-mediated phosphorylation) and noncanonical (kinase-independent molecular interactions) functions.<sup>50,51</sup> For example, it has been shown in nucleated cells that GRK2 is recruited to the membrane and binds activated G<sub>qα</sub> via its RGS domain.<sup>37,52</sup> However, the lack of endogenous binding of G<sub>qα</sub> by GRK2 in platelets is consistent with a previous report regarding the lack of binding of G<sub>qα</sub> by GRK2 in the

heart.<sup>37</sup> In addition, GRK2 contains a C-terminal pleckstrin homology domain, which allows for G<sub>βγ</sub> binding after dissociation of the heterotrimeric G protein. The binding of GRK2 with G<sub>βγ</sub> facilitates membrane targeting of GRK2, which also stimulates phosphorylation of activated GPCRs.<sup>53</sup> We were able to observe this interaction in activated platelets, and this interaction may have consequences for the activation of GRK2. In cells other than platelets, GRK2 has been reported to interact directly with phosphatidylinositol 3-kinase γ<sup>54,55</sup> and Akt.<sup>56</sup> These binding partners of GRK2 may indicate the noncanonical functions of GRK2 in regulating GPCR signaling, which might contribute to the observed distinct functions of GRK2 from other GRKs, such as GRK5 and GRK6. Elucidation of novel GRK2 substrates, other GRK2-mediated signaling events, and GRK2's noncanonical functions and binding partners in platelets is worthy of investigation in a future study.

In summary, we have shown for the first time that GRK2 regulates the hemostatic response by limiting platelet activation via ADP P2Y<sub>1</sub>- and P2Y<sub>12</sub>-dependent signaling. The unique regulatory role of GRK2 in ADP signaling in platelets is highly relevant to recent efforts in developing small molecule GRK2 inhibitors in the treatment of heart disease. Our finding regarding the role of GRK2 in platelets highlights the potential for significant off-target effects of GRK2 inhibition, such as platelet hyperactivity and subsequent thrombosis, due to the loss of GRK2-mediated rate-limiting effects on ADP signaling in platelets. Thus, it emphasizes the benefits of maintaining a certain level of GRK2 activity in platelets in preventing thrombotic events. The efforts and knowledge gained in this study should enable better development of GRK2-based therapeutic options for the treatment of cardiovascular diseases and thrombotic disorders.

## Acknowledgments

The authors thank Xi Chen for technical assistance.

These studies were supported by the National Institutes of Health, National Heart, Lung, and Blood Institute (R01HL144574, P.M.).

## Authorship

Contributions: X.Z., M.C., J.V.M., Y.Y., and A.B. performed experiments; X.Z., M.C., J.K.C., J.V.M., W.J.K., S.E.M., M.T., T.J.S., L.Z., and P.M. designed experiments; and X.Z., and P.M. wrote the paper.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: X.Z., 0000-0002-8651-6248; Y.Y., 0000-0002-7584-6903; W.J.K., 0000-0002-8522-530X; P.M., 0000-0002-5486-8695.

Correspondence: Peisong Ma, Cardeza Foundation for Hematologic Research, Department of Medicine, Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, PA 20036; e-mail: Peisong.Ma@jefferson.edu; and Li Zhu, Cyrus Tang Hematology Center, Soochow University, Suzhou, China; e-mail: zhul@suda.edu.cn.

## References

1. Offermanns S. Activation of platelet function through G protein-coupled receptors. *Circ Res.* 2006;99(12):1293-1304.
2. Brass LF, Newman DK, Wannemacher KM, Zhu L, Stalker TJ. Signal transduction during platelet plug formation. In: Michelson AD, ed. *Platelets*. New York, NY: Elsevier; 2013:367-398.
3. Chen X, Gupta S, Cooper M, et al. GRK6 regulates the hemostatic response to injury through its rate-limiting effects on GPCR signaling in platelets. *Blood Adv.* 2020;4(1):76-86.
4. Chaudhary PK, Kim S, Jee Y, Lee SH, Park KM, Kim S. Role of GRK6 in the regulation of platelet activation through selective G protein-coupled receptor (GPCR) desensitization. *Int J Mol Sci.* 2020;21(11):E3932.
5. Chen X, Zhao X, Cooper M, Ma P. The roles of GRKs in hemostasis and thrombosis. *Int J Mol Sci.* 2020;21(15):E5345.
6. Downes K, Zhao X, Gleadall NS, et al. G protein-coupled receptor kinase 5 regulates thrombin signaling in platelets via PAR-1. *Blood Adv.* 2022; 6(7):2319-2330.
7. Rodriguez BAT, Bhan A, Beswick A, et al; FinnGen Study. A platelet function modulator of thrombin activation is causally linked to cardiovascular disease and affects PAR4 receptor signaling. *Am J Hum Genet.* 2020;107(2):211-221.
8. Ribas C, Penela P, Murga C, et al. The G protein-coupled receptor kinase (GRK) interactome: role of GRKs in GPCR regulation and signaling. *Biochim Biophys Acta.* 2007;1768(4):913-922.
9. Ungerer M, Böhm M, Elce JS, Erdmann E, Lohse MJ. Altered expression of beta-adrenergic receptor kinase and beta 1-adrenergic receptors in the failing human heart. *Circulation.* 1993;87(2):454-463.
10. Ungerer M, Kessebohm K, Kronsbein K, Lohse MJ, Richardt G. Activation of beta-adrenergic receptor kinase during myocardial ischemia. *Circ Res.* 1996;79(3):455-460.
11. Hardy AR, Conley PB, Luo J, Benovic JL, Poole AW, Mundell SJ. P2Y1 and P2Y12 receptors for ADP desensitize by distinct kinase-dependent mechanisms. *Blood.* 2005;105(9):3552-3560.
12. Jaber M, Koch WJ, Rockman H, et al. Essential role of beta-adrenergic receptor kinase 1 in cardiac development and function. *Proc Natl Acad Sci USA.* 1996;93(23):12974-12979.
13. Matkovich SJ, Diwan A, Klanke JL, et al. Cardiac-specific ablation of G-protein receptor kinase 2 redefines its roles in heart development and beta-adrenergic signaling. *Circ Res.* 2006;99(9):996-1003.
14. Tiedt R, Schomber T, Hao-Shen H, Skoda RC. Pf4-Cre transgenic mice allow the generation of lineage-restricted gene knockouts for studying megakaryocyte and platelet function in vivo. *Blood.* 2007;109(4):1503-1506.
15. Stalker TJ, Traxler EA, Wu J, et al. Hierarchical organization in the hemostatic response and its relationship to the platelet-signaling network. *Blood.* 2013;121(10):1875-1885.
16. Shen J, Sampietro S, Wu J, et al. Coordination of platelet agonist signaling during the hemostatic response in vivo. *Blood Adv.* 2017;1(27): 2767-2775.
17. Stalker TJ. Mouse laser injury models: variations on a theme. *Platelets.* 2020;31(4):423-431.
18. Foster CJ, Prosser DM, Agans JM, et al. Molecular identification and characterization of the platelet ADP receptor targeted by thienopyridine antithrombotic drugs. *J Clin Invest.* 2001;107(12):1591-1598.
19. Hollopeter G, Jantzen HM, Vincent D, et al. Identification of the platelet ADP receptor targeted by antithrombotic drugs. *Nature.* 2001;409(6817): 202-207.
20. Zhang FL, Luo L, Gustafson E, et al. ADP is the cognate ligand for the orphan G protein-coupled receptor SP1999. *J Biol Chem.* 2001;276(11): 8608-8615.
21. Jin J, Kunapuli SP. Coactivation of two different G protein-coupled receptors is essential for ADP-induced platelet aggregation. *Proc Natl Acad Sci USA.* 1998;95(14):8070-8074.
22. Varga-Szabo D, Braun A, Nieswandt B. Calcium signaling in platelets. *J Thromb Haemost.* 2009;7(7):1057-1066.
23. Ma P, Gupta S, Sampietro S, et al. RGS10 shapes the hemostatic response to injury through its differential effects on intracellular signaling by platelet agonists. *Blood Adv.* 2018;2(16):2145-2155.
24. Dorsam RT, Kunapuli SP. Central role of the P2Y12 receptor in platelet activation. *J Clin Invest.* 2004;113(3):340-345.
25. Chhatriwala M, Ravi RG, Patel RI, Boyer JL, Jacobson KA, Harden TK. Induction of novel agonist selectivity for the ADP-activated P2Y1 receptor versus the ADP-activated P2Y12 and P2Y13 receptors by conformational constraint of an ADP analog. *J Pharmacol Exp Ther.* 2004;311(3): 1038-1043.
26. Battram AM, Durrant TN, Agbani EO, et al. The phosphatidylinositol 3,4,5-trisphosphate (PI(3,4,5)P3) binder Rasa3 regulates phosphoinositide 3-kinase (PI3K)-dependent integrin  $\alpha\text{IIb}\beta\text{3}$  outside-in signaling. *J Biol Chem.* 2017;292(5):1691-1704.
27. Woulfe D, Jiang H, Mortensen R, Yang J, Brass LF. Activation of Rap1B by G(i) family members in platelets. *J Biol Chem.* 2002;277(26): 23382-23390.
28. Chrzanowska-Wodnicka M, Smyth SS, Schoenwaelder SM, Fischer TH, White GC II. Rap1b is required for normal platelet function and hemostasis in mice. *J Clin Invest.* 2005;115(3):680-687.

29. Stefanini L, Paul DS, Robledo RF, et al. RASA3 is a critical inhibitor of RAP1-dependent platelet activation. *J Clin Invest*. 2015;125(4):1419-1432.
30. Kim S, Jin J, Kunapuli SP. Akt activation in platelets depends on Gi signaling pathways. *J Biol Chem*. 2004;279(6):4186-4195.
31. Woulfe D, Jiang H, Morgans A, Monks R, Birnbaum M, Brass LF. Defects in secretion, aggregation, and thrombus formation in platelets from mice lacking Akt2. *J Clin Invest*. 2004;113(3):441-450.
32. Nagy Z, Mori J, Ivanova VS, Mazharian A, Senis YA. Interplay between the tyrosine kinases Chk and Csk and phosphatase PTPRJ is critical for regulating platelets in mice. *Blood*. 2020;135(18):1574-1587.
33. Angelillo-Scherrer A, de Frutos P, Aparicio C, et al. Deficiency or inhibition of Gas6 causes platelet dysfunction and protects mice against thrombosis. *Nat Med*. 2001;7(2):215-221.
34. Fabre JE, Nguyen M, Latour A, et al. Decreased platelet aggregation, increased bleeding time and resistance to thromboembolism in P2Y1-deficient mice. *Nat Med*. 1999;5(10):1199-1202.
35. Yang J, Wu J, Kowalska MA, et al. Loss of signaling through the G protein, Gz, results in abnormal platelet activation and altered responses to psychoactive drugs. *Proc Natl Acad Sci USA*. 2000;97(18):9984-9989.
36. Ma P, Cierniewska A, Signarvic R, et al. A newly identified complex of spinophilin and the tyrosine phosphatase, SHP-1, modulates platelet activation by regulating G protein-dependent signaling. *Blood*. 2012;119(8):1935-1945.
37. Schumacher SM, Gao E, Cohen M, Lieu M, Chuprun JK, Koch WJ. A peptide of the RGS domain of GRK2 binds and inhibits G $\alpha$ (q) to suppress pathological cardiac hypertrophy and dysfunction. *Sci Signal*. 2016;9(420):ra30.
38. Pitcher JA, Inglese J, Higgins JB, et al. Role of beta gamma subunits of G proteins in targeting the beta-adrenergic receptor kinase to membrane-bound receptors. *Science*. 1992;257(5074):1264-1267.
39. Koch WJ, Inglese J, Stone WC, Lefkowitz RJ. The binding site for the beta gamma subunits of heterotrimeric G proteins on the beta-adrenergic receptor kinase. *J Biol Chem*. 1993;268(11):8256-8260.
40. Thal DM, Yeow RY, Schoenau C, Huber J, Tesmer JJ. Molecular mechanism of selectivity among G protein-coupled receptor kinase 2 inhibitors. *Mol Pharmacol*. 2011;80(2):294-303.
41. Ruggeri ZM. Platelets in atherothrombosis. *Nat Med*. 2002;8(11):1227-1234.
42. Speich HE, Bhal V, Houser KH, et al. Signaling via P2Y12 may be critical for early stabilization of platelet aggregates. *J Cardiovasc Pharmacol*. 2014;63(6):520-527.
43. Trumel C, Payrastra B, Plantavid M, et al. A key role of adenosine diphosphate in the irreversible platelet aggregation induced by the PAR1-activating peptide through the late activation of phosphoinositide 3-kinase. *Blood*. 1999;94(12):4156-4165.
44. Eckly A, Gendraul JL, Hechler B, Cazenave JP, Gachet C. Differential involvement of the P2Y1 and P2YT receptors in the morphological changes of platelet aggregation. *Thromb Haemost*. 2001;85(4):694-701.
45. Gachet C, Hechler B. Platelet purinergic receptors in thrombosis and inflammation. *Hamostaseologie*. 2020;40(2):145-152.
46. Zeiler M, Moser M, Mann M. Copy number analysis of the murine platelet proteome spanning the complete abundance range. *Mol Cell Proteomics*. 2014;13(12):3435-3445.
47. Paul DS, Blatt T, Murcia S, et al. Loss of P2Y1 receptor desensitization does not impact in vivo hemostasis or thrombosis despite increased platelet reactivity in vitro. In: ISTH Congress; 17 July 2021; Philadelphia.
48. Chaudhary PK, Kim S, Kim S. The predominant role of arrestin3 in general GPCR desensitization in platelets. *J Clin Med*. 2021;10(20):4743.
49. Hutchinson JL, Zhao X, Hill R, Mundell SJ. Arrestin-3 differentially regulates platelet GPCR subsets. *Platelets*. 2020;31(5):641-645.
50. Murga C, Arcones AC, Cruces-Sande M, Briones AM, Salices M, Mayor F Jr. G protein-coupled receptor kinase 2 (GRK2) as a potential therapeutic target in cardiovascular and metabolic diseases. *Front Pharmacol*. 2019;10:112.
51. Pflieger J, Gresham K, Koch WJ. G protein-coupled receptor kinases as therapeutic targets in the heart. *Nat Rev Cardiol*. 2019;16(10):612-622.
52. Shankaranarayanan A, Thal DM, Tesmer VM, et al. Assembly of high order G alpha q-effector complexes with RGS proteins. *J Biol Chem*. 2008;283(50):34923-34934.
53. Kozasa T. The structure of GRK2-G beta gamma complex: intimate association of G-protein signaling modules. *Trends Pharmacol Sci*. 2004;25(2):61-63.
54. Naga Prasad SV, Barak LS, Rapacciuolo A, Caron MG, Rockman HA. Agonist-dependent recruitment of phosphoinositide 3-kinase to the membrane by beta-adrenergic receptor kinase 1. A role in receptor sequestration. *J Biol Chem*. 2001;276(22):18953-18959.
55. Schlegel P, Reinkober J, Meinhardt E, et al. G protein-coupled receptor kinase 2 promotes cardiac hypertrophy. *PLoS One*. 2017;12(7):e0182110.
56. Liu S, Premont RT, Kontos CD, Zhu S, Rockey DC. A crucial role for GRK2 in regulation of endothelial cell nitric oxide synthase function in portal hypertension. *Nat Med*. 2005;11(9):952-958.