

10-7-2023

## Identification of Post-cardiac Arrest Blood Pressure Thresholds Associated With Outcomes in Children: An ICU-Resuscitation Study

Monique M Gardner

David A Hehir

Ron W Reeder

Follow this and additional works at: <https://jdc.jefferson.edu/medfp>

Tageldin Ahmed

 Part of the [Critical Care Commons](#)

Michael J Bell

[Let us know how access to this document benefits you](#)

See next page for additional authors

### Recommended Citation

Gardner, Monique M; Hehir, David A; Reeder, Ron W; Ahmed, Tageldin; Bell, Michael J; Berg, Robert A; Bishop, Robert; Bochkoris, Matthew; Burns, Candice; Carcillo, Joseph A; Carpenter, Todd C; Dean, J Michael; Diddle, J Wesley; Federman, Myke; Fernandez, Richard; Fink, Ericka L; Franzon, Deborah; Frazier, Aisha H; Friess, Stuart H; Graham, Kathryn; Hall, Mark; Harding, Monica L; Horvat, Christopher M; Huard, Leanna L; Maa, Tensing; Manga, Arushi; McQuillen, Patrick S; Meert, Kathleen L; Morgan, Ryan W; Mourani, Peter M; Nadkarni, Vinay M; Naim, Maryam Y; Notterman, Daniel; Pollack, Murray M; Sapru, Anil; Schneiter, Carleen; Sharron, Matthew P; Srivastava, Neeraj; Tilford, Bradley; Viteri, Shirley D.; Wessel, David; Wolfe, Heather A; Yates, Andrew R; Zuppa, Athena F; Sutton, Robert M; and Topjian, Alexis A, "Identification of Post-cardiac Arrest Blood Pressure Thresholds Associated With Outcomes in Children: An ICU-Resuscitation Study" (2023). *Department of Medicine Faculty Papers*. Paper 425.  
<https://jdc.jefferson.edu/medfp/425>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Medicine Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

---

## Authors

Monique M Gardner, David A Hehir, Ron W Reeder, Tageldin Ahmed, Michael J Bell, Robert A Berg, Robert Bishop, Matthew Bochkoris, Candice Burns, Joseph A Carcillo, Todd C Carpenter, J Michael Dean, J Wesley Diddle, Myke Federman, Richard Fernandez, Ericka L Fink, Deborah Franzon, Aisha H Frazier, Stuart H Friess, Kathryn Graham, Mark Hall, Monica L Harding, Christopher M Horvat, Leanna L Huard, Tensing Maa, Arushi Manga, Patrick S McQuillen, Kathleen L Meert, Ryan W Morgan, Peter M Mourani, Vinay M Nadkarni, Maryam Y Naim, Daniel Notterman, Murray M Pollack, Anil Sapru, Carleen Schneiter, Matthew P Sharron, Neeraj Srivastava, Bradley Tilford, Shirley D. Viteri, David Wessel, Heather A Wolfe, Andrew R Yates, Athena F Zuppa, Robert M Sutton, and Alexis A Topjian

RESEARCH

Open Access



# Identification of post-cardiac arrest blood pressure thresholds associated with outcomes in children: an ICU-Resuscitation study

Monique M. Gardner<sup>1\*</sup>, David A. Hehir<sup>1</sup>, Ron W. Reeder<sup>2</sup>, Tageldin Ahmed<sup>3</sup>, Michael J. Bell<sup>4</sup>, Robert A. Berg<sup>1</sup>, Robert Bishop<sup>5</sup>, Matthew Bochkoris<sup>6</sup>, Candice Burns<sup>7</sup>, Joseph A. Carcillo<sup>6</sup>, Todd C. Carpenter<sup>5</sup>, J. Michael Dean<sup>2</sup>, J. Wesley Diddle<sup>1</sup>, Myke Federman<sup>8</sup>, Richard Fernandez<sup>9</sup>, Ericka L. Fink<sup>6</sup>, Deborah Franzone<sup>10</sup>, Aisha H. Frazier<sup>11</sup>, Stuart H. Friess<sup>12</sup>, Kathryn Graham<sup>1</sup>, Mark Hall<sup>9</sup>, Monica L. Harding<sup>2</sup>, Christopher M. Horvat<sup>6</sup>, Leanna L. Huard<sup>8</sup>, Tensing Maa<sup>9</sup>, Arushi Manga<sup>12</sup>, Patrick S. McQuillen<sup>10</sup>, Kathleen L. Meert<sup>3</sup>, Ryan W. Morgan<sup>1</sup>, Peter M. Mourani<sup>13</sup>, Vinay M. Nadkarni<sup>1</sup>, Maryam Y. Naim<sup>1</sup>, Daniel Notterman<sup>14</sup>, Murray M. Pollack<sup>4</sup>, Anil Sapru<sup>8</sup>, Carleen Schneider<sup>5</sup>, Matthew P. Sharron<sup>4</sup>, Neeraj Srivastava<sup>8</sup>, Bradley Tilford<sup>3</sup>, Shirley Viteri<sup>15</sup>, David Wessel<sup>4</sup>, Heather A. Wolfe<sup>1</sup>, Andrew R. Yates<sup>9</sup>, Athena F. Zuppa<sup>1</sup>, Robert M. Sutton<sup>1</sup> and Alexis A. Topjian<sup>1</sup>

## Abstract

**Introduction** Though early hypotension after pediatric in-hospital cardiac arrest (IHCA) is associated with inferior outcomes, ideal post-arrest blood pressure (BP) targets have not been established. We aimed to leverage prospectively collected BP data to explore the association of post-arrest BP thresholds with outcomes. We hypothesized that post-arrest systolic and diastolic BP thresholds would be higher than the currently recommended post-cardiopulmonary resuscitation BP targets and would be associated with higher rates of survival to hospital discharge.

**Methods** We performed a secondary analysis of prospectively collected BP data from the first 24 h following return of circulation from index IHCA events enrolled in the *ICU-RESUSCitation* trial (NCT02837497). The lowest documented systolic BP (SBP) and diastolic BP (DBP) were percentile-adjusted for age, height and sex. Receiver operator characteristic curves and cubic spline analyses controlling for illness category and presence of pre-arrest hypotension were generated exploring the association of lowest post-arrest SBP and DBP with survival to hospital discharge and survival to hospital discharge with favorable neurologic outcome (Pediatric Cerebral Performance Category of 1–3 or no change from baseline). Optimal cutoffs for post-arrest BP thresholds were based on analysis of receiver operator characteristic curves and spline curves. Logistic regression models accounting for illness category and pre-arrest hypotension examined the associations of these thresholds with outcomes.

**Results** Among 693 index events with 0–6 h post-arrest BP data, identified thresholds were: SBP > 10th percentile and DBP > 50th percentile for age, sex and height. Fifty-one percent ( $n = 352$ ) of subjects had lowest SBP above threshold and 50% ( $n = 346$ ) had lowest DBP above threshold. SBP and DBP above thresholds were each associated with survival to hospital discharge (SBP: aRR 1.21 [95% CI 1.10, 1.33]; DBP: aRR 1.23 [1.12, 1.34]) and survival to hospital discharge with favorable neurologic outcome (SBP: aRR 1.22 [1.10, 1.35]; DBP: aRR 1.27 [1.15, 1.40]) (all  $p < 0.001$ ).

\*Correspondence:

Monique M. Gardner  
gardnerm@chop.edu

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Conclusions** Following pediatric IHCA, subjects had higher rates of survival to hospital discharge and survival to hospital discharge with favorable neurologic outcome when BP targets above a threshold of SBP > 10th percentile for age and DBP > 50th percentile for age during the first 6 h post-arrest.

**Keywords** Pediatric, Infant, Neonatal, Cardiopulmonary resuscitation, Post-cardiac arrest, Blood pressure, Hypotension, Outcomes

## Introduction

More than 15,000 children per year in the USA have an in-hospital cardiac arrest (IHCA), with survival rates of 45–50% [1, 2]. Children who survive the cardiac arrest event are at high risk for post-cardiac arrest syndrome (PCAS), which includes brain injury, myocardial dysfunction, ischemia and reperfusion injury, and the ongoing underlying precipitating condition [3]. Hypotension, which is common during PCAS, is due to a combination of myocardial dysfunction and the ischemia–reperfusion syndrome [4] and can lead to secondary cerebral and myocardial injury resulting in worse outcomes.

Post-cardiac arrest systolic hypotension has previously been defined as systolic blood pressure (SBP) less than the 5th percentile for age and sex based on normative data of blood pressures in the healthy pediatric population [5]. Prior studies in both adults [6–10] and pediatrics [11–14] have focused on early post-arrest blood pressure measurements (i.e., within the first 6 h after return of spontaneous circulation [ROSC]) when organ reperfusion may be particularly critical and have demonstrated that early hypotension is associated with higher rates of post-arrest hospital mortality and worse neurologic outcomes. Consequently, pediatric guidelines recommend avoiding blood pressure less than the 5th percentile for age in the post-arrest period [3, 15], though this cutoff was selected without prior evidence, and no previous study has sought to identify best blood pressure thresholds in pediatrics. In contrast, adult studies have established that survival to hospital discharge is associated with minimum blood pressure thresholds much higher than the 5th percentile [16] and that myocardial injury can be mitigated with post-arrest mean arterial pressure (MAP) > 80 mmHg [17].

Using prospectively collected data from the *ICU-RESUSCitation* (*ICU-RESUS*) clinical trial [18, 19], we aimed to define post-arrest SBP and diastolic blood pressure (DBP) percentile thresholds based on the probability of survival to hospital discharge with favorable neurologic outcome and then evaluate the association of these new thresholds at 0–6 h and 6–24 h post-arrest with survival to hospital discharge and survival to hospital discharge with favorable neurologic outcome. In a subset of subjects with intra-arrest

hemodynamic data, we aimed to evaluate the association of intra-arrest BP targets with post-arrest BP thresholds.

## Methods

The *ICU-RESUS* trial (ClinicalTrials.gov Identifier: NCT02837497) was a parallel-stepped-wedge hybrid cluster-randomized interventional trial in 18 pediatric intensive care units and pediatric cardiac intensive care units (ICUs, clusters) from 10 clinical sites in the USA between October 1, 2016, and March 31, 2021. The methods [18] and primary results [19] have been previously published. Inclusion criteria were age between 37 weeks corrected gestation and ≤ 18 years and CPR of any duration in the ICU. Subjects were excluded if goals of care limited aggressive ICU therapies prior to arrest, they were brain dead, or they had an out-of-hospital cardiac arrest associated with the current hospitalization. The intervention included a two-part ICU resuscitation quality improvement bundle of point-of-care cardiopulmonary resuscitation (CPR) training on a manikin and structured cardiac arrest debriefs. Debriefs included review of physiologic goals for CPR and post-arrest care targets including hemodynamics. The primary outcome was survival to hospital discharge with favorable neurologic outcome (defined as Pediatric Cerebral Performance Category [PCPC] score of 1–3 or no change from baseline). The secondary outcome was survival to hospital discharge.

Data elements collected for the *ICU-RESUS* trial included subject demographics, arrest characteristics and post-cardiac arrest care data, and survival to hospital discharge and survival to hospital discharge with favorable neurologic outcome, per standard Utstein-style reporting [20]. Pediatric Risk of Mortality (PRISM) III score [21] was assessed 2–6 h prior to cardiac arrest. The vasoactive inotropic score (VIS) was calculated with the following equation: dopamine dose ( $\mu\text{g}/\text{kg}/\text{min}$ ) + dobutamine dose ( $\mu\text{g}/\text{kg}/\text{min}$ ) +  $100 \times$  epinephrine dose ( $\mu\text{g}/\text{kg}/\text{min}$ ) +  $10 \times$  milrinone dose ( $\mu\text{g}/\text{kg}/\text{min}$ ) +  $10\,000 \times$  vasopressin dose (unit/kg/min) +  $100 \times$  norepinephrine dose ( $\mu\text{g}/\text{kg}/\text{min}$ ) at 2 h prior to cardiac arrest, and 0–6 h and 6–24 h post-arrest [22, 23].

The *ICU-RESUS* parent study collected the highest and lowest invasive and non-invasive SBP and DBP were

collected for each subject at 0–6 h and 6–24 h post-arrest [19], as was done in previous pediatric studies [12–14]. We focused primarily on early (0–6 h) hypotension for comparison to prior adult and pediatric studies [6–10, 12–14]. When available, intra-arrest blood pressure was measured via arterial catheters and analyzed in 30-s epochs for the first 10 min of CPR. Providers were trained to target previously reported intra-arrest blood pressure targets: intra-arrest DBP  $\geq 25$  mmHg for subjects  $< 1$  year of age, and  $\geq 30$  mmHg for subjects  $\geq 1$  year [24, 25]. Post-arrest variables collected were highest and lowest temperature ( $^{\circ}\text{C}$ ), arterial PaO<sub>2</sub> (mmHg), PCO<sub>2</sub> (mmHg), pH, glucose, pulse oximetry (SpO<sub>2</sub>), and highest lactate in the 0–6 h and 6–24 h post-arrest intervals.

### Post-arrest blood pressure study

All patients from the *ICU-RESUS* trial were eligible for this study. We excluded patients without ROSC, who were treated with ECMO within 6 h of cardiac arrest, or who did not have blood pressure values available at 0–6 h post-arrest. The lowest SBP and DBP for each patient at 0–6 h and 6–24 h were percentile adjusted for age, sex, and height based on normative data of blood pressures in the healthy pediatric population [5]. All blood pressures were reviewed by trained staff to limit erroneous or artifactual values [19].

### Identification of blood pressure thresholds associated with outcome

Using this cohort, we determined the threshold probability of survival to hospital discharge with favorable neurological outcome versus adjusted percentiles of lowest SBP and DBP at 0–6 h post-arrest. Probability splines were generated with logistic regression, controlling for illness category (medical cardiac, surgical cardiac, medical non-cardiac, surgical non-cardiac) and pre-arrest hypotension (defined as SBP  $< 5$ th percentile for age, sex, and height; mean arterial blood pressure  $< 5$ th percentile for age; or presence of vasopressor/inotropic support after volume expansion, excluding low-dose dopamine). A natural cubic spline with internal knots at the 10th, 50th, and 90th percentiles was used to accurately capture the relationship between post-arrest blood pressures and survival to hospital discharge with favorable neurologic outcome. We determined post-arrest threshold percentile cutoff points of SBP at 0–6 h and DBP at 0–6 h with the use AUROC curves (Additional file 1: Figure S1). Cutoffs were approximated to the nearest clinically relevant integer value. We generated and inspected natural cubic spline with internal knots at the 10th, 50th, and 90th percentiles to graphically depict the relationship

between post-arrest blood pressures and survival to hospital discharge with favorable neurologic outcome. The spline curve provided further support for the choice of post-arrest threshold percentile cutoff points of SBP at 0–6 h and DBP at 0–6 h.

### Association of blood pressure thresholds associated with outcome

The primary exposure was whether the lowest documented SBP or DBP at 0–6 and 6–24 h post-arrest met the prospectively identified SBP and DBP percentile thresholds. The primary outcome was survival to hospital discharge with favorable neurologic outcome and the secondary outcome was survival to hospital discharge. We hypothesized that patients with SBP and DBP above identified thresholds would have higher rates of survival to hospital discharge with favorable neurologic outcome.

For description and comparison of patient and event characteristics, subjects were dichotomized into groups based on prospectively identified SBP and DBP thresholds. Frequencies and percentages were used to summarize categorical variables, and medians and interquartile ranges were used to summarize continuous variables. Associations of covariates with post-arrest blood pressure above threshold were assessed with Fisher's exact test for categorical variables and the Wilcoxon rank-sum test for ordinal variables. The relationship between vasoactive inotropic score at 6- and 24-h post-arrest and post-arrest SBP and DBP above threshold at 0–6 h post-arrest was assessed with the Wilcoxon rank-sum test. Vasoactive inotropic scores of zero were reported as 'none', 0–20, or  $> 20$ . Survival to hospital discharge and survival to hospital discharge with favorable neurologic outcome were summarized and assessed with Fisher's exact test for categorical variables and the Wilcoxon rank-sum test for ordinal variables. These analyses were repeated at the 6–24 h post-arrest.

The association between SBP and DBP thresholds at 0–6 h post-arrest with (1) survival to hospital discharge with favorable neurologic outcome and (2) survival to hospital discharge were assessed using Poisson regression with robust error estimates, controlling for illness category and pre-arrest hypotension. These covariates were selected a priori by the writing group, prior to any data analysis [26], and were consistent among the *ICU-RESUS* planned secondary analyses [27, 28]. Analyses were repeated for lowest SBP and DBP categories at 6–24 h. To test for the impact of the original trial's intervention, we added the variable of treatment category during the parent study (control group

versus transition group versus *ICU-RESUS* bundle) and repeated all analyses.

**Association of intra-arrest blood pressure and post-arrest blood pressure**

Intra-arrest invasively measured blood pressure data were prospectively collected and analyzed as a component of the *ICU-RESUS* trial [18, 19, 25, 29]. We evaluated the association between intra-arrest mean DBP targets and post-arrest study SBP and DBP thresholds at 0–6 h post-arrest using Poisson regression with robust error estimates, controlling for illness category and pre-arrest hypotension. We evaluated the interaction between intra-arrest and post-arrest blood pressure thresholds on outcome.

Covariates for modeling were specified a priori. *P* values were reported based on a 2-sided alternative and considered statistically significant when less than 0.05. Analyses were performed using SAS software v9.4 (Cary, NC, USA).

**Results**

The cohort included 1,129 index events; 773 subjects were evaluable (Fig. 1). After excluding for initiation of ECMO and those who died during the time period, there were 693 analyzable subjects at 0–6 h post-arrest and 693 subjects at 6–24 h post-arrest. Of those that had BP data analyzed for 0–6 h post-arrest, 513 (74.0%) survived


to hospital discharge, and 477 (68.8%) survived to hospital discharge with favorable neurologic outcome. Of the subjects with BP data for 6–24 h, 504 (79.3%) survived to hospital discharge, and 469 (73.7%) survived to hospital discharge with favorable neurologic outcome.

**Identification of systolic and diastolic blood pressure thresholds**

Optimal cutoffs on the AUROC curves were SBP at the 7th percentile adjusted for age, sex, and height and DBP at the 49th percentile adjusted for age, sex, and height. These were supported by splines. For clinical relevance, post-arrest lowest SBP above the 10th percentile at both 0–6 h was the threshold where the probability of survival to hospital discharge with favorable neurologic outcome plateaued. Lowest DBP above the 50th percentile at 0–6 h post-arrest was the threshold where the probability of survival to hospital discharge with favorable neurologic outcome plateaued (Fig. 2). These 0–6 h thresholds for lowest SBP of 10th percentile and lowest DBP of 50th percentile were used for subsequent analyses.


**Systolic blood pressure 0–6 h post-arrest**

There were 352 subjects (50.8%) with the lowest SBP at 0–6 h post-arrest above the 10th percentile threshold (Table 1). These subjects received shorter durations of CPR; fewer doses of epinephrine during arrest; less frequently received sodium bicarbonate and calcium during



**Fig. 1** Consort Diagram





**Fig. 2** Splines of the association of age-, sex-, and height-adjusted systolic (Panel **A**) and diastolic (Panel **B**) blood pressure percentiles at 0–6 h post-arrest by survival to hospital discharge with favorable neurologic outcome controlling for pre-arrest hypotension and illness category. Light blue boundaries represent 95% confidence intervals. Dashed line represents identified blood pressure threshold

arrest; and had lower VIS scores in the first 6 h post-arrest (Table 2). Fewer subjects with lowest SBP above threshold had hypotension as an immediate cause of arrest, but more had respiratory insufficiency.

In univariate analysis, subjects with lowest SBP above threshold had higher rates of survival to hospital discharge with favorable neurologic outcome (76.4% vs 61.0%,  $p < 0.001$ ) and survival to hospital discharge (82.1% vs. 65.7%,  $p < 0.001$ ) (Table 2). In a multivariable model adjusting for illness category and pre-arrest hypotension, subjects with lowest SBP above the 10th percentile threshold were more likely to survive to hospital discharge with favorable neurologic outcome (RR 1.22 [95% CI 1.10, 1.35],  $p < 0.001$ ) and survive to hospital discharge (RR 1.21 [95% CI 1.10, 1.33],  $p < 0.001$ ) (Table 3). Controlling for *ICU-RESUS* treatment category in the initial interventional trial in addition to illness category and pre-arrest hypotension revealed similar results (Additional file 5: Table S4).

#### Diastolic blood pressure 0–6 h post-arrest

Half of the cohort (346 subjects, 49.9%) had lowest DBP above the 50th percentile threshold (Table 1). Subjects with lowest DBP above threshold more frequently had pre-existing congenital heart disease and medical-cardiac or surgical-cardiac illness categories (Table 1) and had lower vasoactive inotropic score in the first 6 h after arrest compared to those with lowest DBP below threshold (Table 2). Fewer subjects with lowest DBP above threshold had hypotension as the immediate cause of arrest (Table 2). Arrest duration was shorter in those with lowest DBP above threshold (3 vs. 4 min,  $P = 0.012$ ) and these subjects less frequently received intra-arrest

epinephrine, calcium, sodium bicarbonate, and fluid resuscitation.

In univariate analysis subjects with lowest DBP above threshold at 0–6 h post-arrest had higher rates of survival to hospital discharge with favorable neurologic outcome (77.2% vs. 60.5%,  $p < 0.001$ ) and survival to hospital discharge (81.8% vs. 63.3%,  $p < 0.001$ ; Table 2). In a multivariable model controlling for illness category and pre-arrest hypotension, lowest DBP above threshold was associated with higher rates of survival to hospital discharge with favorable neurologic outcome (aRR 1.27 [95% CI 1.15, 1.40],  $p < 0.001$ ) and survival to hospital discharge (aRR 1.23 [95% CI 1.12, 1.34],  $p < 0.001$ ; Table 3). Adding *ICU-RESUS* treatment category in the original study to the multivariate analysis yielded similar results (Additional file 6: Table S5).

#### Systolic and diastolic blood pressure post-arrest 6–24 h

We compared groups with lowest SBP and lowest DBP above threshold to those below threshold at 6–24 h post-arrest and found similar patterns to the 0–6 h post-arrest (Additional file 2: Table S1). In contrast to findings at the 0–6-h post-arrest, subjects with lowest SBP and DBP above threshold at 6–24 h did not differ in intra-arrest administration of sodium bicarbonate compared to those with lowest SBP and DBP below threshold (Additional file 3: Table S2). Similar to 0–6 h post-arrest, lowest SBP and DBP above threshold at 6–24 h post-arrest were associated with higher rates of survival to hospital discharge with favorable neurologic outcome, and survival to hospital discharge (Additional file 3: Table S2), including after controlling for illness category and pre-arrest hypotension (Additional file 4: Table S3).

**Table 1** Demographics and pre-event characteristics by systolic and diastolic blood pressure thresholds at 0–6 h post-arrest

	Post-arrest systolic BP (0–6 h) <sup>a</sup>		P value	Post-arrest diastolic BP (0–6 h) <sup>a</sup>		P value
	> 10th percentile (N = 352)	≤ 10th percentile (N = 341)		> 50th percentile (N = 346)	≤ 50th percentile (N = 347)	
Demographics						
Age			<.001 <sup>c</sup>			<.001 <sup>c</sup>
≤ 1 month	27 (7.7%)	74 (21.7%)		63 (18.2%)	38 (11.0%)	
1 month–< 1 year	174 (49.4%)	133 (39.0%)		193 (55.8%)	114 (32.9%)	
1 year–< 8 years	108 (30.7%)	70 (20.5%)		67 (19.4%)	111 (32.0%)	
8 years–< 19 years	43 (12.2%)	64 (18.8%)		23 (6.6%)	84 (24.2%)	
Male	199 (56.5%)	175 (51.3%)	0.171 <sup>c</sup>	191 (55.2%)	183 (52.7%)	0.542 <sup>c</sup>
Race			0.017 <sup>c</sup>			0.812 <sup>c</sup>
White	156 (44.3%)	172 (50.4%)		158 (45.7%)	170 (49.0%)	
Black or African American	106 (30.1%)	68 (19.9%)		87 (25.1%)	87 (25.1%)	
Other	25 (7.1%)	22 (6.5%)		21 (6.1%)	26 (7.5%)	
Unknown or not reported	65 (18.5%)	79 (23.2%)		80 (23.1%)	64 (18.4%)	
Hispanic or Latino	51 (14.5%)	55 (16.1%)	0.396 <sup>c</sup>	59 (17.1%)	47 (13.5%)	0.287 <sup>c</sup>
Pre-existing medical conditions						
Respiratory insufficiency	319 (90.6%)	289 (84.8%)	0.020 <sup>c</sup>	310 (89.6%)	298 (85.9%)	0.164 <sup>c</sup>
Hypotension	154 (43.8%)	221 (64.8%)	<.001 <sup>c</sup>	182 (52.6%)	193 (55.6%)	0.446 <sup>c</sup>
Congestive heart failure	34 (9.7%)	28 (8.2%)	0.510 <sup>c</sup>	31 (9.0%)	31 (8.9%)	1.000 <sup>c</sup>
Pneumonia	58 (16.5%)	35 (10.3%)	0.019 <sup>c</sup>	45 (13.0%)	48 (13.8%)	0.824 <sup>c</sup>
Sepsis	44 (12.5%)	53 (15.5%)	0.274 <sup>c</sup>	33 (9.5%)	64 (18.4%)	<.001 <sup>c</sup>
Renal insufficiency	33 (9.4%)	46 (13.5%)	0.095 <sup>c</sup>	21 (6.1%)	58 (16.7%)	<.001 <sup>c</sup>
Congenital heart disease	175 (49.7%)	210 (61.6%)	0.002 <sup>c</sup>	215 (62.1%)	170 (49.0%)	<.001 <sup>c</sup>
Pre-event characteristics						
Illness category			<.001 <sup>c</sup>			0.003 <sup>c</sup>
Medical cardiac	76 (21.6%)	75 (22.0%)		89 (25.7%)	62 (17.9%)	
Medical non-cardiac	159 (45.2%)	111 (32.6%)		123 (35.5%)	147 (42.4%)	
Surgical cardiac	87 (24.7%)	129 (37.8%)		115 (33.2%)	101 (29.1%)	
Surgical non-cardiac or trauma	30 (8.5%)	26 (7.6%)		19 (5.5%)	37 (10.7%)	
Baseline Pediatric Cerebral Performance Category <sup>b</sup>			0.828 <sup>d</sup>			<.001 <sup>d</sup>
1—Normal	208 (59.1%)	206 (60.4%)		231 (66.8%)	183 (52.7%)	
2—Mild disability	59 (16.8%)	53 (15.5%)		55 (15.9%)	57 (16.4%)	
3—Moderate disability	46 (13.1%)	41 (12.0%)		38 (11.0%)	49 (14.1%)	
4—Severe disability	35 (9.9%)	38 (11.1%)		19 (5.5%)	54 (15.6%)	
5—Coma/vegetative state	4 (1.1%)	3 (0.9%)		3 (0.9%)	4 (1.2%)	

<sup>a</sup> Post-arrest systolic or diastolic thresholds defined as the minimum recorded systolic or diastolic blood pressure in the time period less than or equal to the 10th or 50th percentile for age, sex, and height, respectively

<sup>b</sup> Baseline pediatric cerebral performance category and functional status scale were evaluated prior to the event leading to hospitalization

<sup>c</sup> Fishers exact test

<sup>d</sup> Wilcoxon rank-sum test

### Intra-arrest blood pressures and CPR quality measures and early post-arrest hemodynamics

Intra-arrest hemodynamics were available for analysis in 227 patients (Table 4). Compared to infants (≤ 1 year of age) with post-arrest SBP below threshold at 0–6 h, infants with post-arrest SBP above threshold had higher mean intra-arrest SBP. Similarly, infants with post-arrest

DBP above threshold at 0–6 h had higher mean intra-arrest SBP and DBP, compared to infants below respective thresholds. A higher percentage of all patients (i.e., infants + older children) with post-arrest DBP above threshold at 0–6 h met intra-arrest age-based SBP targets than those with post-arrest DBP below threshold. An association between post-arrest SBP above threshold and



**Table 2** Event characteristics and outcomes by systolic and diastolic blood pressure thresholds at 0–6 h post-arrest

	Post-arrest systolic BP (0–6 h) <sup>a</sup>		P value	Post-arrest diastolic BP (0–6 h) <sup>a</sup>		P value
	> 10th percentile (N = 352)	≤ 10th percentile (N = 341)		> 50th percentile (N = 346)	≤ 50th percentile (N = 347)	
Interventions in place prior to event						
Vascular access	308 (87.5%)	320 (93.8%)	0.006 <sup>b</sup>	301 (87.0%)	327 (94.2%)	0.001 <sup>b</sup>
Arterial catheter	130 (36.9%)	174 (51.0%)	< .001 <sup>b</sup>	152 (43.9%)	152 (43.8%)	1.000 <sup>b</sup>
Central venous catheter	208 (59.1%)	232 (68.0%)	0.018 <sup>b</sup>	220 (63.6%)	220 (63.4%)	1.000 <sup>b</sup>
Vasoactive infusion	111 (31.5%)	167 (49.0%)	< .001 <sup>b</sup>	138 (39.9%)	140 (40.3%)	0.938 <sup>b</sup>
Invasive mechanical ventilation	224 (63.6%)	241 (70.7%)	0.052 <sup>b</sup>	224 (64.7%)	241 (69.5%)	0.196 <sup>b</sup>
Non-invasive ventilation	78 (22.2%)	63 (18.5%)	0.258 <sup>b</sup>	75 (21.7%)	66 (19.0%)	0.397 <sup>b</sup>
End-tidal CO <sub>2</sub> monitoring	186 (52.8%)	226 (66.3%)	< .001 <sup>b</sup>	193 (55.8%)	219 (63.1%)	0.053 <sup>b</sup>
Immediate causes of event						
Arrhythmia	55 (15.6%)	47 (13.8%)	0.521 <sup>b</sup>	49 (14.2%)	53 (15.3%)	0.748 <sup>b</sup>
Cyanosis without respiratory decompensation	19 (5.4%)	9 (2.6%)	0.082 <sup>b</sup>	18 (5.2%)	10 (2.9%)	0.128 <sup>b</sup>
Hypotension	118 (33.5%)	196 (57.5%)	< .001 <sup>b</sup>	139 (40.2%)	175 (50.4%)	0.008 <sup>b</sup>
Respiratory decompensation	241 (68.5%)	192 (56.3%)	< .001 <sup>b</sup>	225 (65.0%)	208 (59.9%)	0.182 <sup>b</sup>
Vasoactive inotropic score (2 h prior to CPR)	0.0 [0.0, 0.8]	0.0 [0.0, 5.0]	< .001 <sup>c</sup>	0.0 [0.0, 5.0]	0.0 [0.0, 3.0]	0.113 <sup>b</sup>
Duration of CPR (minutes) <sup>d</sup>	3.0 [1.0, 6.0]	4.0 [2.0, 8.0]	0.005 <sup>c</sup>	3.0 [1.0, 6.0]	4.0 [2.0, 8.0]	0.012 <sup>c</sup>
Duration of CPR categories (minutes) <sup>d</sup>			0.129 <sup>b</sup>			0.011 <sup>b</sup>
< 6	247 (70.2%)	212 (62.2%)		244 (70.5%)	215 (62.0%)	
6–15	72 (20.5%)	83 (24.3%)		74 (21.4%)	81 (23.3%)	
16–35	26 (7.4%)	34 (10.0%)		24 (6.9%)	36 (10.4%)	
> 35	7 (2.0%)	12 (3.5%)		4 (1.2%)	15 (4.3%)	
CPR time <sup>d</sup>			0.904 <sup>b</sup>			0.899 <sup>b</sup>
Weekday	185 (52.6%)	184 (54.0%)		183 (52.9%)	186 (53.6%)	
Weeknight	57 (16.2%)	56 (16.4%)		55 (15.9%)	58 (16.7%)	
Weekend	110 (31.3%)	101 (29.6%)		108 (31.2%)	103 (29.7%)	
First documented rhythm			0.514 <sup>b</sup>			0.219 <sup>b</sup>
Pulseless electrical activity / asystole	132 (37.5%)	134 (39.3%)		124 (35.8%)	142 (40.9%)	
Ventricular fibrillation / tachycardia	17 (4.8%)	22 (6.5%)		17 (4.9%)	22 (6.3%)	
Bradycardia with poor perfusion	203 (57.7%)	185 (54.3%)		205 (59.2%)	183 (52.7%)	
Pharmacologic interventions during event						
Epinephrine	214 (60.8%)	267 (78.3%)	< .001 <sup>b</sup>	226 (65.3%)	255 (73.5%)	0.021 <sup>b</sup>
Minutes to first epinephrine bolus	1.0 [0.0, 2.0]	1.0 [0.0, 2.0]	0.817 <sup>c</sup>	1.0 [0.0, 2.0]	1.0 [0.0, 2.0]	0.099 <sup>c</sup>
Number of epinephrine boluses	2.0 [1.0, 3.0]	2.0 [1.0, 3.0]	0.713 <sup>c</sup>	1.5 [1.0, 3.0]	2.0 [1.0, 3.0]	0.517 <sup>c</sup>
Atropine	46 (13.1%)	33 (9.7%)	0.188 <sup>b</sup>	46 (13.3%)	33 (9.5%)	0.122 <sup>b</sup>
Calcium	60 (17.0%)	117 (34.3%)	< .001 <sup>b</sup>	76 (22.0%)	101 (29.1%)	0.036 <sup>b</sup>
Sodium bicarbonate	81 (23.0%)	135 (39.6%)	< .001 <sup>b</sup>	95 (27.5%)	121 (34.9%)	0.040 <sup>b</sup>
Vasopressin	5 (1.4%)	10 (2.9%)	0.199 <sup>b</sup>	5 (1.4%)	10 (2.9%)	0.296 <sup>b</sup>
Fluid bolus	56 (15.9%)	72 (21.1%)	0.079 <sup>b</sup>	53 (15.3%)	75 (21.6%)	0.040 <sup>b</sup>
Vasoactive-inotropic score						
Vasoactive inotropic score 6 h post-arrest			< .001 <sup>b</sup>			0.006 <sup>b</sup>
None	207 (58.8%)	133 (39.0%)		178 (51.4%)	162 (46.7%)	
1–20	126 (35.8%)	158 (46.3%)		146 (42.2%)	138 (39.8%)	
> 20	19 (5.4%)	50 (14.7%)		22 (6.4%)	47 (13.5%)	
Vasoactive inotropic score 24 h post-arrest			< .001 <sup>b</sup>			0.188 <sup>b</sup>
None	214 (60.8%)	153 (44.9%)		181 (52.3%)	186 (53.6%)	
1–20	123 (34.9%)	154 (45.2%)		146 (42.2%)	131 (37.8%)	
> 20	15 (4.3%)	34 (10.0%)		19 (5.5%)	30 (8.6%)	

**Table 2** (continued)

	Post-arrest systolic BP (0–6 h) <sup>a</sup>		P value	Post-arrest diastolic BP (0–6 h) <sup>a</sup>		P value
	> 10th percentile (N = 352)	≤ 10th percentile (N = 341)		> 50th percentile (N = 346)	≤ 50th percentile (N = 347)	
Outcomes						
Survival to hospital discharge	289 (82.1%)	224 (65.7%)	<.001 <sup>b</sup>	283 (81.8%)	230 (66.3%)	<.001 <sup>b</sup>
Survival to hospital discharge with favorable neurologic outcome	269 (76.4%)	208 (61.0%)	<.001 <sup>b</sup>	267 (77.2%)	210 (60.5%)	<.001 <sup>b</sup>

<sup>a</sup> Post-arrest systolic or diastolic thresholds defined as the minimum recorded systolic or diastolic blood pressure in the time period less than or equal to the 10th or 50th percentile for age, sex, and height, respectively

<sup>b</sup> Fishers Exact test

<sup>c</sup> Wilcoxon rank-sum test

<sup>d</sup> CPR = Cardiopulmonary resuscitation

**Table 3** Multivariable associations between systolic and diastolic pressure 0–6 h post arrest and survival

	Overall (N = 693)	Survival to hospital discharge with favorable neurologic outcome <sup>a</sup>		Survival to hospital discharge with favorable neurologic outcome <sup>a</sup>	
		Relative risk (95% CI)	P value	Relative risk (95% CI)	P value
Post-arrest systolic threshold (0–6 h) <sup>c</sup>			<.001		<.001
Above (> 10th percentile)	352 (50.8%)	1.22 (1.10, 1.35)		1.21 (1.10, 1.33)	
Below—inclusive (≤ 10th percentile)	341 (49.2%)	Reference		Reference	
Post-arrest diastolic threshold (0–6 h) <sup>c</sup>			<.001		<.001
Above (> 50th percentile)	346 (49.9%)	1.27 (1.15, 1.40)		1.23 (1.12, 1.34)	
Below—inclusive (≤ 50th percentile)	347 (50.1%)	Reference		Reference	
Controlling also for intra-arrest diastolic pressure (met target: yes, no) <sup>b,c</sup> (N = 227)					
Post-arrest systolic threshold (0–6 h) <sup>c</sup>			0.500		0.769
Above (> 10th percentile)	100 (44.1%)	1.06 (0.90, 1.25)		1.02 (0.88, 1.19)	
Below—inclusive (≤ 10th percentile)	127 (55.9%)	Reference		Reference	
Post-arrest diastolic threshold (0–6 h) <sup>c</sup>			0.586		0.636
Above (> 50th percentile)	121 (53.3%)	1.05 (0.89, 1.24)		1.04 (0.89, 1.20)	
Below—inclusive (≤ 50th percentile)	106 (46.7%)	Reference		Reference	
Interaction between intra- and post-arrest diastolic pressure (N = 227)			0.736		0.332
Above intra-arrest target and above post-arrest threshold	112 (16.2%)	1.21 (0.79, 1.87)		1.27 (0.81, 1.98)	
Above intra-arrest target and below post-arrest threshold	94 (13.6%)	1.19 (0.77, 1.84)		1.27 (0.80, 2.00)	
Below intra-arrest target and above post-arrest threshold	9 (1.3%)	1.33 (0.80, 2.19)		1.51 (0.92, 2.46)	
Below intra-arrest target and below post-arrest threshold	12 (1.7%)	Reference		Reference	

<sup>a</sup> Favorable neurologic outcome was defined as no more than moderate disability or no worsening from baseline Pediatric Cerebral Performance Category (PCPC). Baseline PCPC represents subject status prior to the event leading to hospitalization

<sup>b</sup> All models control for illness category and pre-arrest hypotension. Two, noted above, also control for whether the intra-arrest diastolic blood pressure target was met

<sup>c</sup> Target intra-arrest diastolic pressure was ≥ 25 mmHg for subjects < 1 year old or ≥ 30 for subjects ≥ 1 year. Post-arrest systolic or diastolic thresholds defined as the minimum recorded systolic or diastolic blood pressure in the time period ≤ the 10th or 50th percentile for age, sex, and height, respectively

intra-arrest SBP was not observed. There were no differences in measures of CPR quality including compression rate and chest compression fraction over the first 10 min of CPR between patients above and below post-arrest SBP or DBP thresholds. There was no interaction between intra-arrest and post-arrest DBP at 0–6 h on

outcomes (Table 3). In this smaller cohort, when controlling for intra-arrest mean DBP threshold in addition to illness category and pre-arrest hypotension, post-arrest lowest DBP and SBP were not associated with survival to hospital discharge with favorable neurologic outcome or survival to hospital discharge (Table 3). Repeating this

**Table 4** Intra-arrest hemodynamics and chest compression mechanics by systolic and diastolic blood pressure thresholds 0–6 h post

	Post-arrest systolic BP (0–6 h) <sup>a</sup>		P value	Post-arrest diastolic BP (0–6 h) <sup>a</sup>		P value
	> 10th percentile (N = 100)	≤ 10th percentile (N = 127)		> 50th percentile (N = 121)	≤ 50th percentile (N = 106)	
Intra-arrest event characteristics						
Average systolic blood pressure (mmHg)						
Age ≤ 1 year	82.0 [68.9, 103.3]	72.2 [58.4, 87.9]	0.006 <sup>c</sup>	81.7 [64.8, 94.8]	67.2 [54.6, 82.3]	0.004 <sup>c</sup>
Age > 1 year	101.6 [85.1, 124.6]	90.3 [69.3, 116.2]	0.205 <sup>c</sup>	104.5 [95.1, 127.1]	90.3 [72.6, 115.4]	0.070 <sup>c</sup>
Average systolic blood pressure ≥ 60 mmHg for age < 1 year or ≥ 80 mmHg for age ≥ 1 year	78 (78.0%)	87 (68.5%)	0.283 <sup>b</sup>	96 (79.3%)	69 (65.1%)	0.009 <sup>b</sup>
Average diastolic blood pressure (mmHg)						
Age ≤ 1 year	41.1 [32.3, 48.6]	36.3 [30.1, 47.0]	0.176 <sup>c</sup>	41.5 [32.0, 49.9]	34.6 [29.5, 46.0]	0.026 <sup>c</sup>
Age > 1 year	43.4 [37.5, 60.6]	44.5 [34.5, 54.7]	0.628 <sup>c</sup>	53.8 [40.4, 63.7]	42.9 [34.5, 53.4]	0.052 <sup>c</sup>
Average diastolic blood pressure ≥ 25 mmHg for age < 1 year or ≥ 30 mmHg for age ≥ 1 year	92 (92.0%)	114 (89.8%)	0.648 <sup>b</sup>	112 (92.6%)	94 (88.7%)	0.363 <sup>b</sup>
Average chest compression rate over the first 10 min	121.6 [112.7, 132.7]	124.6 [113.4, 132.8]	0.719 <sup>c</sup>	121.97 [112.14, 132.00]	124.74 [116.30, 133.01]	0.256 <sup>c</sup>
Average chest compression fraction ≥ 0.80	96 (96.0%)	117 (92.1%)	0.276 <sup>b</sup>	116 (95.9%)	97 (91.5%)	0.268 <sup>b</sup>
Average chest compression fraction over the first 10 min	1.0 [0.9, 1.0]	1.0 [0.9, 1.0]	0.078 <sup>c</sup>	0.98 [0.95, 1.00]	0.98 [0.92, 1.00]	0.336 <sup>c</sup>

<sup>a</sup> Post-arrest systolic or diastolic blood pressure thresholds defined as the minimum recorded systolic or diastolic blood pressure in the time period ≤ the 10th or 50th percentile for age, sex, and height, respectively

<sup>b</sup> Fishers Exact test

<sup>c</sup> Wilcoxon rank-sum test

analysis on lowest SBP and DBP 6–24 h post-arrest, no significant associations were found (Additional file 4: Table S3).

## Discussion

In this study, we used a large prospectively collected dataset to define thresholds of lowest SBP and DBP at 0–6 h post-arrest that are associated with survival to hospital discharge with favorable neurologic outcome. After controlling for illness category and pre-arrest hypotension, absence of SBP below the 10th percentile and DBP below the 50th percentile during the first 6 h after ROSC were associated with higher rates of survival to hospital discharge with favorable neurologic outcome and survival to hospital discharge. This is the first study to prospectively identify post-arrest BP thresholds associated with outcomes and therefore establishes potential new targets for pediatric post-arrest care.

Hypotension is common after pediatric cardiac arrest and is associated with poor post-arrest outcomes [4, 11, 13, 14]. Prior work has focused on lower thresholds of hypotension, typically the 5th percentile for SBP for age, sex and height, which was selected a priori from normative blood pressure data in the general population [5]. In this study, we performed a robust analysis using

a large prospectively collected cardiac arrest dataset to define lowest SBP and DBP thresholds during the first 6 h post-arrest based on probability of survival to hospital discharge with favorable neurologic outcome. These newly derived thresholds for lowest BP during the first 6 h post-arrest, the 10th percentile for SBP and 50th percentile for DBP, are higher than lowest BP thresholds in previous studies and the targets recommended in current guidelines [3]. Our findings suggest that we should target higher blood pressure than previously recommended to potentially impact outcomes. Furthermore, it is not clear if titrating to specific blood pressures impacts outcomes after pediatric or adult cardiac arrest. Although these findings are limited by available data which only had highest and lowest SBP and DBP, these newly derived thresholds could facilitate future evaluations for interventional studies.

Post-cardiac arrest blood pressure and the impact on outcome is complicated, as hypotension can be both a manifestation of the post-arrest myocardial dysfunction [30, 31] and systemic inflammation [32], which are highly inter-related and both modifiable. Myocardial dysfunction can be treated with vasoactive-inotropic infusions. Post-arrest systemic inflammation can be treated by steroids and/or induced hypothermia, and the associated

vasodilatory shock can be treated with vasoconstrictor medications; importantly, we did not have time-synchronized post-arrest vasoactive and inotropic infusion rates available for further analyses of these relationships. Although the optimal medications and dosing for post-arrest myocardial dysfunction and post-arrest systemic inflammation are unknown, this study and others suggest that proactively maintaining adequate blood pressure is a reasonable therapeutic goal. Particularly, the importance of the association of early (0–6 h after arrest) hypotension and outcomes requires close clinical coordination to start monitoring and responding to blood pressure as quickly as possible after arrest to meet targets.

Several patient factors evaluated in this study were strongly associated with failure to exceed study BP thresholds. Not surprisingly, pre-arrest hypotension and vasoactive or inotropic support were associated with lowest SBP post-arrest below threshold. Pre-arrest renal insufficiency and congenital heart disease were associated with lowest SBP and lowest DBP post-arrest below threshold. While more granular analysis of these subpopulations was outside the scope of this study, these patients may require particular focus to understand their physiology and resuscitation, both intra- and post-arrest, to achieve optimal post-arrest outcomes. Proactive monitoring and targeted care for these high-risk groups may help decrease rates of hypotension and impact outcomes.

We found that subjects who had post-arrest DBP above threshold at 0–6 h were more likely to have met a priori intra-arrest SBP targets during the first 10 min of CPR. This association may be due to use of epinephrine or other vasoactive medications during resuscitation; though interestingly, this association did not persist for SBP thresholds and intra-arrest DBP or other CPR quality measures such as rate or compression fraction. Furthermore, the interactions between intra-arrest and post-arrest thresholds were not associated with outcomes, consistent with previous studies [33]. Previous studies show that intra-arrest DBP above target was associated with higher rates of survival to hospital discharge and with favorable neurologic outcome [24, 25]. While many would surmise that worse CPR quality, manifested as not achieving desired intra-arrest blood pressure targets, would be associated with worse post-arrest hypotension, we did not find this in our study. These findings highlight the complex and not well understood interactions between intra-arrest and post-arrest blood pressure. Notably, when we evaluated the association of post-arrest lowest SBP and DBP thresholds with outcomes and controlled for intra-arrest mean DBP, the association was no longer significant, suggesting that intra-arrest DBP confounds the relationship between the two. Further investigations of these interactions are needed.

This work has several limitations. The limitations of the primary study have been previously discussed [19]. Our analyses focused on the lowest DBP and SBP for each subject during each of the time intervals (0–6 h, and 6–24 h), consistent with most post-arrest investigations to date [12–14]. This may not reflect the total burden of hypotension during the time intervals [4]. Future work can focus on burden, as well as whether SBP or DBP thresholds are the main driver of outcomes. Mean arterial blood pressure (MAP) was not collected because normative data were not available at the time of the parent study design [18].

Additionally, post-arrest care was not standardized (though post-arrest goals were reviewed as part of the intervention), so sites monitored and treated post-arrest blood pressure at their discretion; we did not examine post-arrest myocardial dysfunction. As mentioned above, post-arrest time-synchronized VIS was not available for analysis. Variability of care among sites could impact both treatment of post-arrest hypotension and outcomes.

## Conclusions

We identified new thresholds of post-arrest blood pressure associated with survival to hospital discharge with favorable neurologic outcome and survival to hospital discharge: lowest SBP > 10th percentile for age and lowest DBP > 50th percentile for age. When controlling for illness category and pre-arrest hypotension, lowest DBP and SBP above these thresholds were associated with better outcomes. These findings continue to illustrate the importance of post-arrest blood pressure monitoring and management in efforts to improve outcomes from pediatric cardiac arrest.

## Abbreviations

CPR	Cardiopulmonary resuscitation
BP	Blood pressure
DBP	Diastolic blood pressure
ICU	Intensive care unit
IHCA	In-hospital cardiac arrest
PCPC	Pediatric Cerebral Performance Category
PCAS	Post-cardiac arrest syndrome
SBP	Systolic blood pressure
VIS	Vasoactive-inotropic score

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13054-023-04662-9>.

**Additional file 1. Figure S1:** (A) Association of outcome of survival to hospital discharge with favorable neurologic outcome with systolic blood pressure 0–6 post-arrest. The optimal cut point is at the 7th percentile adjusted for age, sex, and height (sensitivity 66%, specificity 57%). Area under the curve is 0.64. (B) Association of outcome of survival to hospital discharge with favorable neurologic outcome with diastolic blood pressure 0–6 post-arrest. The optimal cut point is at the 49th percentile

adjusted for age, sex, and height (sensitivity 60%, specificity 61%). Area under the curve is 0.64.

**Additional file 2. Supplemental Table 1.** Demographics and pre-event characteristics by systolic and diastolic blood pressure thresholds 6–24 h post-arrest.

**Additional file 3. Supplemental Table 2.** Event characteristics by diastolic and systolic blood pressure threshold 6–24 h post arrest.

**Additional file 4. Supplemental Table 3.** Multivariable associations between diastolic pressure 6–24 h post arrest and survival.

**Additional file 5. Supplemental Table 4.** Multivariable associations including treatment category between systolic and diastolic pressure 0–6 h post arrest and survival.

**Additional file 6. Supplemental Table 5.** Multivariable associations including treatment category between diastolic pressure 6–24 h post arrest and survival.

### Acknowledgements

Members of the CPCCRN/ICU RESUScitation Project Investigators are the authors listed on the first page and the following individuals: Robert Bishop, MD (Department of Pediatrics, University of Colorado School of Medicine and Children's Hospital Colorado, Aurora, CO), Matthew Bochkoris, MD (Department of Critical Care Medicine, UPMC Children's Hospital of Pittsburgh, University of Pittsburgh, Pittsburgh, PA), Richard Fernandez, MD (Department of Pediatrics, Nationwide Children's Hospital, The Ohio State University, Columbus, OH), Deborah Franzon, MD (Department of Pediatrics, Benioff Children's Hospital, University of California, San Francisco, San Francisco, CA), William Lardis (Department of Anesthesiology and Critical Care Medicine, The Children's Hospital of Philadelphia, University of Pennsylvania, Philadelphia, PA), Ashley Siems, MD (Department of Pediatrics, Children's National Hospital, George Washington University School of Medicine, Washington, DC), Sarah Tabbutt, MD, PhD (Department of Pediatrics, Benioff Children's Hospital, University of California, San Francisco, San Francisco, CA), Bradley Tilford, MD (Department of Pediatrics, Children's Hospital of Michigan, Central Michigan University, Detroit, MI), Shirley Viteri, MD (Department of Pediatrics, Nemours Cardiac Center, Nemours/Alfred I. duPont Hospital for Children, Wilmington, DE), and Anil Sapru (Department of Pediatrics, Mattel Children's Hospital, University of California Los Angeles, Los Angeles, CA).

### Author contributions

DAH, AAT, and RMS were primarily involved in submission of this a priori secondary analysis of ICU Resuscitation data. RWR and MLH performed statistical analysis and data presentation. MMG, RWR, RAB, MLH, RWM, RMS, and AAT were responsible for review of statistical analysis, interpretation of results, and were major contributors in writing the manuscript, with MMG serving as primary and corresponding author. All authors read and approved the final manuscript.

### Funding

The ICU RESUScitation Project was supported, in part, by the following grants from the National Institutes of Health National Heart, Lung, and Blood Institute and the Eunice Kennedy Shriver National Institute of Child Health and Human Development: R01HL131544, U01HD049934, UG1HD049981, UG1HD049983, UG1HD050096, UG1HD063108, UG1HD083166, UG1HD083170, UG1HD083171, and K23HL148541.

### Availability of data and materials

The dataset from the ICU Resuscitation trial is available for public use via the National Institutes of Health (NIH) Biologic Specimen and Data Repository Information Coordinating Center (BioLINCC).

### Declarations

#### Ethics approval and consent to participate

The institutional review board (IRB) at the University of Utah served as the single IRB and approved the study (ID: 00093320) with waiver of parental permission.

### Competing interests

Drs. Reeder's, Carcillo's, Carpenter's, Dean's, Fink's, Frazier's, Friess', Hall's, Manga's, Morgan's, Mourani's, Nadkarni's, Naim's, Palmer's, Pollack's, Wessel's, Wolfe's, Yates', Zuppa's, Sutton's, and Meert's institutions received funding from the National Institutes of Health (NIH). Drs. Reeder, Berg, Carcillo, Carpenter, Dean, Fink, Frazier, Friess, Hall, Horvat, Maa, McQuillen, Meert, Morgan, Mourani, Nadkarni, Naim, Pollack, Wessel, Wolfe, Yates, and Zuppa received support for article research from the NIH. Dr. Berg's institution received funding from the National Institute of Child Health and Human Development (NICHD) Collaborative Pediatric Critical Care Research Network grant and the National Heart, Lung, and Blood Institute (NHLBI) ICU-RESUS trial grant. Dr. Fink's institution received funding from the Neurocritical Care Society; she received funding from the American Board of Pediatrics and the Child Neurology Society. Dr. Friess received funding from an expert witness testimony. Dr. Hall received funding from Abbvie, La Jolla Pharmaceuticals, and Kiadis. Drs. Horvat, Maa, and McQuillen's institutions received funding from the NICHD. Dr. Horvat's institution received funding from the National Institute of Neurological Disorders and Stroke. Dr. Maa's institution received funding from the NHLBI. Dr. Mourani disclosed the off-label product use of sodium bicarbonate. Dr. Pollack disclosed work for hire. Dr. Wolfe received funding from The Debriefing Academy and Zoll. The remaining authors have disclosed that they do not have any potential conflicts of interest.

### Author details

<sup>1</sup>Department of Anesthesiology and Critical Care Medicine, The Children's Hospital of Philadelphia, University of Pennsylvania, 3401 Civic Center Boulevard, Philadelphia, PA 19104, USA. <sup>2</sup>Department of Pediatrics, University of Utah, Salt Lake City, UT, USA. <sup>3</sup>Department of Pediatrics, Children's Hospital of Michigan, Central Michigan University, Detroit, MI, USA. <sup>4</sup>Department of Pediatrics, Children's National Hospital, George Washington University School of Medicine, Washington, DC, USA. <sup>5</sup>Department of Pediatrics, University of Colorado School of Medicine and Children's Hospital Colorado, Aurora, CO, USA. <sup>6</sup>Department of Critical Care Medicine, UPMC Children's Hospital of Pittsburgh, University of Pittsburgh, Pittsburgh, PA, USA. <sup>7</sup>Department of Pediatrics and Human Development, Michigan State University, Grand Rapids, MI, USA. <sup>8</sup>Department of Pediatrics, Mattel Children's Hospital, University of California Los Angeles, Los Angeles, CA, USA. <sup>9</sup>Department of Pediatrics, Nationwide Children's Hospital, The Ohio State University, Columbus, OH, USA. <sup>10</sup>Department of Pediatrics, Benioff Children's Hospital, University of California, San Francisco, San Francisco, CA, USA. <sup>11</sup>Nemours Cardiac Center, Nemours Children's Health and Thomas Jefferson University, Wilmington, DE, USA. <sup>12</sup>Department of Pediatrics, Washington University School of Medicine, St. Louis, MO, USA. <sup>13</sup>Department of Pediatrics, University of Arkansas for Medical Sciences and Arkansas Children's Hospital, Little Rock, AR, USA. <sup>14</sup>Department of Molecular Biology, Princeton University, Princeton, NJ, USA. <sup>15</sup>Department of Pediatrics, Nemours Children's Health and Thomas Jefferson University, Wilmington, DE, USA.

Received: 21 July 2023 Accepted: 26 September 2023

Published online: 07 October 2023

### References

- Holmberg MJ, Ross CE, Fitzmaurice GM, Chan PS, Duval-Arnould J, Grosse-estreuere AV, et al. Annual incidence of adult and pediatric in-hospital cardiac arrest in the United States. *Circ Cardiovasc Qual Outcomes*. 2019;12(7):e005580.
- Holmberg MJ, Wiberg S, Ross CE, Kleinman M, Hoeyer-Nielsen AK, Donnino MW, et al. Trends in survival after pediatric in-hospital cardiac arrest in the United States. *Circulation*. 2019;140(17):1398–408.
- Topjian AA, De Caen A, Wainwright MS, Abella BS, Abend NS, Atkins DL, et al. Pediatric post-cardiac arrest care: a scientific statement from the American Heart Association. *Circulation*. 2019;140(6):e194–233.
- Laverriere EK, Polansky M, French B, Nadkarni VM, Berg RA, Topjian AA. Association of duration of hypotension with survival after pediatric cardiac arrest. *Pediatr Crit Care Med*. 2020;21(2):143–9.
- Rosner B, Cook N, Portman R, Daniels S, Falkner B. Determination of blood pressure percentiles in normal-weight children: some methodological issues. *Am J Epidemiol*. 2008;167(6):653–66.

6. Kilgannon JH, Roberts BW, Reihl LR, Chansky ME, Jones AE, Dellinger RP, et al. Early arterial hypotension is common in the post-cardiac arrest syndrome and associated with increased in-hospital mortality. *Resuscitation*. 2008;79(3):410–6.
7. Trzeciak S, Jones AE, Kilgannon JH, Milcarek B, Hunter K, Shapiro NI, et al. Significance of arterial hypotension after resuscitation from cardiac arrest\*. *Critical care medicine*. 2009;37(11).
8. Kaji AH, Hanif AM, Thomas JL, Niemann JT. Out-of-hospital cardiac arrest: Early in-hospital hypotension versus out-of-hospital factors in predicting in-hospital mortality among those surviving to hospital admission. *Resuscitation*. 2011;82(10):1314–7.
9. Kilgannon JH, Roberts BW, Jones AE, Mittal N, Cohen E, Mitchell J, et al. Arterial blood pressure and neurologic outcome after resuscitation from cardiac arrest\*. *Crit Care Med*. 2014;42(9):2083–91.
10. Chiu YK, Lui CT, Tsui KL. Impact of hypotension after return of spontaneous circulation on survival in patients of out-of-hospital cardiac arrest. *Am J Emerg Med*. 2018;36(1):79–83.
11. Topjian AA, French B, Sutton RM, Conlon T, Nadkarni VM, Moler FW, et al. Early postresuscitation hypotension is associated with increased mortality following pediatric cardiac arrest. *Crit Care Med*. 2014;42(6):1518–23.
12. Topjian AA, French B, Sutton RM, Conlon T, Nadkarni VM, Moler FW, et al. Early post-resuscitation hypotension is associated with increased mortality following pediatric cardiac arrest. *Crit Care Med*. 2014;42(6):1518.
13. Topjian AA, Telford R, Holubkov R, Nadkarni VM, Berg RA, Dean JM, et al. Association of early postresuscitation hypotension with survival to discharge after targeted temperature management for pediatric out-of-hospital cardiac arrest: secondary analysis of a randomized clinical trial. *JAMA Pediatr*. 2018;172(2):143–53.
14. Topjian AA, Telford R, Holubkov R, Nadkarni VM, Berg RA, Dean JM, et al. The association of early post-resuscitation hypotension with discharge survival following targeted temperature management for pediatric in-hospital cardiac arrest. *Resuscitation*. 2019;141:24–34.
15. Maconochie IK, Aickin R, Hazinski MF, Atkins DL, Bingham R, Couto TB, et al. Pediatric life support: 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Circulation*. 2020;142(16\_suppl\_1):S140–84.
16. Beylin ME, Perman SM, Abella BS, Leary M, Shofer FS, Grossestreuer AV, et al. Higher mean arterial pressure with or without vasoactive agents is associated with increased survival and better neurological outcomes in comatose survivors of cardiac arrest. *Intensive Care Med*. 2013;39(11):1981–8.
17. Ameloot K, Jakkula P, Hästbacka J, Reinikainen M, Pettilä V, Loisa P, et al. Optimum blood pressure in patients with shock after acute myocardial infarction and cardiac arrest. *J Am Coll Cardiol*. 2020;76(7):812–24.
18. Reeder RW, Girling A, Wolfe H, Holubkov R, Berg RA, Naim MY, et al. Improving outcomes after pediatric cardiac arrest - the ICU-Resuscitation Project: study protocol for a randomized controlled trial. *Trials*. 2018;19(1):213.
19. Sutton RM, Wolfe HA, Reeder RW, Ahmed T, Bishop R, Bochkoris M, et al. Effect of physiologic point-of-care cardiopulmonary resuscitation training on survival with favorable neurologic outcome in cardiac arrest in pediatric ICUs: a randomized clinical trial. *JAMA*. 2022;327(10):934–45.
20. Nolan JP, Berg RA, Andersen LW, Bhanji F, Chan PS, Donnino MW, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update of the Utstein resuscitation registry template for in-hospital cardiac arrest: a consensus report from a task force of the international Liaison committee on resuscitation (American heart association, European resuscitation Council, Australian and New Zealand Council on resuscitation, heart and stroke foundation of Canada, InterAmerican heart foundation, resuscitation Council of southern africa, resuscitation Council of asia). *Circulation*. 2019;140(18):e746–57.
21. Pollack MM, Patel KM, Ruttimann UE. PRISM III: an updated Pediatric Risk of Mortality score. *Crit Care Med*. 1996;24(5):743–52.
22. Gaies MG, Gurney JG, Yen AH, Napoli ML, Gajarski RJ, Ohye RG, et al. Vasoactive-inotropic score as a predictor of morbidity and mortality in infants after cardiopulmonary bypass. *Pediatric Crit Care Med: J Soc Crit Care Med World Fed Pediatric Intens Crit Care Soc*. 2010;11(2):234–8.
23. McIntosh AM, Tong S, Deakyn SJ, Davidson JA, Scott HF. Validation of the vasoactive-inotropic score in pediatric sepsis. *Pediatric Crit Care Med: J Soc Crit Care Med World Fed Pediatric Intens Crit Care Soc*. 2017;18(8):750–7.
24. Berg RA, Sutton RM, Reeder RW, Berger JT, Newth CJ, Carcillo JA, et al. Association between diastolic blood pressure during pediatric in-hospital cardiopulmonary resuscitation and survival. *Circulation*. 2018;137(17):1784–95.
25. Berg RA, Morgan RW, Reeder RW, Ahmed T, Bell MJ, Bishop R, et al. Diastolic blood pressure threshold during pediatric cardiopulmonary resuscitation and survival outcomes: a multicenter validation study. *Crit Care Med*. 2023;51(1):91–102.
26. Lederer DJ, Bell SC, Branson RD, Chalmers JD, Marshall R, Maslove DM, et al. Control of Confounding and Reporting of Results in Causal Inference Studies. Guidance for Authors from Editors of Respiratory, Sleep, and Critical Care Journals. *Ann Am Thorac Soc*. 2019;16(1):22–8.
27. Cashen K, Reeder RW, Ahmed T, Bell MJ, Berg RA, Burns C, et al. Sodium bicarbonate use during pediatric cardiopulmonary resuscitation: a secondary analysis of the ICU-RESUSCitation project trial. *Pediatr Crit Care Med*. 2022;23(10):784–92.
28. Morgan RW, Reeder RW, Ahmed T, Bell MJ, Berger JT, Bishop R, et al. Outcomes and characteristics of cardiac arrest in children with pulmonary hypertension: a secondary analysis of the ICU-RESUS Clinical Trial. *Resuscitation*. 2023:109897.
29. Morgan RW, Berg RA, Reeder RW, Carpenter TC, Franzone D, Frazier AH, et al. The physiologic response to epinephrine and pediatric cardiopulmonary resuscitation outcomes. *Crit Care*. 2023;27(1):105.
30. Checchia PA, Sehra R, Moynihan J, Daher N, Tang W, Weil MH. Myocardial injury in children following resuscitation after cardiac arrest. *Resuscitation*. 2003;57(2):131–7.
31. Conlon TW, Falkensammer CB, Hammond RS, Nadkarni VM, Berg RA, Topjian AA. Association of left ventricular systolic function and vasopressor support with survival following pediatric out-of-hospital cardiac arrest. *Pediatr Crit Care Med: J Soc Crit Care Med World Fed Pediatr Intens Crit Care Soc*. 2015;16(2):146–54.
32. Bro-Jeppesen J, Johansson PI, Kjaergaard J, Wanscher M, Ostrowski SR, Bjerre M, et al. Level of systemic inflammation and endothelial injury is associated with cardiovascular dysfunction and vasopressor support in post-cardiac arrest patients. *Resuscitation*. 2017;121:179–86.
33. Topjian AA, Sutton RM, Reeder RW, Telford R, Meert KL, Yates AR, et al. The association of immediate post cardiac arrest diastolic hypertension and survival following pediatric cardiac arrest. *Resuscitation*. 2019;141:88–95.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.