

Table 1. Hippocampal subfield volume change associated with different medication treatment

| | All | |
|--------------------------|------------|----------|
| | F | p |
| Left Hippocampus | | |
| Tail | 0.603 | 0.439 |
| Subiculum | 4.770E-04 | 0.983 |
| CA1 | 8.740E-04 | 0.976 |
| Fissure | 0.144 | 0.705 |
| Presubiculum | 0.225 | 0.636 |
| Parasubiculum | 1.519 | 0.220 |
| Molecular layer | 0.022 | 0.884 |
| GC-ML-DG | 0.0131 | 0.909 |
| CA3 | 0.143 | 0.706 |
| CA4 | 0.002 | 0.963 |
| Fimbria | 0.011 | 0.917 |
| HATA | 0.0079 | 0.929 |
| Whole | 0.0338 | 0.854 |
| Right Hippocampus | | |
| Tail | 3.101 | 0.081 |
| Subiculum | 0.279 | 0.598 |
| CA1 | 0.257 | 0.613 |
| Fissure | 0.258 | 0.612 |
| Presubiculum | 0.002 | 0.962 |
| Parasubiculum | 0.015 | 0.904 |
| Molecular layer | 0.013 | 0.911 |
| GC-ML-DG | 1.050E-04 | 0.992 |
| CA3 | 0.145 | 0.704 |
| CA4 | 0.004 | 0.951 |
| Fimbria | 0.198 | 0.657 |
| HATA | 0.242 | 0.624 |
| Whole | 0.254 | 0.615 |