

		Attendings (N=15)	Residents (N=15)	Medical Students (N=33)	Nursing Students (N=28)
E-cigarettes are not approved by the FDA for smoking cessation	Correct	15 (100%)	15 (100%)	28 (84.8%)	22 (78.6%)
E-cigarettes are the most commonly used tobacco product among US middle and high school students	Correct	5 (33.3%)	8 (53.3%)	16 (48.5%)	16 (57.1%)
Among youth and young adults, the use of e-cigarettes is associated with subsequent initiation of cigarette use	Correct	11 (73.3%)	8 (53.35%)	14 (42.4%)	22 (78.6%)
	Don't know	4 (26.7%)	2 (13.3%)	15 (45.5%)	2 (7.1%)
Nicotine intake from e-cigarettes can be comparable to that from cigarettes	Correct	8 (53.3%)	9 (60%)	18 (54.5%)	17 (60.7%)
	Don't know	4 (26.7%)	4 (26.7%)	5 (15.2%)	5 (17.9%)
Except for nicotine, exposure to potentially toxic substances from e-cigarettes is significantly lower compared with conventional cigarettes	Correct	9 (60.0%)	6 (40.0%)	16 (48.5%)	6 (22.2%)
	Don't know	4 (26.7%)	5 (33.3%)	6 (18.2%)	7 (25.9%)