

Table 3. Hippocampal subfield abnormality in treatment naive MDD patients

| | Control vs Baseline | |
|--------------------------|---------------------|------------------|
| | t | p |
| Left Hippocampus | | |
| Tail | 0.3727 | 0.721 |
| Subiculum | 1.965 | 0.051 |
| CA1 | 2.44 | 0.016 |
| Fissure | 0.597 | 0.551 |
| Presubiculum | 1.662 | 0.098 |
| Parasubiculum | 3.37 | <0.001 |
| Molecular layer | 2.83 | 0.005 |
| GC-ML-DG | 3.21 | 0.002 |
| CA3 | 4.76 | <0.001 |
| CA4 | 3.53 | <0.001 |
| Fimbria | 0.738 | 0.461 |
| HATA | 4.54 | <0.001 |
| Whole | 2.67 | 0.008 |
| Right Hippocampus | | |
| Tail | 0.626 | 0.532 |
| Subiculum | 1.78 | 0.077 |
| CA1 | 2.43 | 0.016 |
| Fissure | -0.452 | 0.652 |
| Presubiculum | 1.022 | 0.308 |
| Parasubiculum | 3.74 | <0.001 |
| Molecular layer | 3.06 | 0.003 |
| GC-ML-DG | 4.07 | <0.001 |
| CA3 | 5.49 | <0.001 |
| CA4 | 4.44 | <0.001 |
| Fimbria | 2.99 | 0.003 |
| HATA | 2.67 | 0.008 |
| Whole | 2.962 | 0.003 |