

Table 1. Hemodynamic Variables

		Baseline	Reperfusion (min)		
			30	60	120
LVDP (mmHg)					
Group	I	112 ± 15	39 ± 7	36 ± 6	29 ± 6
	II	101 ± 13	65 ± 7 ^a	60 ± 8 ^a	52 ± 8 ^a
	III	124 ± 4	63 ± 14 ^a	54 ± 12 ^a	55 ± 10 ^a
	IV	107 ± 11	47 ± 9	44 ± 7	38 ± 6
	V	111 ± 19	37 ± 10	38 ± 9	34 ± 6
	VI	110 ± 16	49 ± 14	48 ± 13	44 ± 11 ^a
LVEDP (mmHg)					
Group	I	10 ± 0	50 ± 10	55 ± 16	59 ± 15
	II	10 ± 0	24 ± 7 ^a	23 ± 4 ^a	23 ± 5 ^a
	III	10 ± 0	43 ± 17	28 ± 2 ^a	28 ± 3 ^a
	IV	10 ± 0	32 ± 8	29 ± 9 ^a	27 ± 10 ^a
	V	10 ± 0	34 ± 15	32 ± 14 ^a	28 ± 14 ^a
	VI	10 ± 0	33 ± 11	34 ± 9 ^a	35 ± 10 ^a
CF (mL/min/g)					
Group	I	21 ± 5	16 ± 5	16 ± 7	15 ± 6
	II	21 ± 6	16 ± 3	15 ± 3	15 ± 4
	III	22 ± 3	14 ± 3	12 ± 3	12 ± 2
	IV	22 ± 5	18 ± 4	15 ± 3	14 ± 3
	V	19 ± 3	14 ± 2	12 ± 2	11 ± 2
	VI	22 ± 4	18 ± 4	16 ± 4	14 ± 3

Table 1: Data are presented as mean ± S.D. LVDP=left ventricular developed pressure; LVEDP=left ventricular end-diastolic pressure; CF=coronary flow ; Group I: control; Group II: sevoflurane; Group III: cyclosporine A; Group IV: control with LY294002; GroupV: sevoflurane with LY294002; GroupVI: cyclosporine A with LY294002. ^a P<0.05 vs. Group I , n=8 for each group.