

12-2010

Fever and Infection in the Neurosurgical Intensive Care Unit

Craig Laws
Craig.Laws@jefferson.edu

Jack Jallo
jack.jallo@jefferson.edu

Follow this and additional works at: <http://jdc.jefferson.edu/jhnj>

[Let us know how access to this document benefits you](#)

Recommended Citation

Laws, Craig and Jallo, Jack (2010) "Fever and Infection in the Neurosurgical Intensive Care Unit," *JHN Journal*: Vol. 5 : Iss. 2 , Article 5. Available at: <http://jdc.jefferson.edu/jhnj/vol5/iss2/5>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in JHN Journal by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Fever and Infection in the Neurosurgical Intensive Care Unit

Craig Laws, MS,¹ Jack I. Jallo, MD, PhD, FACS²

¹Medical Student, Jefferson Medical College, Thomas Jefferson University

²Department of Neurological Surgery, Thomas Jefferson University, Philadelphia, PA

Fever is an adaptive response to a variety of infectious, inflammatory, and foreign stimuli. The “febrile response” confers an immunological advantage to the host over invading microorganisms – bacterial, fungal and viral. Fever results from a cytokine-mediated reaction that results in the generation of acute phase reactants and controlled elevation of core body temperature. The anterior hypothalamus coordinates the “febrile response” in reaction to the release of endogenous pyrogenes and subsequent up-regulation of prostaglandin synthesis. An ensuing change in the hypothalamic set point for temperature regulation advances a synchronized physiologic response from CNS to periphery, on a microscopic and macroscopic level, throughout the entire human organism. This differs from hyperthermia which refers to heat retention attributable to unregulated readjustment of the thermoregulatory mechanism. Clinically, an elevation of core body temperature, whether in fever or hyperthermia, is only the most apparent manifestation of an intricate mechanism that orchestrates activation of autonomic, immunologic, neurologic, hematologic, endocrine and behavioral responses.

Fever has been classically described as having three phases.¹ In the “initiation phase,” cutaneous vasoconstriction diverts blood from the body’s surface to its core. Simultaneously, a broad sympathetic impulse commences rigors (shivering) as opposing muscles groups contract to contribute to excess heat production. The shunting of blood from epithelium to core organs and the increase in metabolic activity leads to the elevation of body temperature. In the “plateau phase,” heat production and heat loss equalize and the body maintains an elevated temperature through rigors and diaphoresis. In the third phase, or “defervescence,” cutaneous vasodilation and diaphoresis lead to dissipation and loss of excess heat, thus allowing the body temperature to drop back to normal.

Hyperthermia occurs when there is disturbances to the central mechanisms of thermoregulation and heat dissipating mechanisms have been compromised. Although the exact mechanisms are unknown, “hyperthermia-induced brain injury” has been speculated to cause direct damage to thermoregulatory centers thus obliterating feedback to the temperature set point elevation.² Some have hypothesized that brain injury induces disruption of mesencephalic-diencephalic physiology responsible for tonic inhibition of thermogenesis and the overriding effects of sympathetic activation which impair peripheral mechanisms of heat dissipation.³ Contrary to the normal processes associated with infectious fever, hyperthermia presents as an elevated body temperature in association with the absence of diaphoresis, tachypnea, and tachycardia.⁴ Other characteristics distinguishing hyperthermia from fever include: an inordinately high temperature (greater than 41 C) that may persist for days to weeks, a lack of diurnal variation, a plateau-like pattern of elevations and maintenance.⁴ Interestingly, appropriate treatment hinges upon correct diagnosis of fever versus hyperthermia, as elevated temperature with hyperthermia is lowered more effectively

by external cooling than by antipyretic agents. Although the causes of this phenomenon remain speculative, the neurologic effects of fever with traumatic brain injury, intracerebral hemorrhage, subarachnoid hemorrhage and post-operative recovery are significant – increased temperature has been associated with increased local cytokine activity, increased infarct size, increased intracranial pressure, vasospasm, symptomatic vasospasm, ischemic brain injury and poor outcomes in the acute phase of neurologic injury.

Infectious Fever versus Non-Infectious Fever

In the NICU setting, fevers are classified as “infectious” or “non-infectious”.⁵ Clinicians further characterize them as either “explained” or “unexplained” according to clinical findings, laboratory tests, or imaging studies that elucidate their etiology.⁶ For example, pneumonia is an “explained” and “infectious” cause of fever, confirmed by leukocytosis, purulent tracheal secretion, positive sputum cultures and infiltrate on chest radiograph. On the other hand, atelectasis is an “explained” but “non-infectious” cause of fever confirmed by characteristic findings in two or more lung segments without leukocytosis or positive sputum cultures. Further still, central neurogenic fever or post-traumatic hyperthermia is an “unexplained” and “non-infectious” cause of fever without confirmatory testing or sufficiently descriptive imaging studies. Central fever is speculated to result from damage to hypothalamus, midbrain, or pons enhanced by increased sympathetic activity, opening of the ventricles, damage to the frontal lobes, physical distortion, diffuse axonal injury, or toxic blood metabolites.⁸

Implication of Fever, both Infectious and Non-Infectious, in the NICU

In the critically-ill neurologic or neurosurgical patient, fever is a double-edged sword, characterized by both beneficial and detrimental effects on the acutely injured central nervous system. The beneficial effects include enhanced resistance to infection, local activation of the coagulation cascade, cytokine-mediated T-cell activation, as well as, neutrophil and macrophage recruitment to injured tissues. In the critically-ill patient, the protective effect of fever

Medical Subject Headings from the United States National Library of Medicine	
Fever	An abnormal elevation of body temperature, usually as a result of a pathologic process.
Hyperthermia, induced	Abnormally high temperature intentionally induced in living things regionally or whole body. It is most often induced by radiation (heat waves, infra-red), ultrasound, or drugs.
Malignant hyperthermia	Rapid and excessive rise of temperature accompanied by muscular rigidity following general anesthesia

against the development of infection generally offset the dangers. Unfortunately, the patient that has suffered an acute insult to the CNS is an exception. After the initial injury, secondary brain insults are speculated to be caused by several processes including mitochondrial dysfunction, inflammatory response, free radical generation, and excitatory neurotransmitter release.⁹ These episodes of secondary brain injury appear to be caused by or temporally associated with hypotension, intermittent hypoxia and ischemia, fever and hyperthermia, cerebral hypertension and elevated intracranial pressure.⁹

In terms of trauma to the CNS, it is commonly accepted that elevated temperature exacerbates secondary brain injury. Fever has been demonstrated to increase glutamate release, provoke diffuse oxygen free radical production, increase cytoskeletal protein degradation, and markedly augment the permeability of the blood-brain barrier.¹⁰ Increased permeability allows activated immune cells to acquire increased access to the interstitial space. Furthermore, vasogenic edema leads to increased fluid in the extracellular space and accumulation of intracellular fluid resulting in cytotoxic edema. Despite evidence suggesting that hyperthermia does not impact the partial pressure of oxygen¹¹ or a causal relationship between elevated brain temperature and increased intracranial pressure,¹² current therapy suggests otherwise. Fever exacerbates ischemic neuronal damage and physiologic dysfunction after acute traumatic brain injury, subarachnoid hemorrhage, and major neurosurgery.¹³ In addition to the aforementioned effects, fever also increases circulating catecholamines boosting cardiovascular output to meet heightened metabolic demands, further taxing the already damaged CNS. In sum, the physiologic response to fever in the CNS has been assumed to contribute deleterious effects that exceed any potential benefits of staving off infection.

Relevance to the Neurosurgical Patient

At least half of febrile incidents in the ICU are of non-infectious origin⁵ In the neurocritical and neurosurgical patients the incidence and prevalence of non-infectious fever are even greater. Febrile episodes occur in roughly 50% of patients in the NICU with neurocritical patients having a slightly lower frequency of fever, approximately 23%,⁶ than patients in the neurosurgical ICU, approximately 47%.¹⁴

Fever in the neurocritical and neurosurgical patient population predominates with vascular injuries, such as intracerebral hemorrhage and

subarachnoid hemorrhage. The highest rates of febrile episode occur in patients with subarachnoid hemorrhage (65%), followed by traumatic brain injury (40%) and intracranial hemorrhage (31%); no cause of fever was identified in 28% of patients, suggesting fever of central origin.⁶

Fever has been linked to an increase in the length of stay, subsequent onset of further febrile episodes. Multiple studies have demonstrated that “non-infectious fever” is common in both the neurocritical and neurosurgical ICU patient population. Fever also tends to predominate amongst patients that have suffered either intracranial injury or traumatic brain injury as compared to spinal disorders.

Infectious Fever in the NICU

A new fever in an ICU patient can have a multitude of infectious and non-infectious etiologies. The practice guidelines from the Task Force of the Society for Critical Care Medicine (SSCM) and Infectious Disease Society of America (IDSA) urge for a “careful clinical assessment” and “cost-conscious approach” to obtaining a diagnosis through laboratory or radiological tests.¹⁵ The SSCM and IDSA Task Force recommend perpetual reflection on the impact of diagnostic testing and radiological imaging with respect to patient harm, further nosocomial infection risks, and the likelihood of legitimate life-sparing intervention. They argue that “infection be considered regardless of temperature but that laboratory tests for infection should be initiated for febrile patients only if clinical assessment indicates a reasonable possibility that infection might be present”.¹⁵

The common causes of infectious fever in a NICU patient are the same as those for any critically ill patient. However, specific considerations should be made for nosocomial infections secondary to depressed level of consciousness and surgical opening of the cranial vault. In a 2003 article, Commichau et al.⁶ reported that fever was associated with infection in 50% of the 387 patients admitted to the Columbia University Hospital NICU. In a European 2002 study, Stochetti et al.¹⁰ reported similar infection rates, again acknowledging pneumonia as the most frequent infection in head-injured patients. The most common infectious causes of fever in the NICU (10–20% of patients in the ICU) are presented in **Table I**.

The high incidence of nosocomial infections (NI) amongst patients admitted to Intensive Care Units (ICUs) are equally relevant to patients admitted to Neurocritical and Neurosurgical ICUs (NICUs). In the neurocritical and neurosurgical patient populations,

nosocomial infection rates also depend on the severity of illness at presentation (i.e. Grade of SAH) and the exposure to invasive devices such as ventilators, central venous catheters and urinary catheters in addition to neuroscience specific devices, such as ventricular catheters. In the U.S., nosocomial infection rates in ICUs are tracked by the “National Nosocomial Infections Surveillance” System (NNIS) run by and published annually by the CDC. Unfortunately, the NNIS does not collect or analyze data specific to the NICU setting, thus there is a paucity of data available on the incidence of NI in the neurocritical or neurosurgical patient population.

In 1997, Dettenkofer et al.¹⁶ began a prospective study in a ten-bed NICU tracking the incidence of nosocomial infections over a 30-month study period. They reported data on 505 patients with a total of 4,873 patient days and 122 NI amongst 96 patients (74 patients with one NI, 18 patients with two NI's, 4 patients with three NI's). The overall incidence of NI in the NICU was 24% (24.2 per 100 patients with a incidence density of 25.0 per 1,000 patient days), with nosocomial pneumonia (11.7%) and urinary tract infections (8.7%) the most frequent cause of infection, respectively. Interestingly, of the 59 nosocomial pneumonias, only 22 (37%) were ventilator-associated while 37 (63%) non-ventilator-associated. Contrary to data on medical-surgical ICUs, more than half the cases of nosocomial pneumonia were non-ventilator associated, which corresponds to decreased ventilator usage among NICU patients. Furthermore, the incidence of nosocomial pneumonias can probably be accounted for by the known high risk of aspiration pneumonia for patients with a depressed level of consciousness. With respect to UTI's, the incidence of NI's were strongly associated with device utilization (42 of 44 UTI's were device-associated). The incidence of bloodstream infections and other catheter related infections (local infections) were relatively low when compared to the incidence of pneumonias and UTI's (1.4% and 1.2%, respectively). Finally, the incidence of ventriculitis and meningitis were also extremely low (0.8% and 0.4%, respectively) when also compared to other NI's.

Non-Infectious Fever in the NICU, Explained and Unexplained Etiology

The patient with classic protracted fever in the NICU setting requires a methodical approach. Given the ramifications, an infectious cause is always presumed and its etiology sought for any critically ill patient with new onset of fever

Table 1

	Indication	Work-Up	Notes
<p>Pneumonia:</p> <p>Ventilator-Assisted Pneumonia: occurs in ~ 25% of mechanically ventilated medical-surgical ICU patients; Pulmonary infections (bronchitis and pneumonia) accounted for 82% of all infections and were the predominant cause of fever in patients with stroke;</p>	<p>leukocytosis, purulent tracheal secretions, purulent sputum, decreased breath sounds, rales on rhonchi, new or worsening infiltrates on chest radiograph</p>	<p>Chest imaging study—chest radiograph and/or CT scan (if otherwise indicated)</p> <p>Cultures of secretions from LRTI before administration of antibiotic</p> <p>Cultures: Expecterated sputum, tracheal secretions, bronchoalveolar lavage</p>	<p>In cases of co-morbid pleural effusion, thoracentesis if sufficient volume or juxtapositioned near lobar consolidation, stain, culture and cytology of pleural fluid</p>
<p>Bloodstream Infections:</p> <p>In absence of IV-line or catheters, bloodstream infections typically arise from gastrointestinal and genitourinary tracts flora.</p>	<p>Any new patient with fever; even when clinical findings do not strongly suggest a non-infectious cause, because positive culture has important prognostic and therapeutic implications</p>	<p>No more than three blood samples (10-15 ml each) for culture need to be drawn during the first 24 hours after fever onset, and two blood samples should generally suffice; draw at least two blood samples from separate venipuncture sites at least ten minutes apart using proper sterile techniques</p>	
<p>Intravenous Catheter-Related Infection:</p> <p>Stable catheters with good vascular access are a mainstay of treatment for the critically-ill. Infections caused by pathogens that have colonized an indwelling vascular device (tunneled, surgically implanted, cuffed central venous, or subcutaneous central venous port) can cause localized or systemic infection</p>	<p>Difficulty drawing or infusing through the catheter site, inflammation at the insertion site; Recovery of microorganisms from the blood at multiple blood cultures; Rapid onset of infection associated with fulminant shock; Multiple positive blood cultures</p>	<p>Again, no more than three blood samples (10-15 ml each) for culture need to be drawn during the first 24 hours after fever onset, and two blood samples should generally suffice; draw at least two blood samples from separate venipuncture sites at least ten minutes apart using proper sterile techniques; infection at the insertion site as evidence by inflammation or purulence at the exit site or an intravascular site (cultures and thrombosis)</p>	
<p>Sinusitis:</p> <p>Occurs in 5% of ICU patients, but with increased frequency in neurosurgical patients secondary to chronic intubation and NG tubes</p>	<p>Cough and leukocytosis with purulent nasal discharge, maxillary facial tenderness, periorbital edema, headache, tooth pain, earache, sorethroat, foul breath, wheezing</p>	<p>Remove nasotracheal or nasogastric tube placement, nasal packing, CT scan, coronal radiographs, or MRI, puncture and aspiration of sinuses under sterile conditions, gram stain and culture of organisms</p>	<p>Risk factors include facial fractures, steroid administration, NG/NT tubes, neurosurgery</p>
<p>Nosocomial Diarrhea:</p> <p>Most common febrile diarrhea in the ICU is <i>Clostridium difficile</i>; Antibiotic administration associated diarrhea (check medical record for previous use of Clindamycin or Cephalosporin or Fluoroquinolones)</p>	<p>2+ stools per day that conform to the container</p>	<p>Remove offending AB agent; diagnosis via “Gold Standard” is tissue culture from stool sample; Elisa Immunoassay test for toxins, Rapid C-Diff Test, gram stain and culture of organisms</p>	<p>Diarrhea is caused by enteral feeding, antibiotic overuse or infectious causes</p>

Table 1 *continued*

	Indication	Work-Up	Notes
Urinary Tract Infections: Urinary tract disease in the presence of bladder catheters, obstruction of the urethra, bladder or ureters, leukocyte esterase dipstick test	Pyuria, leukocytosis, purulent discharge from urethral meatus, obstructive symptoms, retention symptoms, suprapubic tenderness, costovertebral angle tenderness, change in mental status, fluid retention, edema, headache, change in mental status	Urine Culture (clean catch or urine port), remove pre-existing Foley catheter, gram stain or urinary sediment, and culture of organisms	
Surgical Site Infections	purulent discharge, tenderness, dehiscence, erythema, purulence, or tenderness	Examine surgical wound; Remove drains or packing, open the wound, aspirate under sterile conditions, irrigate; Gram stain and culture of organisms from pus or material deep within the wound site	
CNS Infection	Leukocytosis and alterations in neurological parameters, change in mental status, focal findings, absence of neurological findings, lateral shift of midline structures on imaging study, loss of supra-chiasmatic and basilar cisterns, obliteration of the fourth ventricle	CT scan or MRI; Lumbar Puncture, determination of glucose, protein and LDH, Gram stain, white cell count and culture of CSF	Takes a high index of suspicion

in NICU. However, there are equally legitimate noninfectious causes that must be considered as well.

Avoidance of fever and hyperthermia remains a major aim in the management of patients in the NICU. However, this goal is not commonly achieved given the uncertain etiology of a potentially infectious or non-infectious fever. A fundamental principle in this setting is that the management of a classic fever of unknown origin is that “therapy should be withheld, whenever possible, until the cause of fever has been determined, so that it can be tailored to a specific diagnosis”.⁷

Within the non-infectious etiologies, there are four subcategories of non-Infectious fever which are the following:

1. **Non-Infectious Inflammatory Disease:**
Autoimmune Diseases, Rheumatologic Phenomenon,
2. **Neoplastic Disease:**
Primary CNS Tumors and Malignancies, Leukemias and Lymphomas, Metastatic Disease, Oncologic Diseases

3. **Pharmacologic Causes:**

Drug Fever, Withdrawal Syndromes, Neuroleptic Malignant Syndrome, Malignant Hyperthermia, Chemical Meningitis, Drug Toxicities

4. **Post-Traumatic or Peri-Operative:**

Central Fever, Neurogenic Fever, Aseptic Meningitis, Atelectasis or ARDS without Pneumonia, Deep Venous Thrombosis, Posterior Fossa Syndrome

Fever is typical in a variety of acute and chronic neurological disorders confronted in the neurocritical and neurosurgical patient population. Copious experimental evidence suggests that fever is associated with neuronal injury in conditions such as cerebral ischemia, subarachnoid hemorrhage, intracerebral hemorrhage, peri-operative neurosurgical recovery and traumatic brain injury. However, conclusive evidence linking control of non-infectious fever to improved outcomes has inherent difficulties because central neurogenic fever and post-traumatic hyperthermia are diagnoses of exclusion demanding a thorough previous infectious work-up. The impact of fever control in the

NICU is at the mercy of a veteran clinician with a high index of suspicion because traditional methods of fever control are ineffective.

Treatment Dilemma: “To treat or not to treat”

In NICU patients with fever, the clinical sign of pneumonia, bacteremia, sinusitis, UTI, or meningitis or ventriculitis should direct the therapy and choice of an appropriate antibiotic relatively immediately. First, all cultures should be obtained, while all central lines placed for >48 hours and the nasal tube should be removed and cultured using semiquantitative or quantitative cultures. If fever persists for 48–96 hours after antibiotic treatment and without the cause or source of the infection being identified, the patient should be reevaluated for risk factors associated with fungal infection (initiation of empiric antifungal treatment is indicated), while additional diagnostic test, including venography, CBC with differential and eosinophil count, and abdominal imaging are indicated.

During recent years, several biomarkers have been proposed as adjunctive markers for the evaluation of fever, aiming to discriminate

between an infectious cause from a non-infectious reaction or clandestine inflammatory diseases. The biomarkers include: serum procalcitonin assays with variable cutoffs, endotoxin detection systems, triggering receptor expressed on myeloid cells-S (TREM-1), C-reactive protein (CRP), tumor necrosis factor- α , and interleukin-6. From all the above biomarkers, serum procalcitonin assay is approved for early detection of bacterial infection/sepsis during the first day of ICU admission, while the rest have yet to be validated.

The treatment of neurocritical or neurosurgical patients with central neurogenic fever hinges upon the axiomatic decision to declare a fever to be of infectious or non-infectious cause. For instance, to operate on the presumption that a patient has an infection until proven otherwise demands the utilization of rigorous empiric antibiotics regimens. The use of these medications is far from benign. In the ICU setting with all its innate risk factors for nosocomial infections, empiric antibiotic utilization can lead the patient to be unnecessarily exposed to courses of antibiotics with inherent toxicities, drug interactions and side effects that are conceivably detrimental to recovery or elucidation of the fevers origin. Furthermore, a response to antibiotics is an unreliable criterion for diagnosing or excluding bacterial or viral CNS infection, especially in the context of neurocritical or neurosurgical patients with intrinsic risks for non-infectious fever anyway.

Conclusion: Treatment Implications of Infectious versus Non-Infectious Fever

Clinical signs of fever are frequent in neurocritical and neurosurgical patients, and antibiotics are often administered without proof of infection. The clinical characteristics of patients with infectious fever and non-

infectious fever markedly overlap. Classical markers of infection cannot differentiate reliably between inflammation and infection after neurosurgery.¹⁷ The neurosurgical patient having suffered a CNS insult but lacking a documented source of fever, has a central neurogenic fever or post-traumatic hyperthermia by definition. To make the diagnosis of non-infectious unexplained fever (central neurogenic fever or post-traumatic hyperthermia) the clinician must have a high index of suspicion. On the NICU, the clinician combats inherent limitations of the physical exam and neurological exam, while racing against the incubation period for cultures, and exponential growth curve of infectious organisms in vivo. The critical sequelae of either untreated infections or unmanaged central neurogenic fever are urgent dilemmas in the neurocritical and neurosurgical intensive care setting. Regardless, the adverse effects of temperature elevation on neurological recovery are clear. The maxim that “time is tissue” and now “temp is tissue” demands further investigation and further trials of antipyretic medications, anti-hyperthermia strategies such as intravascular cooling techniques in addition to surgical interventions for tissue sparing before the onset of further ischemic damage or coagulative necrosis in the CNS.

References

1. Aiyagari, V and Diringler M. (2007). Fever control and its impact on outcomes: What is the evidence? *Journal of the Neurological Science*, 261, 39–46.
2. Tauber MG and Moser B. (1999). Cytokines and chemokines in meningeal inflammation: Biology and clinical implications. *Clinical Infectious Disease*, 28, 1–11.
3. Rabinstein, AA and Sandhu K. (2006). Non-infectious fever in the neurological intensive care unit: incidence, causes, and predictors. *Journal of Neurology, Neurosurgery and Psychiatry*, 78, 1278–1280.
4. Childers MK, Upright J and Smith DW. (1994). Post-traumatic hyperthermia induced brain injury rehabilitation. *Brain Injury*, 8(4), 335–343.
5. Dimopoulos, G and Falagas ME. (2009). Approach to the febrile patient in the ICU. *Infectious Disease Clinics of North America*, 23, 471–484.
6. Commichau C, Scarmes N and Mayer SA. (2003). Risk factors for fever in the neurological intensive care unit. *Neurology*, 60, 837–841.
7. Bryan CS and Ahuja D. (2007). Fever of Unknown Origin: Is there a role for empiric therapy? *Infectious Disease Clinics of North America*, 21, 1213–1220.
8. Shibata M. (1998). Hyperthermia in brain hemorrhage. *Medical Hypothesis*, 50, 185–190.
9. O'Phelan KH et al. (2009). Patterns of increased intracranial pressure after severe traumatic brain injury. *Neurocritical Care*, 10, 280–286
10. Stochetti N et al. (2002). Pyrexia in head-injured patients admitted to intensive care. *Intensive Care Medicine*, 28, 1555–1562.
11. Spiotta AM, Stiefel MF, Heuer GG, Bloom S, Maloney-Wilensky E, Yang W, Grady MS, LeRoux PD. (2008). Brain hyperthermia after traumatic brain injury does not reduce brain oxygen. *Neurosurgery*, 62: 864–872.
12. Huschak, G, Hoell T, Wiegel M, Hohaus C, Stuttmann R, Meisel HJ and Mast H. (2008). Does brain temperature correlate with intracranial pressure? *Journal of Neurosurgery and Anesthesiology*, 20(2), 105–109.
13. Ginsberg MD and Busto R. (1998). Combating hyperthermia in acute stroke: a significant clinical concern. *Stroke*, 29, 529–534.
14. Kilpatrick MM, Lowry DW, Firlirk AD, Yonas H and Marion DW. (2000). Hyperthermia in the neurosurgical intensive care unit. *Neurosurgery*, 47, 850–856.
15. O'Grady NP et al. (1998). Practice Guidelines for Evaluating New Fever in Critically Ill Adult Patients. *Clinical Infectious Disease*, 26, 1042–1059.
16. Dettenkofer, M, Ebner S, Els T, Babikir R, Lucking C, Pelz K, Rüden H, and Daschner F. (2001). Surveillance of nosocomial infections in a neurology intensive care unit. *Journal of Neurology*, 248, 959–964.
17. Laifer et al. (2005). Dynamics of Serum Procalcitonin in patients after major neurosurgery. *Clinical Microbiology and Infection*, 11(8), 679–681.